

Molecular Characterization and Antibiogram of *Staphylococcus* and *Pseudomonas* species in the Ear and Throat among Patients Attending Rivers State Teaching Hospital

ABSTRACT

Disease of ear, nose and throat (ENT) affect the functioning of adults as well as children, often with significant impairment of the daily life of affected patients. *Staphylococcus* and *Pseudomonas* species are the most prevalent etiologic agent of ENT infections. Therefore the aim of this study was to determine the Antibiogram molecularly characterized the *Staphylococcus* and *Pseudomonas* species in the Ear and Throat among Patients Attending Rivers State Teaching Hospital. Total number of Eighty samples of throat swab, and aural (ear) swab were collected aseptically using sterile Evepon swab sticks and the sample collected were properly labelled with patients' number, date and the side. The swabs were immediately transported with a transportation medium in sterile cotton plugged test tube to the microbiology laboratory for further analysis. Questionnaires were administered for demographic data collections. Each throat and aural swab sample was inoculated onto Mannitol salt agar and Cetrimide agar plates for isolation of *Staphylococcus* and *Pseudomonas* species respectively. The plates were incubated at 37°C and examined for growth after 24–48 hours. Pure culture of the isolates were primarily identified biochemically and confirmed using molecular approaches. Antibiogram of the isolates was performed by the Kirby-Bauer disk diffusion method on Mueller-Hinton agar according to Clinical and Laboratory Standards Institute guideline. Results shown that out of the 50 ear swabs and 30 throat swab samples, 17 (34%) and 8 (26.7%) of the throat swabs ear swabs were positive of *Pseudomonas* spp. While for the *Staphylococcus* spp. 8 (16%) out of the 50 ear swabs and 4 (13.7%) out of 30 throat swabs were positive. The molecular results of the isolates show that *Pseudomonas fluorescens* NR_115715.1 and *Pseudomonas aeruginosa* NR_113599.1, *Staphylococcus aureus* NR_113956.1 were the predominant species isolated from both ear and throat swabs. The results of antibiogram of the *Staphylococcus* and *Pseudomonas* species isolated revealed that commonly used antibiotics especially Ceftazidime, Cefuroxime, Cloxacillin, Gentamicin and Augmentin were 100% resistant to the isolates only Ofloxacin, Ceftriaxone and Nitrofurantoin were susceptible to the isolates with a percentage of 80.2%, 85.7% and 90%. The findings of this study points to the high level of frequent and indiscriminate uses of antibiotics among patient suffering from symptoms of ENT resulting to difficulty in the treatment of infections using common antibiotics. It is strongly recommended that continuous monitoring of patients is required for preparing antibiograms. These susceptibility results will help the physician for deciding the empirical regime for the patients.

Introduction

Ear (Aural), Nose (Nasal), and Throat (ENT) infection are among the most widespread and serious infections that compel an individual to seek medical attention [1]. Disease of ear, nose and throat (ENT) affect the functioning of adults as well as children, often with significant impairment of the daily life of affected patients according to [2]. It has been envisaged that with increase in global population, infection remain the most important causes of disease with upper respiratory infections causing hearing loss and learning disability particularly in children [3]. *Staphylococcus*

aureus is a ubiquitous, and highly adaptive pathogen that colonizes the skin and mucous membrane of the anterior nares, gastro-intestinal tracts, perineum, the genitourinary tracts and pharynx [4]. It is the causative agent of a wide range of infections including Chronic Suppurative Otitis Media (CSOM) which is defined as a prolonged inflammation of the middle ear cleft characterized by a persistent otorrhoea through a perforated tympanic membrane, lasting more than 6-12 weeks in humans and animals with a significant impact on public health [5]. Clinically, *S. aureus* is the most pathogenic member of the genus *staphylococci* and the etiologic agent of a wide variety of diseases that

Comment [SAMUEL EK1]: Molecular characterization and antibiogram of *Staphylococcus* and *Pseudomonas* species isolated from patients attending Rivers State Teaching Hospital.

You can talk about where (ear and throat) it was isolated in your methods.

Comment [SAMUEL EK2]: challenge to...

Comment [SAMUEL EK3]: antibiogram molecularly... is not correct. Maybe you meant antibiogram of/and...

Comment [SAMUEL EK4]: eighty

Comment [SAMUEL EK5]: do you mean eighty samples each of throat swabs and another eighty samples of ear swabs? if not, indicate or say eighty samples including throat and ear swabs...

Comment [SAMUEL EK6]: shows

Comment [SAMUEL EK7]: shows

Comment [SAMUEL EK8]: delete. use comma

Comment [SAMUEL EK9]: and before *Staphylococcus*...

Comment [SAMUEL EK10]: predominant

Comment [SAMUEL EK11]: used

Comment [SAMUEL EK12]: the statement is not correct. The antibiotics cannot be resistant/susceptible to the isolates. it is the isolates that will be resistant/susceptible to the antibiotics.

Comment [SAMUEL EK13]: this is not properly stated. check grammar/spelling

ranges from superficial skin abscess, food poisoning and lifethreatening diseases such bacteremia, necrotic pneumonia in children and endocarditis [4]. The anterior nares of nose are the primary reservoir for replication and spread to other body sites.

Ear infection such as chronic otitis (tinnitus), have serious consequences in developing countries such as retarded language development and progress in school among children [5]. Tinnitus which is now known to be the most common childhood infections, lead annually to the death of over 50,000 children under 5 years (Rover et al., 2006) [6], in other cases nasal condition may be distressing, as in the case of nasal myiasis (Kuruville et al., 2006) [7]. Bacterial species such as *Staphylococcus aureus*, *Streptococcus spp*, *Proteus spp.*, *Haemophilus* and *Coli* forms were found to be responsible for most cases of ENT infections. Kumar et al [6] reported that *pseudomonas*, *Staphylococcus aureus*, *Proteus* and *Klebsiella* are the common bacteria that cause ENT infection in Japura India. In addition according to a study done on ENT infection in Benin City, Nigeria by Osozuwe et al.[7].the following bacteria were found in 466 patients enrolled in the study; *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella spp.*, *Streptococcus pneumoniae*, *E. coli* and *Citrobacter frenndi*. Based on their study, *Pseudomonas aeruginosa* was the most prevalent etiologic agent of ENT infections in Benin City.

P. aeruginosa is a well-known Gram-negative opportunistic human pathogen capable of causing both acute and chronic infections, being highly related to nosocomial infections according to Sharma et al.[8] The lungs are one of the most relevant niches for *P. aeruginosa* colonization, making it one of the most common pathogens isolated from respiratory infections, such as cystic fibrosis (CF) lung infection. Moreover, its great adaptability, phenotypic and genomic plasticity, ubiquity, and opportunistic sense enables *P. aeruginosa* association with other types of infection [9] *Staphylococcus aureus* colonises approximately 30% of the population [1]

with the anterior nares historically considered the most frequent carriage site [2, 3]. Colonised persons carry the bacterium asymptotically and may transmit it to susceptible persons either directly or through contact with fomites. While such colonisation often does not harm the host, it increases the risk of infections in both community and hospital settings [4], hence the need for this study to determine the Antibioqram and molecularly Characterized the *Staphylococcus* and *Pseudomonas* species in the Ear and Throat among Patients Attending Rivers State Teaching Hospital

MATERIALS AND METHODS

2.1 Study Area

Rivers State University Teaching Hospital is known to be a centre for massive influx of patients with one form of complications or another in Rivers State; hence its selection for this research study. Eighty patients who reported to ENT- in- patients unit of this hospital with various signs and symptoms of aural, nasal and throat related infections and diseases were selected randomly for this work. Patient's biodata was as well collected

2.2 Sample Collection

Total number of Eighty samples of throat swab, and aural (ear) swab were collected aseptically using sterile Evepon swab sticks and the sample collected were properly labelled with patients' number, date and the side (i.e. ear, and throat). The swabs were immediately transported with a transportation medium in sterile cotton plugged test tube to the microbiology laboratory for further analysis. Qquestionnaires designed with both open-ended questions such as age and closed ended questions with nominal categorical values such as gender were administered. Data including age, gender and symptoms were also collected [4].

Comment [SAMUEL EK14]: full stop

Comment [SAMUEL EK17]: characterize

Comment [SAMUEL EK18]: Isolated from the ear and...

Comment [SAMUEL EK15]: delete Rover et al., 2006 since you have cited (6).

Comment [SAMUEL EK16]: The

Comment [SAMUEL EK19]: state the exact number of samples for ear swabs and throat swabs

Comment [SAMUEL EK20]: Microbiology laboratory of which school/hospital?



Plate 1: ENT Specialist collecting Throat Swab Sample

2.3 Isolation of *Staphylococcus* and *Pseudomonas* species

Each throat and aural swab sample was inoculated onto Mannitol salt agar and Cetrinide agar plates for isolation of *Staphylococcus* and *Pseudomonas* species respectively. The plates were incubated at 37°C and examined for growth after 24–48 hours. *S. aureus* were initially screened based on the presence of golden yellowish colonies on Mannitol salt agar while *Pseudomonas* species were primarily screened based on green pigmentation.

2.4 Identification of the Isolates

Pure culture of the isolates were primarily identified biochemically and confirmed using molecular approaches.

2.4.1 Bacterial genomic DNA extraction

Five milliliters of an overnight broth culture of the bacterial isolate in Luria Bertani (LB) was spun at 14000rpm for 3 min. The cells were re-suspended in 500ul of normal saline and heated at 95°C for 20 min. The heated bacterial suspension was cooled on ice and spun for 3 min at 14000rpm. The supernatant containing the DNA was

transferred to a 1.5ml microcentrifuge tube and stored at -20°C for other downstream reactions.

2.4.2 DNA quantification

The extracted genomic DNA was quantified using the Nanodrop 1000 spectrophotometer. The software of the equipment was launched by double clicking on the Nanodrop icon. The equipment was initialized with 2 ul of sterile distilled water and blanked using normal saline. Two microlitre of the extracted DNA was loaded onto the lower pedestal, the upper pedestal was brought down to contact the extracted DNA on the lower pedestal. The DNA concentration was measured by clicking on the “measure” button.

2.4.3 16S rRNA Amplification

The 16s rRNA region of the rRNA genes of the isolates were amplified using the 27F: 5'-AGAGTTTGATCMTGGCTCAG-3' and 1492R: 5'-CGGTTACCTTGTTACGACTT-3' primers on an ABI 9700 Applied Biosystems thermal cycler at a final volume of 30 microlitres for 35 cycles. The PCR mix included: the X2 Dream Taq Master mix supplied by Inqaba, South Africa (Taq polymerase, DNTPs, MgCl), the primers at a concentration of 0.4M and the extracted DNA as template. The PCR conditions were as follows: Initial denaturation, 95°C for 5 minutes; denaturation, 95°C for 30 seconds; annealing, 52°C for 30 seconds; extension, 72°C for 30 seconds for 35 cycles and final extension, 72°C for 5 minutes. The product was resolved on a 1% agarose gel at 120V for 15 minutes and visualized on a UV transilluminator.

2.4.4 Sequencing

Sequencing was done using the BigDye Terminator kit on a 3510 ABI sequencer by Inqaba Biotechnological, Pretoria South Africa. The sequencing was done at a final

Comment [SAMUEL EK21]: procedure for the swabs if possible

Comment [SAMUEL EK22]: list the biochemical tests conducted

Comment [SAMUEL EK23]: minutes

volume of 10ul, the components included 0.25 ul BigDye® terminator v1.1/v3.1, 2.25ul of 5 x BigDye sequencing buffer, 10uM Primer PCR primer, and 2-10ng PCR template per 100bp. The sequencing conditions were as follows 32 cycles of 96°C for 10s, 55°C for 5s and 60°C for 4min.

2.4.5 Phylogenetic Analysis

Obtained sequences were edited using the bioinformatics algorithm Trace edit, similar sequences were downloaded from the National Center for Biotechnology Information (NCBI) data base using BLASTN. These sequences were aligned using ClustalX. The evolutionary history was inferred using the Neighbor-Joining method in MEGA X (Saitou and Nei, 1987). The bootstrap consensus tree inferred from 500 replicates (Felsenstein, 1985) is taken to represent the evolutionary history of the taxa analyzed. The evolutionary distances were computed using the Jukes-Cantor method (Jukes and Cantor 1969).

2.5 Antimicrobial Susceptibility Testing

Isolates confirmed were subjected to antimicrobial susceptibility testing to 15 antibiotics. Profiling was performed by the Kirby-Bauer disk diffusion method on Mueller-Hinton agar according to Clinical and Laboratory Standards Institute guidelines [9]. An inoculum for each isolate was prepared by emulsifying colonies from an overnight pure culture in sterile normal saline (0.85%) in test tubes with the turbidity adjusted to 0.5 McFarland standard (0.5 mL of 1% w/v BaCl₂ and 99.5 mL of 1% v/v H₂SO₄), equivalent to 1.0 × 10⁸ cfu/mL. The bacterial suspension was uniformly streaked on Mueller Hinton agar plates using sterile swabs and left for 3 min prior to introduction of the antibiotics. Antibiotics commonly used for treatment

of *S. aureus* and *Pseudomonas* species infections were selected for this assay, Plates were incubated at 35 °C for 24 h, and the diameters of zone of inhibition were measured and results interpreted according to Clinical Laboratory Standards institute

2.6 Calculation of Multiple Antibiotic Resistances Index (MAR)

Multiple antibiotic resistance (MAR) index was calculated using the formula, a/b (where, a=number of antibiotics to which the organism was resistant and b = total number of antibiotics to which the organism was tested [18])

2.7 Statistical Analysis

Statistical analysis was carried out on the data obtained during the study using a computer-based program SPSS version 20 for Analysis of Variance (ANOVA)

3. Results and Discussion

A total number of 50 ear swabs and 30 throat swabs were collected from the patients attending Rivers State University Teaching Hospital. Out of these, ear swabs were positive of *Pseudomonas* spp. 17 (34%) and 8 (26.7%) in throat swabs. While *Staphylococcus* spp. 8 (16%) out of the 50 ear swabs and 4 (13.7%) out of 30 throat swabs were positive (Figure 1). Female patients recorded the percentage of positive for both *Staphylococcus* spp and *Pseudomonas* spp. In ear and throat swabs (Table 1 and 2) respectively. The molecular results of the isolates show that *Pseudomonas fluorescens* which has NCBI accession number NR_115715.1 and *Pseudomonas aeruginosa* which has NCBI accession number NR_113599.1 were the predominates species isolated while *Staphylococcus aureus* which has NCBI accession number NR_113956.1 and *Staphylococcus aureus*

Comment [SAMUEL EK26]: hour

Comment [SAMUEL EK24]: maintain the appropriate reference format

Comment [SAMUEL EK25]: use appropriate style

Comment [SAMUEL EK27]: this is not clear.

Comment [SAMUEL EK28]: continue the statement

which has NCBI accession number NR_037007.2.

Table 1: Prevalence of *Pseudomonas* species Based on Sex and sample

Sample	Sex	Negative	Positive	Total sample Examined
Ear Swabs	Male	10 (62.5)	6 (37.5)	16
	Female	23 (67.6)	11 (32.3)	34
Total		33 (66)	17 (34)	50
Throat Swabs	Male	8 (72.7)	3 (27.3)	11
	Female	14 (73.7)	5 (26.3)	19
Total		22 (73.3)	8 (26.7)	30

Comment [SAMUEL EK29]: check your calculations again

Table 2: Prevalence of *Staphylococcus* species Based on Sex and sample

Sample	Sex	Negative	Positive	Total sample Examined
Ear Swabs	Male	13 (81.3)	3 (18.7)	16
	Female	29 (85.3)	5 (14.7)	34
Total		42 (84)	8 (16)	50
Throat Swabs	Male	10 (90.9)	1 (9)	11
	Female	16 (84.2)	3 (15.8)	19
Total		26 (86.7)	4 (13.7)	30




Figure 1 : Percentage Occurrence of *Staphylococcus* and *Pseudomonas* species in the Ear and Throat Swab.




Figure 2: Phylogenetic tree showing the Evolutionary Distance between the Bacterial Isolates

Generally, the most prevalent organisms was *Pseudomonas aeruginosa* (NR_113599.1) while the least was *Staphylococcus aureus* (NR_037007.2.). Nevertheless, both strains of *Staphylococcus aureus* and species of *Pseudomonas* were isolated in both ear and throat swabs.

The finding in this study is in agreement with the report of Kumar *et al.* [23], who find out that *Pseudomonas spp.*, *S. aureus*, *Proteus spp.* and *Klebsiella spp.*, are the common bacteria that cause ENT infection in Japura India, similarly Osazuwa *et al.* (2011) find out that *P. aeruginosa*, *S. aureus*, *Klebsiella spp.*, *S.*

pneumoniae, *E. coli* are the bacteria associated with ENT infection and *Pseudomonas aeruginosa* was the most prevalent aetiological agent of ENT infection in Benin city, El-Mohmood *et al.* [24]. These study collaborate the above findings that *P. aeruginosa*, *Pseudomonas fluorescens* and *Staphylococcus aureus* were found in both Ear, and throat infection.

The percentage of ENT patients attending Rivers State University Teaching Hospital who are currently taking antibiotics before seeking specialist advice is significant high as shown in (Figure 3)

Comment [SAMUEL EK30]: organism

Comment [SAMUEL EK31]: discuss your work in more details before referring to similar or contrary results




Figure 3: Percentage of patients based on History Antibiotic Uses

The results of antibiogram of the *Staphylococcus* and *Pseudomonas* species isolated in this study from ear and throat swab is presented in Figure 4 and 5 respectively. The *Staphylococcus* and *Pseudomonas* species were subjected to eight different types of antibiotics such as Ceftazidime (30µg), Cefuroxime (30µg), Gentamicin (10µg), Cloxacillin (5µg), Ofloxacin (5µg), Augmentin (30µg), Ceftriaxone (30µg), Erythromycin (5µg), and Nitrofurantoin (30µg) but of which only Ofloxacin, Ceftriaxone and Nitrofurantoin were susceptible to the isolates with a

percentage of 80.2%, 85.7% and 90% for the *Pseudomonas* species respectively (Fig 3) while for the *Staphylococcus* species the percentage susceptibility ranged from Erythromycin (8%), Gentamicin (16.7%), Ceftriaxone (50%), Augmentin (60%) to Ofloxacin (90%), as shown in (Figure 4). Other commonly used antibiotics especially Ceftazidime, Cefuroxime, Cloxacillin, Gentamicin and Augmentin were 100% resistant to the isolates. The high percentage resistant recorded in this study could be as a result of overuse and frequent misuse of antibiotics by patients who observed little


Comment [SAMUEL EK33]: restructure your statement to make your point clearer

Comment [SAMUEL EK32]: full stop

symptoms in their ear, nose and throat which have resulted in changes of antibiotic resistance profiles of microorganisms amongst bacterial populations. The high rate of resistance of the isolates to Cefuroxime, Ceftazidime and Augmentin is of concern as most physicians are quick to prescribe these drugs especially Augmentin in the treatment of bacterial infections according to Ogonna and Azuonwu [24]. Resistance of bacteria to Cefuroxime has


been reported by Harrison and Bratcher [25] who investigated the susceptibility of some microorganisms to antibiotics such as cefuroxime. Study carried by Oparaoduet *al* [27] on AntibioGram of Methicillin Resistant *Staphylococcus aureus* isolated from Nasal Carriage of Some University Students of Rivers State University, Port Harcourt, Nigeria also revealed high percentage resistant of commonly uses antibiotic as obtained in this study.

Comment [SAMUEL EK34]: state if the overuse or misuse was during the treatment, before the treatment or after treatment



Key: Ceftazidime (CAZ), Cefuroxime (CRX) Gentamicin (GEN) Cloxacillin (CXM) Ofloxacin (OFL) Augmentin (AUG) Ceftriaxone (CTR) Nitrofurantoin (NIT)

Figure 4 Sensitivity profile of *Pseudomonas* species Isolated from Ear and Throat Swab



Key: Ceftazidime (CAZ), Cefuroxime (CRX) Gentamicin (GEN) Cloxacillin (CXM) Ofloxacin (OFL) Augmentin (AUG) Ceftriaxone (CTR) Erythromycin (ERY)

Figure 5 Sensitivity profile of *Staphylococcus* species Isolated from Ear and Throat Swab

CONCLUSION AND RECOMMENDATIONS

The percentage of *Pseudomonas fluorescens* (NR_115715.1), *Pseudomonas* (NR_113599.1), *Staphylococcus aureus* (NR_113956.1) and *Staphylococcus*

aureus (NR_037007.2.) isolated from ear and throat swabs in this study revealed that *Staphylococcus* and *Pseudomonas* species are the main causative agent of ENT infection. The findings of this study shown low

Comment [SAMUEL EK35]: your study didn't show how they are "main"... you focused on the two bacterial pathogens...

percentage of the *Staphylococcus* and *Pseudomonas* species in the ear and throat of patients as points to the high level of frequent and indiscriminate use of antibiotics which in turn resulted to high level of resistant of common uses antibiotics such as Augmentin, Gentamicin, Erythromycin and Ceftazidime among patient suffering from symptoms of ENT resulting to difficulty in the treatment of infections using the common antibiotics. It is strongly recommended that self medical practices by patient be discouraged. Continuous monitoring of patients is also required for preparing antibiograms. These susceptibility results will help the physician for deciding the empirical regime for the patients.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

REFERENCES

1. Ahmad M.M., Kurawa, Z.M., Shu'aibu, and Yahaya G. Microbiological Assessment of Bacterial Isolates from Ear, Nose And Throat (ENT) Among Patients Attending Aminu Kano Teaching Hospital. *Nigerian Journal of Basic and Applied Science* 2016, 24(1): 15-18
2. Witsell, D.L., Dolor, R.J., Boile, J.N. and Stinnet, S.S.. Exploring health related quality of life in patients with diseases of the ear, nose and throat; A multicenter observation study. *Otolaryngology Head and Neck Survey*, 2001, 125(4): 288-298
3. Chibuiké, I., Reginald, A.O., Solomon, U.C., Ifeanyi A.O., Conrad J., Chinenyenwa, J.N., Nnadozie J. and kelechi U.O. Prevalence and Antibiotic Susceptibility Patterns of Methicillin Resistant *Staphylococcus Aureus* (MRSA) Isolated from Healthy Inhabitants of Uturu Rural Communities, Abia State, Nigeria. *Journal of Natural Science Research*, 2013, 3(10): 85-91
4. Oparaodu U., Odu N. N. and Uzah G. A. Antibigram of Methicillin Resistant *Staphylococcus aureus* isolated from Nasal Carriage of Some University Students of Rivers State University, Port Harcourt, Nigeria, *Journal of Advances in Medicine and Medical Research*. 2021, 33(15): 73-81.
5. Onotai L.O and Oparaodu Ureh. Challenges in the Management of Adult Chronic Suppurative Otitis Media in Port Harcourt Nigeria. *Journal of Dental and Medical Sciences*. 2017, 16 (1), 86-90
6. Ogbonna DN, Azuonwu C. Plasmid profile and antibiotic resistance pattern of bacteria from abattoirs in Port Harcourt City, Nigeria, *International Journal of Pathogen Research*. 2019;2(2):1-11.
7. Odu N, Okonko I. Nasal carriage and antibiotics susceptibility of *Staphylococcus aureus* in healthy students of university of Port Harcourt, Rivers State, Nigeria. *New York Science Journal*. 2012; 5(7):56-63.
8. Chigu CO, Ezeronye OU. Antibiotic resistant *Staphylococcus aureus* in Abia State of Nigeria. *Afr. J. Biotech*. 2013; 2(10):374-378
9. CLSI Clinical and Laboratory Standard Institute Performance standards for Antimicrobial Susceptibility Testing Eighteenth

Comment [SAMUEL EK36]: you must have sought consent of the patients you took the samples from.

informational supplement.
2006;M100-S18,28(1):46- 52

10. Madani TA, Al-Abdulla AA, Al-Sanousi TM, Ghabrah SZ, Afandi, Bajjunid HA. Methicillin resistant *Staphylococcus aureus* in two tertiary care centres in Jeddah, Saudi Arabia. *Infect. Control Hosp. Epidemiol.* 2001;22:211-216.
11. Sharma G, Rao S, Bansal A, Dang S, Gupta S, Gabrani R. 2014. *Pseudomonas aeruginosa* biofilm: potential therapeutic targets. *Biologicals.* 42(1):1-7.
12. Shen K, Sayeed S, Antalis P, Gladitz J, Ahmed A, Dice B, Janto B, Dopico R, Keefe R, Hayes J, et al. 2006. Extensive genomic plasticity in *Pseudomonas aeruginosa* revealed by identification and distribution studies of novel genes among clinical isolates. *Infect Immun.* 74(9):5272-5283

UNDER PEER REVIEW