

1 **INTESTINAL PARASITES IN SCHOOL-AGED CHILDREN OF RUMUODOGO,**  
2 **EMOHUA LOCAL GOVERNMENT AREA, RIVERS STATE, NIGERIA**

3  
4 **ABSTRACT**

5 **Aim:** This epidemiological survey was carried out to assess the prevalence of intestinal parasites  
6 in School aged Children.

7 **Place and Duration of study:** The study was carried out in Rumuodogo Community in Emohua  
8 Local Government area of Rivers State, Nigeria between the months of March to June 2019.

9 **Methodology:** Microscopic examination of stool samples from 200 school-aged pupil between  
10 the ages 3-18 years was carried out using formol-ether concentration and direct wet mount, and  
11 stained iodine. Demographic data as well as epidemiological data was obtained using a  
12 questionnaire. .

13 **Results:** The finding reveals an overall prevalence of 49.5% However, there was / was no  
14 significant difference in the mean value ( $\chi^2 =$  ,  $df =$  ,  $P =$  ) The common intestinal parasites identified  
15 were *Ascaris lumbricoides* 19.5%, *Ancylostoma duodenale* 9.5%, *Trichuris* spp. 6%,  
16 *Strongyloides* spp. 4.5%, *Enterobius* spp. 1%, *Taenia solium*, 1.5%, *Taenia saginata* 1%,  
17 *Entamoeba histolytica* 4.5%, *Giardia lamblia* 4% and *Schistosoma mansoni* 2%. However, there  
18 was / was no significant difference in the mean value ( $\chi^2 =$  ,  $df =$  ,  $P =$  ) The prevalence of  
19 intestinal parasites infections with respect to the available and use of toilet facility revealed that  
20 *Pit toilet users had the highest prevalence of 62 (56.9%)* while no infection was recorded among  
21 users of water closet toilet. The result showed that there was / was no significant difference in the  
22 mean value ( $\chi^2 =$  ,  $df =$  ,  $P =$  ). When age group was compared the result did /did not show  
23 difference in the mean value ( $\chi^2 =$  ,  $df =$  ,  $P =$  ) The finding however, showed that age groups, 6-10  
24 years had the highest prevalence of 55.6%. **Maximum of seven children had multiple infections.**  
25 **The male female ration was 62:37** while the percentage for males was 43.2% and 50% for  
26 females.

27 **Conclusion:** Intestinal parasitic infections are associated with reduced development of children  
28 manifested by a reduced physical fitness which may result from vitamin deficiencies, inducing  
29 intestinal bleeding, and protein energy malnutrition. Therefore, the high prevalence of intestinal  
30 parasites recorded in this study calls for an intensified effort in the control of the infections.  
31 **Regular deworming of children and environmental sanitation should** be carried out to further  
32 reduce the prevalence.

33 **Keywords;** Parasites, Children, School, Rumuodogo

## 37 1.0 INTRODUCTION

38 In spite of the tremendous advances in medicine and clinical Parasitology, globally over the past  
39 few decades, human intestinal parasitic infections remain the single largest cause of human death  
40 and discomfort in school children and poor communities. These infections are the most prevalent  
41 in the tropical and sub-tropical regions of the world, where adequate water and sanitation are  
42 lacking [1]. The major factor that contributes to transmission and prevalence of intestinal  
43 parasitic infection most especially in many endemic areas of the tropics are poor environmental  
44 sanitation, poor hygiene practices, indiscriminate disposal of human and animal waste that  
45 resulted to the contamination of the environment which predisposes humans to infection as a  
46 result of interaction of humans to contaminated environment due to poverty [1,2] . Reports by  
47 the world health organization [3], puts the figure of infected children worldwide at 880 million.  
48 This high infection is due to severe shortage in health care, education, sanitation, transport and  
49 chronic poverty [4].

50 Intestinal parasitic infections could have significant effect on the growth and development of  
51 children which manifests as reduced physical fitness and constrained growth due to problems  
52 such as vitamin deficiencies, inducing intestinal bleeding, and protein energy malnutrition  
53 associated with their effects. There might equally be a subtle but important development effect  
54 on cognition and educational development [4, 5, 6].

55 Intestinal parasitic infection may also pose some serious consequences on human health, such as  
56 hepatomegaly, oesophageal varices and bleeding [7]. Individuals infected with helminths  
57 according to Mulu *et al.* [8], could be susceptible to other infections such as malaria and HIV.

58 Intestinal parasites especially *Ascaris lumbricoides*, *Trichuris trichiura* and hookworm species  
59 are most common in Nigeria. Prevalence of these parasites especially *Ascaris*, according to  
60 Ovutu and Olaniyi [1, 9] has remained unchanged in the last 50 years and poly-parasitism occurs.  
61 Ezey *et al.* [4] reported a 70% prevalence of intestinal helminths in school children of Khana,  
62 Rivers State. Mafiana, [10] equally reported 70.8% prevalence in Ilewo Ogun State Nigeria. This  
63 study was carried out to determine the prevalence of intestinal parasites in school- aged children  
64 in Rumuodogo town Emohua, Rivers State.

## 65 2.0 Materials and Methods

66 **2.1 Study Area**

67 The study was carried out in Rumuodogo community in Emohua Local Government Area of  
68 Rivers State, Nigeria. It is located at latitude 4<sup>0</sup> 53'0 North and longitude 6'52'0 East of the  
69 Greenwich. The area lacks social amenities. The vegetation is rainforest and the humidity is very  
70 high. The community is surrounded by both fresh and saline waters and their major occupations  
71 are subsistence farming and fishing.

72

73

74

75 **2.2 Collection of stool samples**

76 Morning stool samples were collected using sterile vials from 200 pupils with the help of their  
77 parents. The specimens were preserved in 10% formalin and transferred to the Animal and  
78 Environmental Biology laboratory of Rivers State University for analysis.

79 **2.3 Examination of stool samples**

80 **2.3.1 Formol-Ether Concentration Technique**

81 This method was adopted from Cheesbrough [11]. About 1g of the stool was emulsified in about  
82 4ml of 10ml of formol solution in a test tube. The formol solution was prepared by mixing 50%  
83 strong formaldehyde solution with 450ml distilled water. 4 ml of the formol water was added to  
84 the solution and mixed properly by shaking the mixture was filtered into a test tube using a cloth  
85 gauge and about 3-4ml diethyl ether was added and shaken vigorously and allowed to stand for 2  
86 minutes. The mixture was centrifuged at 1000 revolutions per minute for 3 minutes. Using a  
87 glass rod, the faecal debris from the side of the tube was loosened and the test tube inverted to  
88 pour off the supernatant leaving the deposit at the bottom of the test tube in an upright position.  
89 The deposit was mixed by tapping the tube with the finger and using a Pasteur's pipette, a drop  
90 of the deposit was applied on a microscope slide mixed with Lugol's iodine, covered with a

91 cover slip and viewed under the microscope with X10 and X40 objective respectively. The eggs  
92 and larvae of the parasites were identified with reference to Atlas of Parasitology.

### 93 **2.3.2 Direct wet preparations**

94 A little portion of stool was mixed with 2 drops of 0.85% saline solution on a slide. A drop of  
95 iodine was added and examined under the microscope [11].

### 96 **2.4 Data Analysis**

97 Data was entered into Microsoft excel 2010 and analysis done using the Chi-square statistical  
98 tool. Descriptive statistics was calculated and presented in tables [12]

### 99 **3.0 Results**

100 **Table 1.** Age related Incidence of intestinal parasites in School-Aged Children in the study area

Age groups in years	No. Examined (%)	No. Positive (%)	No. Negative (%)
3-6	62 (31)	35 (56.4)	27(43.5)
7-10	72 (36)	40 (55.5)	32 (54.0)
11-14	37 (18.5)	17 (45.9)	20 (54.1)
15-18	29 (14.5)	7 (24.1)	22 (75.9)
Total	200	99 (49.5)	101 (50.5)

101

102 Table 1 shows 99 (49.5%), prevalence of intestinal parasites in the study area, with children  
103 between ages 3-6 having the highest prevalence of 56.4% while ages 15-18 recorded the least.

104

105

106

107 **Table 2. Sex related incidence of intestinal parasites amongst school-Aged Children in the**  
 108 **area.**

109

Sex	No. Examined (%)	No. Positive (%)	No. Negative (%)
Male	125 (62)	54 (43.2)	71 (56.8)
Female	75 (37.5)	45 (60)	30 (40)
Total	200	99 (49.5)	101 (50.5)

110 Table 2 shows a the prevalence of intestinal parasites with respect to sex. The result shows that higher  
 111 prevalence 125 (62:37). Than their female counterpart which had 75(37.5).

112 **Table 3: Incidence of intestinal parasites species in relation to age groups**

113

Species of Parasite	Age groups (Years)				Total
	3-6(%)	7-10(%)	11-14(%)	15-18(%)	
<i>Ascaris lumbricoides</i>	18 (9)	13 (6.5)	4 (2)	4 (2)	39 (19.5)
<i>Hookworm sp.</i>	7 (3.5)	6 (30)	4 (2)	2 (1)	19 (9.5)
<i>Trichuris trichiura</i>	3 (1.5)	5 (2.5)	3 (1.5)	1 (1.5)	12 (6.0)
<i>Strongyloides stercoralis</i>	2 (1)	4 (2)	3 (1.5)	0	9 (4.5)
<i>Enterobius vermicularis</i>	2 (10)	0	0	0	2 (1)
<i>Schistosoma mansoni</i>	2 (1)	1 (0.5)	1 (0.5)	0	4 (2)
<i>Taenia sp</i>	2 (1)	3 (1.5)	0	0	5 (2.5)
<i>Entamoeba histolytica</i>	2 (1)	5 (2.5)	0	2 (1)	9 (4.5)
<i>Giardia lamblia</i>	4 (2)	4 (2)	0	0	8 (4)
Total (200)	42 (21)	41 (20.5)	15 (7.5)	9 (4.5)	107 (53.5)

114 Table 3 shows that *Ascaris lumbricoides* had the highest prevalence of 19.5% while *Shistosoma mansoni*  
115 and *Enterobius vermicularis* had the least prevalence in the study area respectively.

116

117

118 **Table 4: Incidence of intestinal parasites with respect to dominant toilet facility in the study area.**

Toilet Facility	(%) No. Examined	(%) No. Positive	(%) No. Negative
Bush	88	36 (40.9)	52 (59.0)
Pit	109	62 (56.9)	47 (43.1)
Water closet	3	0	3 (100)
Total	200	98 (49)	102 (51)

119

120 Table 4 indicates pit toilet users had the highest prevalence of 56.9% than those who used bush as well  
121 as water closet that didn't record any incidence.

122

#### 123 **4.0 Discussion**

124 There has not been any report of work on the prevalence of intestinal parasites among school-  
125 aged children in the study area. From the results of this study, high prevalence (49.5%) observed  
126 possibly may be due to poor hygiene condition of the area and thus the various intestinal  
127 parasites recorded

128 The prevalence of 49.5 recorded in this study was lower when compared with 66.3% observed  
129 by Agi [13] in Choba community, 70% by Ezey *et al.* [4] in Sii, Gwara and Gure communities in  
130 Khana Local Government Area in Rivers State and 72% finding by [14] in University of Guyana,  
131 Georgetown, Guyana.

132 *Ascaris lumbricoides* was the most prevalent parasite encountered in the study followed by  
133 hookworm and *Trichuris* species. This could be attributed to the involvement of the people in  
134 agricultural practices of farming and fishing or lack of adequate personal hygiene. Hookworm  
135 infection maybe as a result of walking or playing on infested soils bare footed.

136 Findings are equally in line with the report of Michael *et al.*[15] who found a high prevalence of  
137 *Ascaris lumbricoides*, among communities in the vicinity of Port Harcourt. Also, Abah and  
138 Arene [16] reported that *Ascaris lumbricoides* occurred more frequently (51.78%) among  
139 primary school-aged children in Rivers State.

140 The data shows that females had a higher prevalence of 60% which may be as a result of being  
141 involved in domestic and agricultural activities which predisposes them to polluted  
142 environments. The male work in farms, far from homes and are less exposed to heavily polluted  
143 home environment and consequently less infected. This is consistent with [17].

144 In the case of the most toilet facility (pit) used, most people in the area are poor and unable to  
145 afford water closet. Table 4 shows that the pit latrine users had a comparative high prevalence of  
146 (56.9%) than bush toilet system. The result is in agreement with [18] who observed a high  
147 percentage of parasitic infection in Benin City among pit toilet users. But the bush users had a  
148 prevalence of (40.9%) which is equally high. This practice is a product of underdevelopment and  
149 shows lack of personal and community hygiene. The use of bush and pit as toilets is a major  
150 source of soil and water pollution which is responsible for the high incidence of soil transmitted  
151 helminth parasites in Nigeria [1,4].

152 The high prevalence (55.5%) of the parasites among aged groups 6-10 years may be as result of  
153 the children in this age group walking barefooted around homes even when going to school and  
154 swim in contaminated stream. The infection decreased in ages 11-14 and ages 15-18. This is  
155 consistent with Mafiana [10] who noted 82.1% prevalence in children aged 10-11 years. This  
156 sequence may be attributed to the fact that children in older age groups had behavioural patterns  
157 different from those of the younger groups; more knowledge of disease, and a little improvement  
158 on personal hygiene.

159 *Gairdia lambia* and *Entamoeba histolytica* had a prevalence of 4 and 4.5% respectively which  
160 was higher when compared to findings by Akaniwo [19] who recorded a 2.5% prevalence in  
161 Obio/Akpor but lower when compared to 11.7% observed by Mercado *et al.* [20] in Chile. Poly-  
162 parasitism was observed in 7 out of the 200 faecal samples. The co-existence of the different  
163 parasites in the infected individuals is an important feature in the biology of these parasites. This

164 is so because the acquisition of single infection produces a different effect from acquisition of  
165 large number of worms. .

166

## 167 5.0 CONCLUSION

168 It was observed from this study that majority of those infected were not necessarily because of  
169 poverty but because of ignorance as a result of lack of education and good environmental  
170 sanitation (through proper waste disposal system), poor personal hygiene and shortage of  
171 drinking water. Therefore, health facilities, pipe borne water, education through enlightenment  
172 programme and improved sanitary conditions be provided since the infection is by no means than  
173 water and poor sanitary conditions.

### 174 **Ethical Approval and Consent:**

175 Approval for the study was given by the school authorities and written consent was obtained  
176 from the parents and guardians of the children

177

178

179

## 180 REFERENCES

- 181 1. Ovutor, O. , Imafidor H., and Awi-waadu G.D.B (2017). Assessment of Physico-  
182 chemical parameters of soils in fallowing farmlands on the abundance of human infecting  
183 geohelminthes in Mgbuitanwo Emohua, Rivers state. *Journal of Advances in Biology and*  
184 *Biotechnology*, 15(3): 1-8,
- 185 2. Udonsi, J. K., Benke, J M and Gilbert, F.S (1996). Analysis of the prevalence of infection  
186 and association between human gastrointestinal nematodes among different age classes  
187 living in urban and suburban communities of Port Harcourt. *J. Helminthol* 70 pp. 75- 84
- 188 3. World Health Organization (2013). Soil transmitted fact sheets. 2010 report, WHO press,  
189 Geneva.107:16-171

- 190 4. Eze, C.N. , Owhoeli, O. and Ganale , S.S (2016) Assessment of intestinal helminthes in  
 191 community school children of Khana Local Government Area, Rivers State, Nigeria.  
 192 *Nigerian Journal of Parasitology* ;37 (1):pp
- 193 5. Abah, A E and Temple, B. (2015) Prevalence of Malaria Parasite among Asymptomatic  
 194 Primary School Children in Angiama Community, Bayelsa State, Nigeria. *Trop Med*  
 195 *Surg*;4(1):203
- 196 6. Odu, N.N. Okonko, I. O. and Ethi, O. (2011). Study of neglected tropical diseases  
 197 (NTD)Gastrointestinal helminthes among school children in Port Harcourt, Rivers State  
 198 Nigeria. *Report and Opinion* 3 (9): 6-16
- 199
- 200 7. Alum A, Rubino J R,and Ijaz M K(2010) The global war against intestinal parasites  
 201 should be use in a holistic approach? *Int. J infected disease.* 2010; 14:732-730.  
 202
- 203 8. Mulu A., Maier M.,and Liebert U G (2013) Deworming of intestinal helminthes reduces  
 204 HIV- 1subtype C viremia in chronically coinfectd individuals. *International journal of*  
 205 *infectious Diseases*; 10(6):02-016.  
 206
- 207 9. Olaniyi J E, Muktar H A, Pauline E J (2007) A review of intestinal helminthiasis in  
 208 Nigeria and the need for school-based intervention. *Journal of Rural and Tropical Public*  
 209 *Health*;6:33-39  
 210
- 211 10. Mafiana C. F. (1995) Intestinal Helminthiasis (Ascaris) among school children in illewo-  
 212 Orile, Ogun State, Nigeria. *The Nigeria Journal of Parasitology* 16 pp.47-53.
- 213 11. Cheesbrough M. District Laboratory Practice in Tropical Countries. Part 1 (2ed)  
 214 Cambridge University Press 2005: 194-202
- 215 12. Thrushfield, D.M. (1995). Veterinary Epidemiology. Blackwell Science
- 216 13. Agi P. I (1997). Comparative Helminth infection of man in two Rural Communities of  
 217 Niger delta, *Nigeria WAJM* 16 (4) 232-236.
- 218 14. Jonathan W., Abdullah A. A and Kumar S, (2015) Prevalence and Association of  
 219 Parasitic Helminths among the Cross Section of Male and Female Gender Groups at  
 220 University of Guyana, Georgetown, Guyana. *Research Journal of Parasitology*, 10: 50-  
 221 57.
- 222 15. Michael E I., Abah A., Marcus P (2017). Prevalence of gastrointestinal parasitic infection  
 223 among school children in Port Harcourt City Local Government Area, Nigeria. *J*  
 224 *Bacteriol Parasitol* 8:323

225 16. Abah A E., and Arene F.O.A (2015). Status of intestinal parasitic infections among  
226 primary school children in Rivers State, Nigeria, *Journal of Parasitology Research*, vol  
227 2015. Article ID 937096.

228 17. Ephrem T., Tariku B., Seleshi K M., Ahmed Z and Tefera B (2017). Prevalence and  
229 intensity of soil transmitted helminthes among school children of mendera Elementary  
230 school, southwest Ethiopia. *Pan African Medical Journal*; 27:88.

231 18. Mordi R.M and Paul O.A.N. (2007). A study of blood and gastro-intestinal parasites Edo  
232 State. *African Journal of Biotechnology* Vol 6 (19). pp 2201-2207.

233 19. Akaniwo G. (1999) The prevalence of intestinal parasites among school-aged children in  
234 Obio/Akpor Local Government Area Rivers State, Nigeria. Unpublished B.Sc. Zoology  
235 Project.

236 20. Mercado R., Otto J P., Musleh M and Perez M(1997) Human infection by intestinal  
237 protozoa and helminth in Calbuco Country; X Region, Chile. *Bio-cli-parasitol* 52(1-2)  
238 pp.36-38

239  
240  
241