
Case study

Nerve damage: catastrophic complications that need to be vigilant in the clinical application of triamcinolone acetonide, two case reports

Abstract

In this paper, we reported two cases of severe neurological complications caused by triamcinolone acetonide. One case was severe hearing loss on the same side after triamcinolone acetonide was injected around the ear to treat scars, and the other case was severe complications of lower limb paralysis after triamcinolone acetonide was injected into the spinal canal. In clinical practice, we should pay attention to the disastrous neurological complications caused by triamcinolone acetonide hydrochloride, and take relevant preventive and coping measures.

Keyword: Neurological Complications; Triamcinolone Acetonide

Introduction

Triamcinolone acetonide, a long-acting adrenocortical hormone, is commonly used in the treatment of rheumatoid arthritis, allergic dermatitis, skin scars, and chronic pain in the joint cavity or epidural space injection (Song,2019; Van Boxem,2019). However, we should be alert to the disastrous neurological complications caused by triamcinolone acetonide.

Case presentation

Recently, two cases of triamcinolone ~~acetonide~~acetonide-related catastrophic

neurological complications occurred in our hospital. The first case was an 18-year-old female, ASA grade I, without any basic disease, in generally good condition. Because of the scar on the earlobe, the cosmetic physician injected triamcinolone acetonide and lidocaine injection into the scar. The patient immediately developed local large-area ecchymosis, and the ipsilateral hearing was severely damaged, resulting in nausea, vomiting, and vertigo. After 2 months of treatment, the patient's nausea, vomiting, and vertigo improved, but the hearing damage did not improve. The second patient was a 55-year-old female. Due to postherpetic neuralgia, the left T10, T11, and T12 nerve roots were blocked under the guidance of an X-ray. The injection drug was triamcinolone acetonide combined with lidocaine. The operation was smooth. Two hours after the injection of the drug, the patient developed paraplegia in both lower limbs. After one month of treatment, it still did not improve.

Discussion

The above two cases of catastrophic neurological complications are speculated to be related to the complications of triamcinolone acetonide. Triamcinolone acetonide is a suspension, insoluble in water, with a large number of particles gathering, and the largest particle is greater than 500 μ m. It has been reported that the medium-sized particles of the adrenocortical hormone with a diameter of 51-to 1,000 μ m can easily block the blood vessels, especially the anterior spinal artery (Dietrich, 2015; Mehta, 2017). Although intratympanic injection of triamcinolone acetonide can be used to treat hearing impairment, severe hearing impairment caused by topical injection into the earlobe is very rare (Dahm, 2019). The serious hearing impairment

and paraplegia of lower limbs caused by triamcinolone acetonide in the above two cases are likely to block the capillaries of the inner ear or the blood vessels of the spinal cord with triamcinolone acetonide granules, resulting in serious dysfunction of the inner ear and spinal cord. In addition, the preservative of triamcinolone acetonide contains benzyl alcohol, which may have certain neurotoxicity to the adjuvant itself and cause a corresponding neuroinflammatory reaction.

Therefore, ~~in view~~because of the disastrous consequences of triamcinolone acetonide, we should be alert to its serious neurological complications when using triamcinolone acetonide clinically. Try to avoid using it in some parts with abundant blood supply and important nerve function. During use, it should be closely observed, relevant complications should be treated as soon as possible, and irreversible nerve complications should be avoided.

Reference

Dahm, V., Nieratschker, M., Riss, D., Kaider, A., Auinger, A., Honeder, C., & Arnoldner, C. (2019). Intratympanic Triamcinolone Acetonide as Treatment Option for Idiopathic Sudden Sensorineural Hearing Loss. *Otology & neurotology: official publication of the American Otological Society, American Neurotology Society [and] European Academy of Otolgy and Neurotology*, 40(6), 720–727.
<https://doi.org/10.1097/MAO.0000000000002283>

Dietrich, T. J., Sutter, R., Froehlich, J. M., & Pfirrmann, C. W. (2015). Particulate versus

non-particulate steroids for lumbar transforaminal or interlaminar epidural steroid injections: an update. *Skeletal radiology*, 44(2), 149–155.

<https://doi.org/10.1007/s00256-014-2048-6>

Mehta, P., Syrop, I., Singh, J. R., & Kirschner, J. (2017). Systematic Review of the Efficacy of Particulate Versus Nonparticulate Corticosteroids in Epidural Injections. *PM & R: the journal of injury, function, and rehabilitation*, 9(5), 502–512.

<https://doi.org/10.1016/j.pmri.2016.11.008>

Song, H., Tan, J., Fu, Q., Huang, L., & Ao, M. (2019). Comparative efficacy of intralesional triamcinolone acetonide injection during early and static stage of pathological scarring. *Journal of cosmetic dermatology*, 18(3), 874–878.

<https://doi.org/10.1111/jocd.12690>

Van Boxem, K., Rijdsdijk, M., Hans, G., de Jong, J., Kallewaard, J. W., Vissers, K., van Kleef, M., Rathmell, J. P., & Van Zundert, J. (2019). Safe Use of Epidural Corticosteroid Injections: Recommendations of the WIP Benelux Work Group. *Pain practice: the official journal of World Institute of Pain*, 19(1), 61–92.

<https://doi.org/10.1111/papr.12709>