

Genetic Diversity and Phytochemical Studies on Selected Ecotypes of Utasi (*Gongronema latifolium*) Plant using Microsatellite Markers

Abstract

Diversity and phytochemical study was carried out on twelve ecotypes of utasi, *Gongronema latifolium* using simple sequence repeats markers. Screening of ten (10) trinucleotide SSR primers produced 4 primers that were able to amplify the DNA from all the plant ecotype. A total of 34 bands were amplified from the 4 SSR primers which includes SSR 2, SSR 3, SSR 1 and SSR 4. Out of the amplified products using the 4 primers, 17 were found to be polymorphic with an average of 9 bands per primers. Number of amplification products per primer ranged between 10 in SSR 4 to 12 in SSR 2 with SSR 1 having the least number of bands (1). The sequences of the 4 random primers used in this study along with the number of bands generated and the number of polymorphic bands. The results of genetic analysis of the gongronema latifolium ecotypes delineated the 12 ecotypes into five cluster groups. Results of phytochemical analysis of leaves of *Gongronema latifolium* showed (Table 2) mean saponin ($2.09 \pm 0.01\%$), mean tannin ($1.17 \pm 0.01\%$), mean flavonoid ($2.53 \pm 0.01\%$), mean polyphenol ($4.60 \pm 0.01\%$) and mean reducing compound ($7.92 \pm 0.01\%$). The phytochemical indices of the least $1.17 \pm 0.01\%$ and highest $7.92 \pm 0.01\%$ were constituted by tannins and reducing compounds, respectively. Thus, the study suggested the need for future exploitation of *Gongronema latifolium* leaves for phytochemical contents due its high medicinal value.

Introduction

Gongronema latifolium commonly known as Utazi and Arokeke by the South-South and South-West inhabitants in Nigeria is found in Africa, Asia and Oceania. It is a tropical phyto-protein that is used as spice (Ugochukwu *et al* 2003). Apart from the proteinous aspect of this plant, it is also used as a traditional medicinal plant due to its phytochemical composition for the treatment of various gastrointestinal disorders such as diarrhea, ulcers, dyspepsia and also in the management of diabetes mellitus (Okafor *et al* 1996; Nwing *et al* 2005).

Following domestication, genetic diversity in plants has continued to narrow down due to a continuous selection pressure for specific traits i.e. yield, thus rendering them more vulnerable to disease and insect epidemics and jeopardizing the potential for sustained genetic improvement

over a long term. Thus it is extremely important to study the genetic composition of the germplasm of existing modern-day cultivars in comparison with their related species. This will not only provide information on their phylogenetic relationship, but will also indicate a chance of finding new and useful genes, as the ecotypes with most distinct DNA profiles are likely to contain a great number of novel alleles (Harlan, 1981).

Genetic diversity, the level of biodiversity, refers to the total number of genetic characteristics in the genetic makeup of a species. It could also refer to both the vast number of different species as well as the diversity within specie. Genetic diversity serves as a way for populations to adapt to changing environments. With more variation, it is more likely that some individuals in a population will possess variations of alleles that are suited for the environment. Those individuals are more likely to survive to produce offspring bearing that allele. The population will continue for more generations because of the success of these individuals.

A 2007 study conducted by the National Science Foundation that genetic diversity and biodiversity are dependent upon each other, that diversity within a species is necessary to maintain diversity among species and vice versa. According to the lead researcher in the study, “if any one type is removed from the system, the cycle can break down, and the community becomes dominated by single species (Lankau, 2007).

Survival and adaptability of species bring about a high genetic diversity because when a species environment changes, slight gene variations are necessary to produce changes in the species anatomy that enables it to adapt and survive. Species that has a large degree of genetic diversity among its population will have more variations from which to choose the best fit alleles. Species that have very little genetic diversity are at a great risk because with very little genetic diversity within species, healthy reproduction becomes increasingly difficult, and offspring often deals with similar problems. The vulnerability of a population to certain types of diseases can also increase with reduction in genetic diversity.

Amongst the highly useful and medicinal important plants found in the Sub-Saharan regions of Africa, *Gongronema latifolium*, formerly called *Marsdenenia latifolium* Benth, is one of the most widely used species of its genus. It belongs to the Asclepiadaceae family. It is called “Utazi”, “Utezi” and “Arokeke” in south eastern and western Nigeria, respectively (Ugochukwu et al., 2003).

The plant is used as a leafy vegetable in south-eastern Nigeria and a good source of vitamins, protein, iron and minerals (Okafor, 2005). The medicinal importance of *G. latifolium* cannot be over emphasized. The plant plays a vital role in the treatment and prevention of varied health related problems including liver diseases, diabetes mellitus, high blood pressure, loss of appetite, dysentery, stomach pains, worm infectors, cough and malaria fever (Agbo et al., 2005; Okafor, 2005). Medicinal importance of the plant is further elaborated by the presence of five bioactive compounds including alkaloids, saponnins, tannins, flavonoids, and glycosides in leaves, which was suggested to proffer varied pharmacological effects on its specie (Gamaniel and Akah, 1996).

Plants improvement, either by natural selection or through the efforts of breeders, has always relied upon creating, evaluating and selecting the right combination of alleles. The manipulation of a large number of genes is often required for improvement of even the simplest of characteristics. With the use of molecular markers, it is now a routine to trace valuable alleles in segregating population and mapping them. These markers once mapped, enable dissection of the complex trait into component genetic units more precisely, thus providing breeders with new tools to manage these complex units more efficiently in a breeding programme (Hayes, 1993).

Molecular markers have been looked upon as tools for a large number of applications ranging from localization of a gene to improvement of plant varieties by marker-assisted selection. They have also become extremely popular markers for phylogenetic analysis adding new dimensions to the evolutionary theories. If we look at the history of the development of these markers, it is evident that they have been improved over the last two decades to provide easy, fast and automated assistance to scientist and breeders. Plants breeding analysis based on molecular markers has generated to preserve and popularize it (Flavell, 1995).

Microsatellite or simple sequence repeats markers are amplification products of anonymous DNA sequences using single, short di or tri oligonucleotide primers and thus do not require prior knowledge of a DNA sequence. Low expense, efficiency in developing a large number of bands in a short time and requirement for less sophisticated equipment has made the microsatellite technique valuable especially for the studies of genetic diversity amongst crops.

Hence this study seeks to unveil the diversity in different species of Utazi (*Gongronema latifolium*) with regards to its medicinal and molecular characteristics using reliable genetic markers.

Materials and Methods

2.1.1 Sample Collection

Young leaves of *Gongronema latifolium* obtained from different locations in Cross River, Akwa Ibom, Oyo and Edo States, Nigeria, (Table 1) was taken to the International Institute of Tropical Agriculture (IITA) Ibadan, Nigeria. They were given codes for easy identification and then stored in a cold room for future laboratory analysis.

Table 1:

Sources of plant samples used for the study

Designation	Ecotypes	Coordinates /elevation
G1	Oyo State, Ibadan	N06 ⁰ 12. 234 E006 ⁰ 21.235
G2	Oyo State, Ibadan	N06 ⁰ 05.135 E006 ⁰ 25. 328
G3	Oyo State, Ibadan	N06 ⁰ 40. 234 E006 ⁰ 25.235
G4	Oyo State, Ibadan	N06 ⁰ 15.221 E006 ⁰ 45. 328
G5	Akwa Ibom, Ibiakpan Ikot Ekpene L.G.A, Aks (1)	N 03 ⁰ 09. 678 E007 ⁰ 49.163, Elev. 99m
G6	Akwa Ibom Okobo, Ibiono Ibom L.G.A,	N03 ⁰ 05. 236

	Aks	E007 ⁰ 51.100 Elev 86m
G7	Akwa Ibom Ikot Ekpene Ii, Ibiono Ibom L.G.A, Aks	N03 ⁰ 11. 108 E07 ⁰ 55. 133 Elev 124m
G8	Cross River, Biological Science Block (1)	N04 ⁰ 56.981 E008 ⁰ 21.190 Elev. 32m
G9	Cross River, Biological Science Block (2)	N04 ⁰ 56. 981 E008 ⁰ 21.190 Elev. 32m
G10	Cross River, Old Odukpani Road	N04 ⁰ 56.981 E008 ⁰ 21. 191 Elev. 34m
G11	Edo State, Eguare Irrua	N06 ⁰ 44. 234 E006 ⁰ 45.235
G12	Edo State, Ikhideu-Egoro	N06 ⁰ 45.235 E006 ⁰ 05. 328 Alt 373m



Fig. 1 showing leaf samples of *Gongronema latifolium*

2.1.2 Total genomic DNA extraction from Utazi leaves

Genomic DNA was extracted from fresh leaves of each sample using a modified method of Stothardet *et al* (1996). Leaves samples were mascerated and total genomic DNA was extracted from each sample using CTAB extraction buffer containing 2-merccaptoethanol, hexadecyltrimethyl-ammonium bromide (CTAB) (solid), tris (hydroxyl-mehyl) amino-methane, ethlenediamine-tetraacetic acid, disodium salt solution (EDTA), and sodium chloride. Each sample were soaked in TE (10mM tris HCL and 1mM EDTA) to get rid of the remaining

ethanol. Tissue from each of the leaves were placed in sterile 1.5ml Eppendorf tube, 500 μ l of CTAB solution added and 10 μ l of proteinase was added too. The genomic DNA were extracted from the CTAB buffer by adding an equal volume of chloroform and isoamyl alcohol to each tube, the organic and the aqueous layers were gently mixed for 5mins and spun at 13,000 rpm for 20 mins. before the upper aqueous layer were removed into another sterile eppendorf tube and equal volume of 100% ethanol were added, mixed and incubated at -20°C overnight to enhance DNA precipitation, it was spun again for 13,000 rpm for 20mins. The pellet were washed with 70% ethanol and spun for another 20 mins, the supernatant were removed and the pellets dried at room temperature.

2.1.3 DNA quantification and quality determination

The quantity and quality of genomic DNA extracted was examined by comparing the template DNA isolated from samples with a DNA ladder (gene ruler) of 50bps to 2kbp for SSR in a 1% agarose gel using 1 x TBE buffer and viewed in a gel box (G: Box, Syngene). The concentration and quality was further determined at optical density (OD) readings of 260nm and 280nm using a Nano-drop spectrophotometer (Thermo Scientific Nano-Drop 2000C). The concentrations were used to guide the normalization of DNA of each sample at a concentration of 20ng/ μ L. Additionally, the ratio of OD 260/280 was provided by the Nano-drop which gave an indication of purity of the samples. Pure DNA has OD260/OD280 value of 1.8 and a deviation from this signifies the presence of contaminants that inhibit PCR reaction.

2.1.4 Polymerase chain reaction optimization

Polymerase chain reaction was carried out in an Eppendorf Master Thermocycler (Eppendorf, company, USA). Each reaction tube was made of 100ng of template DNA, 2.5mM of MgCl₂, 100 μ M of dNTPs, 1X *Taq* buffer, 20 picomolar of 10-mer Primer, and 1 unit of *Taq* DNA polymerase made to a final volume of 20 μ l. DNA amplification was performed using the following thermal profile or sequence: 94°C for 5 min (35 Cycle); 94°C for 1 min, 37°C for 1 min, 72 °C for 2 min (35 cycles); final extension was performed at 72°C for 10 min (35 cycle) and samples were cooled at 4°C. PCR optimization was carried out using six selected DNA samples. A pre-mix containing dNTPs (dATPs, dCTPs, dGTPs and dTTPs), MgCl₂, Tris-HCl (pH 9.0), KCl and *Taq*DNA was used. A master mix containing 2 μ l of sterile distilled water, 0.5 μ l of 10pmoles forward primer, 0.5 μ L of 10pocomoles reverse primer, 5 μ l premix and 2 μ l of template DNA was prepared. The diluted DNA samples were subjected to polymerase chain reaction (PCR) amplification using simple sequence repeat (SSR) markers.

2.1.5 Gel electrophoresis of PCR products

Leaves samples per marker were separated on 1.0% agarose gel at 80V for 40 minutes. Agarose powder was dissolved in Tris-borate EDTA (1x TBE) buffer by slowly boiling in a microwave oven. The agarose was allowed to cool and 1mg/ml concentration of ethidium bromide was added to the gel. The warm agarose solution was then poured into the gel tray in which combs were inserted to form sample wells. The gel was allowed to solidify for 30 minutes before immersing in the electrophoresis tank containing 100ml TBE buffer. The samples were run alongside 1.0 μ L 1kb DNA ladder at 80 volts for 40 minutes. The amplified products were

viewed under UV light in a gel box (G: Box, Syngene). Four of the primers showed amplification at various degrees while three primers did not amplify at all. Among the 12 primers that showed amplification, twelve (12) primers, six for each of SSR best were selected to amplify the polymorphic ecotypes.

2.2 Phytochemical analysis of the plant extracts of *G. latifolium*

Phytochemical analysis was carried out on the powdered and aqueous extract of samples of *G. latifolium* leaves using standard procedures to identify the constituents as described by Sofowara (1993). The phytochemical analysis was carried out to determine the presence of the following chemicals in the plant extracts: tannins, saponins, flavonoids and alkaloids.

2.2.1 Determination of Tannins

About 0.5g of the dried powdered sample was boiled in 20 .0 millimeters (ml) of water in test tube and then filtered. A few drops of 0.1 (%) ferric chloride were then added. Observation of a brownish green colour indicated the presence of tannins.

2.2.2 Determination of Saponins

About 2.0g of the powdered sample was boiled in 20.0ml of distilled water in a water bath and filtered. Ten ml. of the filtrate was mixed with 5.0ml of distilled water and shaken vigorously for a stable persistent froth to occur. The frothing was mixed with 3 drops of olive oil and shaken vigorously. Formation of emulsion indicated the presence of saponin.

2.2.3 Determination of Flavonoids

One ml. of 10.0% lead acetate solution was added to 1.0ml. of aqueous extract of the plant. The formation of a yellow precipitate indicated a positive test for flavonoids.

2.2.4 Determination of Alkaloids

Three ml. of aqueous extract was stirred in 3ml. of 1% Hcl on a steam bath. Mayer's and Wagner's reagents were added to the mixture. Turbidity of the resulting precipitate indicated the presence of alkaloids.

Results

Genetic diversity is of great significance for breeding programmes as well as taxonomic studies. Molecular markers have frequently been used for the detection of genetic diversity in plants. It is mostly used because of its rapidity, simplicity and lack of any prior genetic information about the plant.

The SSR banding pattern of the 12 ecotypes of *Gongronema* are illustrated in fig. 1. After screening 10 trinucleotide SSR primers, 4 primers were able to amplify the DNA from all the plant ecotype. A total of 34 bands were amplified from the 4 SSR primers which includes SSR 2, SSR 3, SSR 1 and SSR 4. Out of the amplified products using the 4 primers, 17 were found to be polymorphic with an average of 9 bands per primers. Number of amplification products per primer ranged between 10 in SSR 4 to 12 in SSR 2 with SSR 1 having the least number of bands

(1). The sequences of the 4 random primers used in this study along with the number of bands generated and the number of polymorphic bands is given in table 2.

Ward linkage generated a dendrogram which illustrated the overall genetic diversities and relationship amongst the ecotypes surveyed. This is illustrated in fig. 2.

The first cluster consisting of seven ecotypes of *G. latifolium* were further classified into three sub-clusters, 1A, 1B and 1C. "1A" consist of G9, G11 and G5, which were ecotypes gotten from Cross River, Edo State and Akwa Ibom State respectively.

"1B" consists of G6, G8 and G10. These were ecotypes gotten from Akwa Ibom (G6) and G8 and G10, collected from Cross River State. "1C" has an ecotype, G12 which was gotten from Edo State. (Fig. 2)

The second cluster contains two ecotypes G2 and G3 collected from Oyo State, Ibadan, while the third cluster has 3 ecotypes, G4, G7 and G1. These ecotypes G4 and G1 were gotten from Oyo State, Ibadan, while G7 was gotten from Akwa Ibom State.

It can be observed from the Dendrogram that even though the *G. latifolium* ecotypes, G9, G11 and G5 from Sub-cluster 1A were gotten from different locations, they had a genetic similarity. Morphologically the leaves and plant growth habit were same; G11 had high altitudes of 373 as compared to G9 and G5 having low altitudes of 32m and 99m respectively. Also, the leaves of G9 and G5 were broad as compared to G11 having small leaves. This was because people from parts of Edo state where this ecotype was collected complained of lack of water, making most of the plants small and dark in colour.

Sub-cluster 1B, containing G6, G8 and G10 were ecotypes gotten from Akwa Ibom and Cross River and they formed a cluster, showing that they have a genetic similarity. Morphologically leaves were the same, broad, light and had thin stems.

The clustering pattern of G12, collected from Edo state indicates its genetic diversity, which might be due to its location. It had a very high Elevation of 405m above sea level as compared to other ecotypes.

Cluster 2 containing G2 and G3, collected from Oyo state, Ibadan clustered together showing its genetic relatedness amongst the two ecotypes. Also cluster 2 had G4, G7 and G1 collected from Oyo state and Akwa Ibom State. These three ecotypes clustered indicating their relatedness even if they were collected from different locations.

Ecotype from Oyo State Ibadan from 001 -04 were properly amplified these ecotypes. This implies that SSR1 does not have a corresponding sequence with these ecotypes, hence resulting in poor amplification product.

G5, G6, and G7 which were gotten from Akwa Ibom State produced poor amplification products using the primers especially in Ecotypes 5 and 6. Ecotype 7 was only amplified using SSR2 and SSR3, but SSR4, SSR1 produced no bands. This shows some level of genetic diversity as most of the primers do not have a corresponding sequence with the ecotypes.

Ecotypes 8, 9 and 10 collected from Cross River produced poor amplification products in SSR 3, SSR 4. SSR 2 produced bands in ecotype 8 and 10. SSR 1 produced bands in ecotype 10. This however shows genetic diversity amongst this ecotype as selected primers were able to amplify some ecotypes, produce unique bands or produce no band at all.

Ecotype 11 and 12 gotten from Edo State had an amplification product using SSR 1, SSR 3 and SSR 4. Only SSR 2 was able to produce reproducible bands in ecotype 12. SSR 2 therefore has a sequence that corresponds to ecotype 12 but does not correspond to ecotype 11. This shows some extent of diversity between the primers and the ecotype.

All the 4 primers produced different number of polymorphic bands when used to screen the ecotypes. This explains reasons for the diversity observed amongst *G. latifolium* ecotypes suggesting that the more polymorphic the bands are the more diverse the plant. This also explains the variations in the chromosome Aikpokpdion et al., (2012).




Fig. 2: Dendrogram showing genetic relationship between 12 Ecotypes of *Gongronema latifolium*.

Phytochemical analysis of *Gongronema latifolium*

TABLE 2

Phytochemical profiles of *G.latifolium* leaf

Phytochemical Indices	Values (%)
Alkaloids	2.40 ± 0.06

Glycosides	2.26 ± 0.01
Saponins	2.09 ± 0.01
Tannins	1.17 ± 0.01
Flavonoids	2.53 ± 0.01
Polyphenol	4.60 ± 0.01
Reducing sugars	7.92 ± 0.01

Results of phytochemical analysis of leaves of *Gongronema latifolium* showed (Table 2) mean saponin (2.09 ± 0.01%), mean tannin (1.17 ± 0.01%), mean flavonoid (2.53 ± 0.01%), mean polyphenol (4.60 ± 0.01%) and mean reducing compound (7.92 ± 0.01%). The phytochemical indices of the least 1.17 ± 0.01% and highest 7.92 ± 0.01% were constituted by tannins and reducing compounds, respectively.

Discussion

Gongronema latifolium family Asclepiaceae is a creeping plant, geographically distributed in Africa, tropical and sub-tropical Asia and Oceania. In Nigeria, it is commonly called Utazi and arokeke in the south south and south west geopolitical zones. It is a tropical rain forest plant protein source used as vegetable spice (Ugochukwu *et al*; 2003). *Gongronema latifolium* leaf meal flowers have revealed the presence of sure nutritional factors from the leaf, stem and root which varies in health applications respectively (Essien *et al* 2007). It has been used in the traditional system of medicine for various gastrointestinal disorders such as diarrhea, ulcers and dyspepsia and also in the management of diabetes mellitus (Okafor *et al*; 1996; Nwing, *et al* 2005). The leaves have been reported to have a hypoglycaemic effect (Ugochukwu

and Babadyi, 2003; Ogundi *et al* 2003) by decreasing activity of glucokinase enzyme and level of hepatic glycogen, and blood glucose. Phytochemical studies of *G. latifolium* showed that the root contains polyphenol in abundance, alkaloids, glycosides and reducing sugar (Antai *et al* 2009). Mensal *et al*; (2008) analysed the effect of *G. latifolium* on immune system of birds, and reported that it contained important compounds and served as antibiotics for treatment of common pathogenic strains of infective agents in bird and also used for prevention and treatment of diseases that could cause mortality in farm animals (Ugochukwu *et al* (2003) and Agbe *et, al* (2005).

Nutritionally, *G. latifolium* is rich in fats, protein, vitamins, minerals and essential amino acids (Eleginmi, 2007). Ani *et al* (2013) while feeding broiler birds on varying dietary levels of *G. latifolium* leaf meal recorded positive effect on growth performance of chicks at 75% level of *G. latifolium* leaf meal per 25kg of feed. Afolabi, (2007) investigated the chemicals composition and anti-bacterial activity of *G. Latifolium* and observed that *G. latifolium* was a good source of protein content at the rate of 27.2% in dry matter, high and compared favourably with percentage dry matter values reported for chickpea (24.0%), cowpea (24.7%), Lentil (26.1%), fluted pumpkin leaves (22.4%) *Tamarndus indica* (24.3%), *Mucana flagellipies* (24.9). *Hibiscus esculantus* (23%) and *Parkiabig lobosa* (20.9%). Several previous and current studies on *G. Latifolium* reported that it has nutritional high protein, vitamin and minerals. Glew *et al* (1997), Akwaowo *et al* (2000); Ajayi *et al* (2006); Igbal *et al*, (2006) Igbal *et al*, (2006); Okafor (2005); Kubarava *et al* (2007) in their analyses of *G. latifolium* found that it was is the cheapest most available source of important protein, vitamin, minerals and essential amino –acids that boost the physiological status of humans and promoted their growth. Tacon *et al* (2006) findings supported

it as locally available and cheap plant protein source which is very important for future development of phytomedicine.

Conclusion

This study used molecular technique to produce basic information on the genetically diverse nature of *Gongronema latifolium* ecotypes obtained from the different ecologies. The results of genetic analysis further revealed that the utasi ecotypes shared common ancestral origin commonly segregated into five cluster groups. Phytochemical analysis of utasi ecotypes revealed non significant variations among the valuable presence of alkaloids. Flavonoids, saponins, tannins, glycosides and reducing sugars which confers its medicinal properties. It has also shown that even though most plants are the same phenotypically and have same names, there is always a character (at the molecular level) that separates one plant from the other.

References

- Afolabi, F. E. (2007). Chemical composition and antibacterial activity of *Gongronema latifolium*. *Journal of Zhejiang University*, 8(5): 252-358.
- Agbo C. U, K.P. Bayeri and I.U Obi (2005), *Indigenous Knowledge And Utilization Of Gongronema lati folium Benth: A Case Study Of Women In University Of Nigeria*, Nsukka. Bio- Research journal. 3(2) 66-69
- Aikpokpodion P.f O.A. Peggy and U. Edak (2012), *Karyotype Analysis And Ploidy Determination Using Flow Cytometry In African Bitter Milk Plant 'utasi', Gongronema latifolium Benth*. *Cytoiogia* 77(1): 1-10
- Akuodor, G. C, Idris Usman, Ms mban, C. C, megwas, UA, Apkan, J. L, Ugwu, T.C, D. O. Okoroafor and U. A. Osunkwo (2010). *Studies On Antl-Ulcer, Analgesic And Antipyretic Properties Of The Ethanolic Leaf Extract Of Gongrenema latifolium In Rodents*, African journal of Biotechnology 9(15): 2316-2321.
- AIFP (2004). Farming Nigerian water. *News letter of the Aquaculture and Inland Fisheries Project of National Special Programme for Food Security in Nigeria* (F.A.O Office). Abuja, Nigeria.
- Ajani, E. K., Nwana, L. C. & Musa, B. O. (2004). Replacement of fish meal with maggot meal in the diets of Nile tilapia, *Oreochromis niloticus*. *World Aquaculture*, 35: 52-54.

- Ajayi, A. I., Oderinde, R. A., Kajogbola, D. O. & Ukponi, J. U. (2006). Oil content and fatty acid composition of some underutilized legumes from Nigeria. *Food Chemical*, 99(1): 115-120.
- Akinwande, A. A., Alatise, P. S. & Ayanlode, F. V. (2006). Evaluation of pawpaw leaf-based carica papaya, on fingerlings reared in plastic bowes. *Proceeding of the 21st and Annual Conference of the Fisheries Society of Nigeria (FISON)*, Calabar, 13-17 November, 320-324.
- Akwaowo, E. U., Ndon, B. A. & Etuk E. U. (2007). Minerals and antinutrients in fluted pumpkin. *Food Chemical*, 70(2): 235-240.
- Al-faragi, S. K. H & Al-saphar, S. A. A. (2013). Effect of local probiotic on common earp cyprinus carpio growth performance and survival rate. *Journal of Genetic and Environmental Resources Conservation*, 1(2): 89-96.
- Amisah, S., Oteng, M. A. Ofori, J. K. (2009). Growth performance of the African Catfish *Clarias gariepinus* fed varying inclusion levels of *Leucaena leucocephale* leaf meal. *Journal of Applied Science Environmental Management*, 13(1): 21-26.
- Ani, A. O., Ogbu, C. C., Abakasanga, I. U. & Ugwuowo, L. C. (2013). Response of broiler birds to varying dietary levels of *Gongronema latifolium* leaf meal. *Journal of Biology, Agriculture and Health Care*, 3(14): 2224-3208.
- Antai, A. B. (2009). Physiochemistry and some hematological changes following oral administration of ethanolic root extract of *Gongronema Latifolium* in rats. *Nigerian Journal Physiological Science*, 24(1): 79-83.
- AOAC, (2000). *Association of official analysis chemist international (AOAC)*. (17th ed.). USA: AOAC International, Gaitherburg, M.
- AOAC. (2000). *Association of official analytical chemist official method of analysis*. (18th ed.). Washington, DC: AOAC International, Gaitherburg, M.
- Argyropoulou, V. N., Kalogeropoulos, M. I. & Alex, M. N. (1992). Effect of dietary lipids on growth and tissues fatty acid composition off grey Mullet *Mugil Cephalus*. *Biochemistry and Physiology*, 101, 129-135.
- Auta, J. (1993). Water quality management in fish ponds. *Proceeding of National Workshop on Fisheries Extension Delivery*, 2.
- Ayanda, J. O. & Tafida, A. A. (2006). In land fisheries in Nigerian economy: Issues and challenges in Fisheries Society of Nigeira (FISON) Conference Proceedings, Calabar, Nigeria 13th – 17th November, 2016. Pp.174-177.
- Ayoola, S. O., Ajani, E. K. & Fashae, O. F. (2013). Effect of probiotics (Lactobacillus and Bifidobacterium) on Growth performance and haematological profile of *Clarias gariepinus* Juveniles. *Journal of Fish and Marine Sciences*, 5(1): 01-08.

- Babatunde, G. M., Fajimi, A. O. & Oyejide, A. O. (1992). Rubber seed oil versus palm oil in broiler chicken diet, effect on performance nutrient digestibility haematology and carcass characteristics. *Animal Feed Science and Technology*, 35, 133-146.
- Bagenal, T. B. (1978). *Method of assessment of fish production in fresh water*. Oxford: IBP Blackwell Science Publication. Handbook No. 3.
- Bajerjee, S. K., Pcita, B. C., Bamdoyopahyay, P & teway, A. (2002). Change of blood parameters in carp, catla. *Journal of Aquatic Biology*, 17: 79-84.
- Baker, F. J., Silverton, R. E. & Pallister, C. J. (1980). *Introduction to medical laboratory technology*. (7th ed). London: Butter Worths.
- Banjerjee, S. S., Pcita, B. C., Bandoyopahyay, P. & Teway, A. (2002). Change of blood parameters in carp, catla. *Journal of Aquatic Biology*, 17: 79-84.
- Bardach, J. J., Ryther, H. & Mclarney, W. O. (1972). *Aquaculture farming and husbandry of fresh water and marine organisms*. London: Prentice Hall.
- Barker, F.J. & Silverton, R.E (1985). Visual red cell count. In F. J. Baker, R.E. Silverton and C. J. pallister (Eds), *Introduction to medical laboratory technology*. London: Butter Worth.
- Barker, J. (1988). Simply fish London: Faver & Faber, 36-39 J. G. Bell, McEvoy, D. R. Tocher, F. Mcahee & P. J. Campbell. Diets of Atlantic Salmon (*salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid metabolism. *Journal Nutrition*, 4(1): 126-131.
- Berjerano, I. (1984). Detection and control of stress conditions in warm water aquaculture. *Research on Aquaculture*, 2(1): 56-68.
- Bichi, A. H. & Ahmad, M. K. (2010). Growth performance and nutrient utilization of Africa catfish (*Clarias gariepinus*) fed varying dietary levels of processed cassava leaves. Bagero. *Journal of Pure and Applied Science*, 3(1), 118-122.
- Blaxhall, P. C. (1992). The haematological assessment of health of fresh water fish. *Journal of Fish Biology*, 4593-605.
- Bolger, S. R. & Connolloy, P. L. (1989). The selection of suitable indices for the measured and analysis of fish conduction. *Journal of Fish Biology*, 34 171-182.
- Bond, C. E. (1996). *Biology of fishes*. (2nd ed.). New York: Soundness College Publisher.
- Bovisova, T. A., G. E. Souza, D. Miller W. C. Labys, (2007). *Remaining competitive at the regional level developing a local aquaculture industry. I of aquaculture economics & Mg; (II) vol, Las vegas*. USA: Nevada.
- Boyd, C. E. (1979). *Water quality in warmwater fish ponds*. Alabama Auburn: University Agriculture Experiment Station, Auburn.

- Boyd, C. E. & Lickotocoper, F. (1990). *Water quality in pond for aquaculture, Alabama agriculture*. Alabama, Auburn: Experimental Station Auburn University. British Standard of measurement for packed cell volume, (1968). *British Standard*, 4316.
- Bromage, N. R. & Roberts, R. J. (1995). *Brood stock management and egg and larval quality*. Osmead, Oxford, UK: Blackwell Science.
- Bruton, M. N. (1979a). *The food and feeding behaviour of Clarias gariepinus (pisce, claridae) in lak sibaya, South Africa with species. It emphasises on its role a predator of chuids*. London: Transaction of Zoological Society.
- Burkill H.M, (1985). *The Useful Plants of West Tropical Africa. 2nd edition. Volume 1, Families A-D*. Royal Botanic Gardens, Kew, Richmond, United kingdom. 960pp
- Changadeya, W. L., Malekano, B. & Ambau, A. J. D. (2003). Potential of genetics for aquaculture development in Africa in NAGA. *World Fish Center Quarterly*, 26(3): 31-35.
- Chude, L. A. (2001). Identifying Ikwnano catchment area and fish production aquaculture perspective. *Proceedings of the 16th Annual Conference NAAS, Nigeria*. Pp. 172.
- Clark, S., Whiltemore, D. H. J. & McMahon, R. F. (1979). Consideration of blood parameters of large mouth bass (*Micropterus salmoides*). *Journal of Fish Biology*, 14 (2): 147-154.
- Clay, D. (1979). Population biology, growth and feeding the African catfish, *Clarias gariepinus*. With special reference to juveniles and their importance in fish culture. *Archive of Hydrobiology*, 87(4): 453-482.
- Cowey, C. B. & Roberts, R. J. (1978). *Nutritional pathology*. London: Billier Tindall.
- Cowey, C. B., Mackie, A. M. & Bell J. G. (1985). *Nutrition and feeding in fish*. London, Academic Press.
- Dasgupta, S., Davis, K. J. & Wirth, F. F. (2004). Selling freshwater prawn in food festivals: A marketing experiment in Kentucky. *Journal of Aquaculture Economic and Management (II)*, 5(4): 3-16.
- Davis, O. A., Inko, M. B., Tariah, D. & Amache, E. (2006). Growth response and survival of heterobranchus longifiis fingerlings fed at different feeding frequencies. *African Journal of Biotechnology*, 5, 778-780.
- De Graof, G. J., Galemon, F. & Banzoussi, B. (1995). The artificial reproduction and fingerling production of catfish *clarias gariepinus* (Burchell 1988) in protected and unprotected ponds. *Aquaculture Research*, 26, 233-242.
- De Silva, S. S. & Anderson, T. A. (1995). *Fish nutrition in aquaculture*. London: Chapman and Hall.

- Demma, J., Gebre-Mariam, T., Asres, K., Evgetie, W. & Engindawork, E. (2006). Toxicological study on *Glinus lotoides*, a traditionally used technical herb in Ethiopia. *Journal of Ethnopharmacology*, 111: 451-457.
- Docan, A., Cristea, V., Grecu, I. & Dediv, L. (2010). Haematological response of the European catfish, *Silurus glanis* reared at different densities in flow-through production system. *Archiva Zootechnica*, 13, 263-270.
- Edwin, H. R., Menghe, H. L. & Bree, B. M. (2006). *Catfish nutrition nutrients requirements*. London: Msu Publications and Information.
- Ekanem, A. P. & Ekanem, I. E. (2010). Effects of photo-protein and normal diet on the growth and Gonadal development of *Clarias gariepinus*. *Tropical Environmental Research*, 9: 556 -559.
- Ekanem, A. P., Eteng, S. U. Nwosu, F. M. & Eyo, V. O. (2012). Comparative study of the Growth and Gonad development of *Clarias gariepinus* (Burchell 1822) fed diet, with plant and animal –based ingredient in concrete tanks. *Journal of Agricultural Science and Technology*, 2: 1203-1210.
- Ekanem, A. P., Eyo, V. O., James, P. U & Udo, N. E. (2013). Effect of Unical feed on fecundity and gonad development of *Clarias gariepinus*: A comparative study with coppens commercial feed in earthen pond. *International Journal of Science and Research*, 2 (10) 8-14.
- Ekanem, A. P., Victor, O. E., Austin, I. O. Udeme, I. E. & Udo, P. J. (2012). Growth performance and food utilization of the African catfish (*Claris gariepinus*) fed uncial. Aqua feed and coppens commercial feed. *Journal of Marine Biology and Oceanography*, 1: 2.
- Ekprikpo-Wilfred, P. C. (1992) Hematological parameters of healthy catfishes *Hetrobranchus bidorsalis*, University of Calabar Nigeria
- Eleyinmi A.F, (2007). *Chemical Composition and Antibacterial Activity Of Gongronema latifolium*. *Journal of Zhejiang University, Science B* 8(5): 352-358
- El-sayed, A. F. M. (1994). Evaluation of soya bean meal, spirulina meal and chicken offal meal as protein sources for silver sea bream fingerlings. *Aquaculture* 127: 169-176.
- England, J. M. Walford, D. M & waters, D. A. N. (1972). Re-assessment of the reliability of haematocrit (MCHC). *British. Journal of Haematology*, 23, 242-247.
- Erondu, E. S., Nnubia, S. & Nwadukwe, F. O. (1993). Hematological studies on four catfish species raised in fresh water pond in Nigeria. *Journal of Applied Ichthyology*, 9(3-4), 250-256.
- Essien. J. P. Ebong. G. A & Akpan, E. J. (2007) Antioxidant and Antitussive properties of *Gongronema latifolium*. *Applied Science Environmental Management*, 11(4): 47-50.

- Eyo, A. A. (2003). Fundamentals of fish nutrition and diet development. An overview” *National workshop on fish feed development and feeding practices in Aquaculture organized by fisheries society of Nigeria (FISON) in collaboration with institute of freshwater fisheries research (NSPFS) and FAO National Special Programme for food security (NSPFS)*: 1-33.
- Eyo, V. O. & Ekanem, A. P. (2011). Effect of feeding frequency on growth, food utilization and survival of (*clarias gariepinus*) using locally formulated diet. *African journal of environmental pollution and health* 9(2): 11-17.
- Eyo, V.O., Ekanem, A. P. & Jimmy, U. U. (2004). A comparative study of the gonadosomatic index (GSI) and Gonad gross morphology of African catfish (*Clarias gariepinus*) fed Unical aqua feed and coppens commercial feed. *Crotian Journal of Fisheries*, 72: 63-69.
- Ezemot Nwadiaro, R. I. (2009). A guide to biostatistics and health research method. *Hypothesis Testing*, 6: 142-147.
- Ezenwa, B. I. O. (1994). Aquaculture development in Nigeria, 41-80. In A. Coche, (Ed). *Aquaculture development and research in sub-Saharan Africa*. National Reviews CIFA Technical Paper No. 23. Suppl. Rome. FAO. 397.
- Fafioye, O.O., Falade, S. O., Adebisi A. A., Jonyo-oni & Omoyinmi G. A. K. (2005). Effects of dietary soybeans (alycine max) on growth and body composition of African catfish (*clarias gariepinus*, Burchell) Fishgerlings. *Turkey Journal of Aquatic Science*, 5: 11-
- Flavell, R.B (1995). *Generating Tools For Marker-Assisted Selection In Plant Breeding*. Tibtech, 13: 313-319.
- Gamaniell, K.S and P. A .Akah (1996). *Analysis Of The Gastrointestinal Relaxing Effect Of The Stem Extracts Of Gongronema latifolium*, *Phytomedicine* 2(4): 293-296.
- Global Biodiversity Information Faciitiy(GBIF). Accessed November 24, 2011.
<http://www.gbif.org>
- Harlan J.R (1981). *The Origin Of Indeginous African Agriculture*. *Cambridge History of Africa, vol 1: From Earliest Times to caSOOBC* (3.D. Clark, ed.). Cambridge University Press, Cambridge UK. PP 623-658.
- Hayes P.M (1993). *Theoritcal Applied Genetics*. 87, pg 392-401
- Lankau R.A (2007), *Mutual Feedbacks Maintain Both Genetic and Species Diversity in a Plant Community*. *Science* 317: 1561-1563.
- Mahmood, T. S. Muhammad and Z. K Shimwari (2010). *Molecular and Morphological Characterization of Caralluma species*. *Pak. S. Bot*, 42(2): 1163-1171

- Morebise O., M.A .Fafunso, J.M. Makinde, O.A. Oiajide and E.O.Awe (2002). *Antiinflammatory Property of the Leaves of Gongronema latifolium*. *Phytother Res*; 16 (si): 575-577. doi : 10.1002/ ptr. 784 (Pubmed) (Cross Ref)
- Nair,S and R. Keshavachandran (2006). *Molecular Diveirsity on Chakkarakoli (Gymnema sylvestre R.Br,f) Assessed Through Isozyme and RAPD Analysis*, 3. *Trop. Agric*, 44(1-2): 31-36
- National Biological Information Infrastructure^ BII). Accessed March 1, 2011. <http://www.nbli.gov>
- Sanibare, M. A and Z. O. Gbile (2008). *Ethnobotanical Survey Of Anti-Asthmatic Plants in Southern Nigeria African Journal of Traditional, Complementary and Alternative Medicine* S(4): 340-345
- Siddiqui,A.A, B.Ahmand and A.Dongra (2000). *Development in Chemistry and Pharmacology of Gymnema sylvestre*. 3 *med. Aroni plant- sci*, 22:223-231
- Ugochukwu N. H and Babady, N. E. (2003), *Antinyperglycemic Effects of Aqueous and Ethanolic Extracts of Gongronema latifolium Leaves on Glucose and Glycogen Metabolism in Livers of Normal and Streptozotoan Induced Diabetic Rats*, *Life sci*: B(15): 1920-1938, doUIQ. 1016/50024-3205(03) 00543-5. (pub med) [cross ref],
- Ugochukwu N. H, and N. E Babady (2002). *Antioxidant Effects of Gongronema latifolium in Hepatocytes of Rat Models of Non-Insulin Dependent diabetes (Diabetes melHtus*. *Fitorapia*. 73 (7-8): 612-618. DoUIO. 1016/50367-326(02) 00218-6 {pubmed} cross ref}.
- Williams J. G. K et al, (1990). *DMA Polymorphisms Amplified by Arbitrary Primers are Useful as Genetic Markers*. *Nucleic acids Res*. 18, 6531-6535
- Zabeau, M and P, Vos (1993). *Selective Restriction Fragment Amplification: a General Method for DNA Fingerprinting*. European patent office, publication 0534 854 A1, bulletin 93/13.