

Short communication

Comment [LC1]:

First Report of the Crapemyrtle aphid, *Tinocallis kahawaluokalani* (Hemiptera : Aphididae) and its local predaceous insects in Gabon

Comment [LC2]: Crape Myrtle aphids or Crapemyrtle aphid

Comment [LC3]: Topic is not easy to understand. I suggest,

Impact of Crape Myrtle aphids *Tinocallis kahawaluokalani* Kirkaldy (Hemiptera : Aphididae) and its related Natural Enemies

ABSTRACT

A study was carried out among some green spaces in Gabon, mainly in Libreville, Lambarene, Mouila, Franceville and Mitzic, from August 2019 to October 2021. The aim was to assess the presence and impact of the Japanese aphid *Tinocallis kahawaluokalani* Kirkaldy (Hemiptera : Aphididae), and its related natural enemies, using one-time visual observations on Crapemyrtle shrubs. This ornamental plant insect pest was recorded, occurring on leaves of *Lagerstroemia indica* (L.) Pers. (Lythraceae). Seven local beneficial insect species were observed preying on this aphid, mainly three hoverflies *Ischiodon aegyptius* Wiedemann, *Paragus borbonicus* Macquart, *Asarkina* sp. (Diptera : Syrphidae), three ladybird species *Platynaspis capicola* Crotch, *Scymnus interruptus* Goeze and *Chilochorus nigratus* Fabricius (Coleoptera : Coccinellidae), and the brown lacewing *Micromus* sp. (Neuroptera : Hemerobiidae), as natural enemies of *T. kahawaluokalani*, highlighting a local promising way for biocontrol of this pest. These findings are a new record for this insect pest from Gabon and Africa.

Comment [LC4]: Among or amongst?

Comment [LC5]: Capitalise months of the year

Comment [LC6]: Check spelling

Keywords: *Tinocallis kahawaluokalani*, leaf infestation, *Lagerstroemia indica*, Natural enemies, biocontrol

Comment [LC7]: Check spelling

Comment [LC8]: Check spelling

1. INTRODUCTION

Ornamental plants are widely present in all of the human artificially green spaces because of their beautiful flowers and ability to be managed in various aesthetic shapes within home and public gardens. Some of them are frequently attacked by insect pest species comprising such as aphids (Hemiptera : Aphididae) [1]. The crape myrtles, represented by *Lagerstroemia* spp (Lythraceae) are among the commonly woody plants cultivated for their ornamental attractiveness [2]. The major insect pest species reported as infesting this plant belongs to the sap sucking group, which major foliage pest is the Crapemyrtle aphid (CMA) *Tinocallis kahawaluokalani* Kirkaldy (Hemiptera : Aphididae) [1,3, 4, 5], followed by the Crapemyrtle Bark Scale *Acanthococcus Lagerstroemiae* (Kuwana) (Hemiptera : Eriococcidae) [6]. The main damages resulting from their attacks consist of black sooty mold on leaves due to their honey-dew excretions [6]. Heavy infestations may lead to aesthetic alteration and reduce visual appeal and lead to leaf drop. The plant health could decline in cases of heavy attacks of the CMA [5]. Further works on this aphid species reported its occurrences in some Asian countries as China, Japan, Thailand [2], India [7], Taiwan, some of North America as USA [4, 6], Hawaii [2], Central America as Mexico [6], Latin America as Argentina [8], Brazil [9], Eastern South of Europe [10,11] and Middle East as Iran [12]. For long time the CMA *T. kahawaluokalani* was an exotic pest species for Africa, for it has only been cited in Cameroon [13] but not in any other region nor country of this continent. The CMA impact and role in this continent are not yet known, especially in favourable condition areas. This first report in Gabon aims to highlight main knowledge on its distribution and some of its local natural predators.

Comment [LC9]: CMA acronym means they are individual words right? so please space out the words.

Comment [LC10]: Roles and impact are not known yet you call them exotic pests?

Comment [LC11]: Grammar should be properly structured

Comment [LC12]: Rephrase please

2. MATERIALS AND METHODS

Field surveys were conducted within shrubs of cultivated *Lagerstroemia indica* (L.) Pers. (Lythraceae) from avenue sides from 30 sampling sites of respective localities of Libreville, Lambarene, Mouila, Franceville and Mitzic, from august 2019 to October 2021 (Table 1).

Comment [LC13]: Statement no clear

Table1. Sampling sites of survey from ~~august~~August, 2019 to ~~october~~October, 2021, in Gabon.

No.	Locality (Province)	City location	Coordinates (latitude, longitude)
1	Libreville (Estuaire)	Carrefour Démocratie	0°25'33"N ; 9°27'24"E
2		Carrefour Awendje	0°23'12"N ; 9°28'26"E
3		Carrefour Sogatol	0°22'38"N ; 9°28'19"E
4		Oloumi	0°22'26"N ; 9°27'49"E
5		Nzeng Ayong	0°25'46"N ; 9°29'06"E
6	Franceville (Haut-Ogooué)	Yéné	1°38'24"S ; 13°34'21"E
7		Gare ferroviaire	1°37'54"S ; 13°34'49"E
8		Carrefour Sogafric	1°37'51"S ; 13°35'11"E
9		Echangeur	1°37'46"S ; 13°34'57"E
10		Avenue tribunal	1°37'35"S ; 13°36'00"E
11		Hôtel Evoula Palace	1°38'28"S ; 13°36'17"E
12		Ngobounda	1°37'20"S ; 13°36'09"E
13		Potos	1°37'54"S ; 13°34'49"E
14		Ngoungoulou	1°36'24"S ; 13°36'13"E
15		Mega mall	1°35'43"S ; 13°36'13"E
16		Djamiti	1°35'43"S ; 13°36'13"E
17		Ondzei	1°37'28"S ; 13°36'50"E
18		Ongali	1°37'19"S ; 13°36'55"E
19		IDA**	1°38'09"S ; 13°34'19"E
20		Makana	1°38'02"S ; 13°33'34"E
21	USTM	1°38'24"S ; 13°34'21"E	
22	Mouila (Ngounié)	Carrefour École St Martin	1°51'58"S ; 11°03'34"E
23		Complexe administratif	1°58'24"S ; 13°03'37"E
24		Hôtel Lac Bleu	1°52'18"S ; 11°03'34"E
25		Palais de Justice	1°52'05"S ; 11°03'25"E
26		Place indépendance	1°51'54"S ; 11°03'25"E
27	Lambaréné (Moyen-Ogooué)	Lycée Adiwa	0°41'31"S ; 10°13'42"E
28		Hôtel de Ville	0°42'30"S ; 10°13'21"E
29	Mitzic (Woleu Ntem)	Hôpital Général	0°46'31"N ; 13°34'21"E
30		Village Afia	1°04'12"S ; 11°41'31"E

With the purpose of recording the CMA occurrence on other host plants, complementary observations were carried out on *Lagerstroemia speciosa* (L) Pers and *Lawsonia alba* L. (Lythraceae) at Franceville, Okondja and Onguia village, because literature reported these species as other host plants of CMA [5,7]. All the visible trees were examined for observation of the CMA *T. kahawaluokalani* and relate predators. All the infested plants were taken into consideration basing on presence of the ~~crapemyrtle~~ aphid and predators, and the aphids infestation levels were established on five twigs per tree, and per site using following basic formula :

$$\text{Infestation} = \frac{\text{Number of infested stools}}{\text{Total of infested stools}} \times 100 [\%]$$

Comment [LC14]: Edit

After some authors [21], since the total number of stools in a sampling unit was 100, any stool infested with the aphid species was counted as 1% infestation.

Five twigs were randomly chosen per tree, and aphids and enemies were counted on five leaves from tip of each twig, following OILB/SROP recommended method [14] and used in this area [15]. The CMA colonies Abundance rate on five leaves per twig (ARFLT) were evaluated from a free scale of 1 to 10, 10 to 30 (weak), 30 to 100, and 100 to 200 individual insects, corresponding respectively to very weak, weak, middle and high levels of aphid infestation. The impact on infested twigs was observed according literature levels pointing out honey-dew, black sooty mold and plant decline [5].

The earliest aphid samples were collected with the help of fine paintbrush and preserved in 70% ethyl alcohol in 15ml glass vials for later identification at laboratories, as well for some larvae or adults of natural enemies kept in rearing plastic boxes on twigs bearing their preys (CMA). To confirm the field direct identification, the collected insects (CMA and predators) from Franceville, Lambarene, Mitzic and Mouila were brought at the Crop Plants Protection Laboratory (CPPL) of the Institut National Supérieur d'Agronomie et Biotechnologies (INSAB) from the University of Sciences and Technologies of Masuku (USTM) (Haut-Ogooué Province), whereas those collected from Libreville were analyzed at the Institut des Recherches Agronomiques et Forestières (IRAF) from the Centre National de la Recherche Scientifique et Technologique (CNAREST) (Estuaire Province).




Figure 1. *Tinocallis kahawaluokalani* observed in Gabon : heavy infestation of the CMA on a young leaf of *Largestroemia indica* showing a green syrphid larva (a), upper side of leaves covered with black sooty mold growing of aphid honeydew (b), nymph (c) and adult (d), typical tubercles on abdomen (e).

The hoverfly, coccinellid and the brown lacewing larvae observed feeding on the aphid *T. kahawaluokalani* were collected and reared into plastic rearing vials at the CPPL of INSAB still the imago hatching, for identification. All the CMA and related natural predators were identified following systematic keys provided by some available literature references [4,5,9, 23, 24, 25, 26]. Real nature and manifesting pictures (under a Binocular stereoscopic microscope Leica SZB 200) were taken with a Techno park4® camera.

Descriptive statistics, such as frequency tables, charts, simple means and percentages were used to describe and present data generated.

3. RESULTS AND DISCUSSION

3.1 The pest prevalence

The assessment of the *crapemertyle* aphid throughout 30 sites of five Gabonese localities showed presence of *Tinocallis kahawaluokalani* (Fig. 1) on 75% of *Largestroemia indica* of all the investigated provinces (table 2). The major observed *crapemertyle* were highly infested, as for the western province (Estuaire), the center (Moyen-Ogooué and Ngounié), the ~~eastern~~-Eastern (Haut-Ogooué) and the ~~northern~~-Northern (Woleu-Ntem). The insect was present in all the sampling regions of the country. The earliest infestations were recorded on ~~august~~-August 2019 at Libreville (infested trees were 85,71%), ~~july~~-July 2020 at Lamabaréné (60%) and Mouila (83,33%), whereas latest records occurred on ~~october~~-October 2021 at Franceville (84,09%) and Mitzic (71,43 %) (table 2). Observations on potential hosts at Franceville and surrounding localities, as ten individuals of *Largestroemia speciosa* (L) Pers and twelve *Lawsonia alba* L. (Lythraceae) showed any CMA infestation, certainly because the CMA host preference in this town goes exclusively on *L. indica*. Weak, heavy and very heavy infestation were recorded, inducing presence of honey-dew, black sooty mold, and plant decline (table 2). Seven beneficial predaceous species were also recorded (table 3) from Franceville, Libreville and Mitzic.

Comment [LC15]: Check all and be consistent

Formatted: Font: Not Italic

Comment [LC16]: Italicize all scientific name

Formatted: Font: Italic

Table 2. Occurrence and distribution of *Tinocallis kahawaluokalani* Kirkaldy (1906)(Hemiptera : Aphididae) (on 5 twigs per tree)

Locality (Province)	City location	Tree samples		Aphidpest		
		N	% infested trees	ARFLT*	Infestation level	Damage level
Libreville (Estuaire)	Carrefour Démocratie	2	100	500 - 1000	+++++	Much honey-dew, Leaves drop
	Carrefour Awendje	8	100	30 - 100	+++	Little honey-dew
	Carrefour Sogatol	3	100	30 - 100	+++	Little honey-dew
	Oloumi	3	100	30 - 100	+++	Little honey-dew
	Nzeng Ayong	2	0	0	-	No damage
	Yéné	3	0	0	-	No damage
Franceville (Haut-Ogooué)	Gare ferroviaire	2	100	500 - 1000	+++++	Black sooty mold
	Carrefour Sogafric	2	0	0	-	Green leaves
	Echangeur	7	62,5	150 - 500	++++	Much honey-dew, Little honey-dew
	Avenue tribunal	1	100	50 - 150	+++	Little honey-dew
	Evoula Palace	12	100	50 - 150	+++	Little honey-dew
	Ngobounda	4	100	30 - 100	+++	Honey-dew
	Potos	1	100	50 - 150	+++	Very little honey-dew
	Ngougoulou	2	100	30 - 100	+++	Honey-dew
	Mega mall	2	0	0	-	No damage
	Djamiti	3	100	5 - 50	++	Very little honey-dew
	Ondzei	1	100	30 - 100	+++	Little honey-dew, Black sooty mold
	Ongali	3	100	150 - 500	++++	Black sooty mold
	IDA**	3	0	0	-	No damage
	Makana	3	66,67	5 - 50	++	No damage
USTM	10	66,67	1-10	+	No damage	
Mouila (Ngounié)	Ecole St Martin	3	66,67	30 -100	+++	Little honey-dew
	Complexe administratif	2	100	30 - 100	+++	Little honey-dew
	Hôtel Lac Bleu	3	66,67	30 - 100	+++	Little honey-dew
	Palais de Justice	2	100	30 - 100	+++	Little honey-dew
	Place indépendance	2	100	30 - 100	+++	Little honey-dew
Lambaréné (Moyen-Ogooué)	Lycée Adiwa	2	0	0	-	No damage
	Hôtel de Ville	3	100	30 - 100	+++	Little honey-dew
Mitzic (Woleu Ntem)	Hôpital Général	5	100	50 - 150	+++	Black sooty mold
	Village Afia	2	0	0	-	No damage
TOTAL	30	104	75			

Legend : *ARFLT = Abundance rate on five leaves per twig ; **IDA = Inspection Déléguée d'Académie

3.2 The predators' occurrence

The predominant hoverfly species were respectively *Asarkina* sp present on 20 to 25% of twigs, *Ischidon aegyptius* (50 to 100%), and *Paragus borbonicus* (10 to 100%) (Fig. 2) ; coccinellid species were *Platynaspis capicola* (20%), *Chilochrus nigrinus* (20 to 80%) and *Scymnus interruptus* (20%), and the hemerobiid brown lacewing *Micromus* sp. (8 to 20%)(Fig. 3). In addition, numerous individuals of a *Deræocoris* sp. (Hemiptera : Miridae) and *Orius* sp. (Hemiptera : Anthocoridae) were found associated with the CMA populations.

Table3. Occurrence of beneficials (on 5 twigs per tree) according to the pest aphid level

Localities (Province)	City location	Pestaphid <i>Tirocallis kahawaluokalani</i> Kirkaldy(1906) (Hemiptera : aphididae)	Beneficials'insects (natural enemies of CMA)												Associated insects					
			Asarkina sp (Diptera : Syrphidae)		Ischidon aegyptiusWiedemann (1830) (Diptera : Syrphidae)		Paragus borbonicus(Macquart (1482) (Diptera : Syrphidae)		Platynaspis capicolaCrotch (1874) (Coleoptera : coccinellidae)		Chilochorus nigrinusFabricus (1798) (Coleoptera : coccinellidae)		Scymnus interruptusGoeze(1885) (Coleoptera : coccinellidae)		Micromus sp (Neuroptera : Hemerobiidae)		Deræocoris sp (Hemiptera : Miridae)		Orius sp (Hemiptera : Anthocoridae)	
			n	% tb*	n	% tb	n	% tb	n	% tb	n	% tb	n	% tb	n	% tb	n	% tb	n	% tb
Libreville(Estuaire)	Carrefour Démocratie	+++++	-	-	6	80	-	-	-	-	-	-	-	-	-	-	-	-	-	
Franceville(Haut- Ogooué)	Gare ferroviaire	+++++	1	20	10	100	1	10	-	-	1	20	-	-	-	-	3	30	3	20
	Echangeur	++++	2	25	9	60	5	100	1	20	-	-	3	20	2	8	-	-	-	-
	Ngobounda	+++	-	-	7	50	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	Quartier sable	+++	-	-	6	80	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mitzié (Woleu Ntem)	Ongali	++++	-	-	6	80	-	-	-	-	11	80	-	-	-	-	-	-	-	-
	Hôpital Général	+++	-	-	-	-	-	-	-	-	-	-	-	-	1	20	-	-	-	-
TOTAL			3	23,08	44	69	6	15	1	20	12	63,16	3	20	3	12	3	30	3	20

Legend : *tb= twigs bearing beneficial or associated insects



Figure 2. Hoverflies (Diptera : Syrphidae) preying in CMA : *Asarkina* sp. (a : larval instar, b : pupae, c : adult) ; *Ischidon aegyptius* (d : larval instar, e, pupae, f : male, g : female), *Paragus borbonicus* (h : larval instar, i : pupae, j : laying female, k : male).



Figure 3. Coccinellid and Hemrobiid predators: *Chilocorus nigritus* (a : larva, b : nymph, c : adulte on twig, d : 3.5x magnifying adult), *Platynaspis capicola* (e : minute and 5x magnifying adult), *Scymnus interruptus* (f : adult), *Micromus* sp. (g : nympe, h : adult)

3.4 Discussion

Among prospected localities, Franceville revealed highest natural enemies abundance and diversity because of deeper sampling effort. This study is a first time report of the *T. kahawaluokalani* occurrence and relate aphidophagous species in Gabon, and certainly a second record in Africa, because within [litteratures-litteratures](#) studying CMA, only one mention notices observations of the CMA at Younde and Douala in Cameroon [13], at our knowledge. Any other study was found emphasizing on this pest occurrence in Africa. Moreover, the latest entomofaunistic studies in inhabiting localities of Gabon [15, 18] did not report any infestation of the CMA. Nevertheless, monitoring of this pest at Franceville provided unsuccessful results from 2019 to 2020. The record of CMA that very 2021 year is to indicate its newly arrival and expansion in this locality. Furthermore, the introduction of the CMA in Gabon seems to be recent, and could have been occurred from Cameroon, or another neighboring or faraway country, in the context of global fast world and increasing exchanges. Heavy infestations on shrubs of the train station site (Gare ferroviaire) could lead to suspect the train as having been a transport mean that facilitates that insect occurrence and outspread from Libreville to Franceville. Elsewhere, within the local enemies observed, the syrphid species were the major predator beneficial group, frequently being predated on the crapemertyle colonies, whereas the other predators occurred occasionally in some sites. The species *Orius* sp, possibly *Orius*

Comment [LC17]: spelling

Comment [LC18]: spelling error

insidiosus, observed on blossoming crape myrtles is an interesting case needing deeper scientific investigations because, as known as predator of the crape myrtle aphid in USA [19]. This presumable predator, recognized as an omnivore predeflecting small hemipterans and flower resources, with behaviour to aggregate in areas with high concentrations of pollen [20]. Their synergist aphidophagous predation on CMA allowed to record very few presence of the Aphid one month later after first observations. The study did not find momified aphids, as to confirm unsuccessful assessment from former authors pointing out the lack of parasitoids attacking CMA [13]. Some of these predators previously had been recorded by local former authors [15, 18], and are potential biological candidates for mass rearing to reinforce their effectiveness as an IMP promising way. Considering current observations, efforts are to be made to develop deeper knowledge and efficient strategies to control the CMA in this environment.

4. CONCLUSIONS

The Japan aphid is well established and spreading in Gabon. From the Estuaire ~~western~~-Western province, the Ngounie center province, the Haut-Ogooué ~~eastern~~-Eastern province, and the Woleu-Ntem ~~northern~~-Northern province, the crape myrtle aphid occurrence is recorded, with local natural enemies belonging to syrphids, coccinellids, hemerobiids and minute pirate bugs (anthocorid). Regular monitoring is recommended and control measures might take in account rearing, release and preservation of natural enemies of *T. kahawaluokalani*. Nevertheless, since this insect pest is newly recorded in Gabon and now reported from the African continent, assessment efforts are needed to state on its economic threshold of infestation, in order to develop global management strategies against this ornamental threat.

Comment [LC19]: spelling

Comment [LC20]: check

Comment [LC21]: check spelling and rephrase

COMPETING INTERESTS

All authors disclose any financial and personal relationships with other people or organizations that could inappropriately influence their work. This work was performed without funding, consultancies or other potential involvement that could lead to conflicts of interest. Authors declare that no competing interests exist.

REFERENCES

1. Blackman RL, Eastop VF. Aphids on the World's trees. An identification and information guide. CAB International, Wallingford, Oxon, 1994.
2. Gilman E F, Watson D. G. *Lagerstroemia indica* Crape-Myrtle. Fact Sheet ST-342. Environmental Horticulture Department, Florida Cooperative Extension Service, IFAS, University of Florida, USA. 1993.
3. Baker J, Carter C C, Horn KF, Kline D, Scott J, Singletary H, Stephan D. Pests of Crape Myrtle : Key to Crape Myrtle Pests. *NC State Extension Publications* USA (published online). 1993. <https://content.ces.ncsu.edu/insect-and-related-pests-of-shrubs/pests-of-crape-myrtle>
4. Herbert J J, Mizell R, Mc Auslane H. Host preference of the Crape Myrtle Aphid (Hemiptera : Aphididae) and host suitability of Crape Myrtle cultivars. *Environ. Entomol.* 2009 ; 38 : 1155-1160.
5. Herbert J J, Mizell R. Crape Myrtle Aphid, *Tinocallis kahawaluokalani* (Kirkaldy) (Insecta : Hemiptera : Aphididae). Fact Sheet EENY365. Department of Entomology and Nematology, UF/IFAS Extension, University of Florida, USA. 2018.
6. Wang Z. Biology and Ecology of Crape Myrtle Bark Scale *Acanthococcus lagerstroemiae* (Kuwana) (Hemiptera : Eriococcidae) LSU Master's Theses. 2017.
7. Agarwala BK, Mahapatra SK, Ghosh AK. Description of sexual morphs of *Tinocallis kahawaluokalani* (Kirkaldy) (Homoptera: Aphididae) from India. *Entomon.* 1989 ; 14 : 273-274.
8. Alfonsina S. Aphididae (Hemiptera) on ornamental plants in Córdoba (Argentina). —*Rev. Soc. Entomol. Arg.* 2008 ; 67(1-2) : 49-56.
9. Lúcia A, Peronti BG, Sousa-Silva CR. Aphids (Hemiptera : Aphidoidea) of ornamental plants from São Carlos, São Paulo state, Brazil. *Rev. Biol. Trop.* 2002 ; 50(1) : 137-144.
10. Tsitsipis JA, Nikos KI, John MT, Dionyssios LP, Apostolos AD, Ioanna G. A contribution to the aphid fauna of Greece. *Bull. Insectology.* 2007 ; 60(1) : 31-38.

11. Schoeny A, Gognalons P. Data on winged insect dynamics in melon crops in southeastern France. Data in Brief, Elsevier, 29. 2020 ;105132pp. <https://doi.org/10.1016/j.dib.2020.105132>.
12. Chitgar MG. First report of the aphid *Tinocallis kahawaluokalani* (Hemiptera : Aphididae) from Iran. J. E. S. I. 2017 ; 37(3) :369-370.
13. Mizell R F, Bennett F D, Reed DK. Unsuccessful search for parasites of the crapemyrtle aphid, *Tinocallis kahawaluokalani* (Homoptera : Aphididae). Fla. Entomol. 2002 ; 85(3) : 521-523.
14. OILB/SROP. Contrôles, seuils, et indications pour la lutte (Pommier III). In Acta (Eds.). Contrôles périodiques en verger. Acta, Paris, France. 1977.French
15. Poligui RN. Étude de l'entomofaune au sein d'associations culturales comprenant le Safoutier (*Dacryodes edulis*) dans le Haut-Ogooué (Gabon). Thèse de doctorat. Gembloux université de Liège Belgique. 2014. French
16. van den Berg H, Cock MJW. African bollworm and its natural enemies in Kenya. CABI Africa Regional Centre, Second Edition. 2000.
17. Ssymank AA. contribution to the Syrphidae (Diptera) fauna of Cameroon, with a preliminary checklist of the family. Afr. Invertebr. 2012 ; 53 (1) : 249–266.
18. Bayendi Loudit SM, Poligui RN, Verheggen F, Francis F. Occurrence of Aphids and their predators within vegetable crops in peri-urban areas in Libreville (Gabon). Commun. Agric. Appl. Biol. Sci. 2017 ; 82(2) :199–205.
19. Parsons SE, Frank SD. Urban tree pests and natural enemies respond to habitat at different spatial scales. J Urban Econ. 2019 ;5(1) :1–15.
20. Isehour D J, Marston N L 'Seasonal Cycles of *Orius insidiosus* (Hemiptera : Anthorcoridae) in Missouri Soybeans', Journal of Kansas Entomological Society .1981 ; 54 : 129–42.