

# Taylor Flap Reconstruction and its Impact on Healing and Psychological Comfort: [A Therapeutic Challenge Incurred Within a Developing Country](#)

## **Abstract**

**Purpose:** The aim of this study was to assess the feasibility and then report the short, medium and long-term results of Taylor flap reconstructions after abdomino-perineal amputation extended to the posterior wall of the vagina, taken by the tumor mass.

The abdomino-perineal amputation is burdened with significant morbidity closely related to the inevitable large perineal defect due to wide excision.

Hence the importance of pelvipereineal filling techniques to facilitate perineal healing and to ensure a better quality of life, especially sexual, in women after posterior colectomy, instead of a mutilating posterior pelvic exenteration.

We describe and analyze below two cases of perineal reconstruction by modified Taylor flap.

During 2020, two of our patients underwent rectus abdominis musculocutaneous flap reconstruction pedicled on the inferior epigastric vessels and with a transverse skin paddle, for abdominoperineal amputations performed for malignant melanoma of the anal canal (Fig.1) and adenocarcinoma of the lower rectum respectively.

Our two patients underwent a posterior colectomy associated with abdominoperineal amputation, which improved the quality of life of the patients, in particular an almost normal sexual activity in the medium and long term.

The postoperative follow-up was simple with an average hospital stay of 09 days. Abdominoperineal healing completed with a median of 19.5 days.

Unremarkable follow-up with a follow-up of one year.

The realization of modified Taylor flaps allowed a complete and fast perineal healing, and a psychological comfort in connection with a satisfactory sexual activity in both patients.

### **Keywords:**

Taylor flap; Abdominal-perineal amputation; posterior colpectomy; Melanoma; Rectal Cancer; Rectus Abdominus

### **Introduction:**

The APA (**AbdominoPerineal Amputation**) is the gold standard procedure for lower rectal cancers.

In the latest published series, postoperative complication rates without perineal reconstruction vary from 35 to 66% [1].

Since 2002, we have resorted to plastic surgery using musculocutaneous flaps to overcome the postoperative complications inherent to the extent of the excision and the difficulty of closing the perineal defect, in order to accelerate healing, limit the aesthetic after-effects and preserve sexual activity, particularly in women, by preserving the lower genital tract, which results in a considerable improvement in the quality of life of the patients.

We report two cases of perineal reconstruction using a modified Taylor flap. We analyzed the postoperative course, specific complications, and results of this surgery.

### **Case presentation**

We report two cases of lower rectal cancer treated during the year 2020.

They were two women in their 50s with a WHO (world health organization) score of 0, preserved nutritional status, and no surgical history that could compromise vascular flow tributary to the lower epigastric vessels.

One of them had a malignant melanoma of the anal canal (Fig.1), while the other patient had a well-differentiated adenocarcinoma of the lower rectum (Fig.2), treated with APA after concomitant radio-chemotherapy.

In both patients, the abdominoperineal amputation extended to the lower 2/3 of the posterior vaginal wall to obtain an R0 resection and the reconstruction required an extension of the skin flap to ensure the plasticity of the posterior vaginal wall.

Thus, the reconstruction began with the removal of the musculocutaneous flap with an average size of eight centimeters in width and twenty-one centimeters in length, initially traced (Fig. 3) along a transverse axis at the height of the navel on the right side, giving way to the definitive colostomy on the left at the end of the operation,

Once the flap has been mobilized (Fig. 4), it is tilted downwards by a 180° rotation intra-abdominally to fill in the perineum. The proximal musculocutaneous part is fixed in two planes to cover the perineal region and the distal part of the skin paddle fills the posterior aspect of the vagina, without having to open and section the broad and round ligament.

Reconstruction is performed as soon as the abdominoperineal amputation is completed. The distal colonic stump concerned by the stoma is left to stand, hemmed in to the skin at the last stage of the operation.

Finally, the flap will be fixed to the perineal floor by separate stitches going up the posterior surface of the vagina (Fig. 5).

Furthermore, the fascia is closed simply by bringing the fascial edges of the anterior rectus fascia together, without any prosthetic reinforcement, in order to reduce the risk of sepsis, which is increased by the proximity of the definitive left iliac colostomy, and taking care not to plicate or injure the feeder pedicle of the flap.

In performing the perineo-vaginal plasty, the objective is to evaluate the duration incurred for optimal healing and the quality of our patient's lives, particularly sexual quality, by means of close clinical monitoring, initially weekly for one month, then every 15 days during the first trimester, and then monthly, with a current follow-up of 8 months.

Thus, healing is declared when the 2 sites; abdominal donor and perineo-vaginal receptor have completely healed.

The quality of sexual life will be judged by the extent of the period of coital arrest, relating the degree of vaginal dryness and dyspareunia.

### **Discussion:**

In the hospital, the patient operated on a melanoma of the anal canal had presented phlyctenoid bullae on the proximal cutaneous side of the flap which have collapsed by application of fatty plaster changed daily. Apart from this, both patients progressed well without the emergence of any complication specific to the excision surgery or specific to the plastic surgery procedure.

The length of hospitalization for both patients was 10 days.

Complete perineal healing was obtained at the end of the 3rd week in both patients.

With an average follow-up of one year, no complications arose at either the donor or recipient sites.

Coital resumption started progressively from the second week, just after complete perineo-vaginal healing (Fig.6); delayed by one week, following dyspareunia in the patient who underwent excision for anal melanoma.

Perineal healing and psychological ease, particularly sexual ease, represent the essential goal in patients who have undergone APA associated with posterior colectomy [3].

In 1983, Taylor described a musculocutaneous flap of the rectus abdominis pedicled on the deep inferior epigastric vessels [2]. The skin racket was initially traced along an oblique axis that follows a line from the umbilicus to the tip of the scapula.

The technical variant eventually adopted in our two patients consisted in harvesting a paddle with a large transverse axis. This transverse rectus

abdominis myocutaneous flap (TRAM flap) with an inferior pedicle is mainly indicated in slim individuals.

This option is one of the safest and most effective solutions for the coverage and filling of pelvic-perineal tissue loss [4].

Direct closure with omentoplasty as one of the alternatives to the Taylor flap had a significantly higher complication rate, healing time and occurrence of perineal hernias (48.9% vs. 26.8% and 117 d vs. 18.7 d; 15.4% vs. 0% respectively in the series of Lefèvre et al.) than with a Taylor flap, with an almost identical incidence of occurrence of abdominal eventrations [5].

During pelvic-perineal plasties, we can also resort to the use of other flaps such as the Gracilis flap, at the cost of a questionable vascular reliability and a much smaller size compared to the Taylor flap, which most often results in samples being taken from both sides, or even by combining several at the same time, especially at the time of enlarged reconstructions (example: reconstruction by using petal flap of the lotus and the pudendal fascio-cutaneous flap) which are complex and difficult techniques, and which can be complicated by partial or total necrosis of the flaps, of ischemic or suppurative origin in 25% of the cases; elucidated during innumerable series of studies [6-7].

Thus, the Taylor flap represents the reference technique for covering large pelvic-perineal losses of substance, particularly in enlarged APAs. The other plasty techniques are considered as a second choice, or in case of impossibility of harvesting the pedicled rectus flap (anterior laparotomies complicated by wide parietal ventrations) [8].

### **Conclusion:**

Vascular reliability and the large volume of the musculocutaneous flap are key assets to ensure optimal filling of the pelvic-perineal defect associated with pelvic exenteration and abdomino-perineal amputations.

According to previous published series and our two reported cases [10], the Taylor flap is the technique of choice with a much reduced healing time, in all patients who have undergone PAA for anal or lower rectal cancer without

worsening the morbidity figures, related to the radical surgery, with a benefit on psychological comfort and sexual quality of life.

**Author contributions** : All authors contributed to the study concept and design, data interpretation, drafting, final approval, and accountability for all aspects of the work.

**Funding** : This study was not funded by any grant.

**Conflict of interest**: The authors declare that they have no conflict of interest.

### **Consent**

**As per international standard or university standard, patient (s) written consent has been collected and preserved by the author(s).**

### **Ethical Approval:**

**As per international standard or university standard written ethical approval has been collected and preserved by the author(s).**

### **References:**

1. Sinna R, Qassemyar Q, Benhaim T, Lauzanne P, Mauvais F. Perforator flaps: a new option in perineal reconstruction. Review article. J Plast Reconstr Aesthet Surg 2010;63(11):e766-74.
2. Taylor GI, Corlett R, Boyd JB. The extended deep inferior epigastric flap: a clinical technique. Plast Reconstr Surg 1983;72:751-65.
3. Lawhead Jr RA, Clark DG, Smith DH, Pierce VK, Lewis Jr JL. Pelvic exenteration for recurrent or persistent gynecologic malignancies: a 10-year

review of the Memorial Sloan-Kettering Cancer Center experience (1972—1981). *Gynecol Oncol* 1989;33(3):279—82.

4. Harrison LB, Enker WE, Anderson LL. High dose rate intraoperative radiation therapy for colorectal cancer. *Oncology*. 1995; 9:737-741.
5. Lefèvre JH, Corte H, Tiret E, Boccara D, Chaouat M, Touboul E, et al. Abdomino-perineal resection for squamous cell anal carcinoma: survival and risk factors for recurrence. *Ann Surg Oncol* 2012;19(13):4186-92.  
<http://dx.doi.org/10.1245/s10434-012-2485-1>.
6. Wee JT, Joseph VT. A new technique of vaginal reconstruction using neurovascular pudendal-thigh flaps: a preliminary report. *Plast Reconstr Surg* 1989; 83:701-9.
7. Bell SW1, Dehni N, Chaouat M, Lifante JC, Parc R, Tiret E. Primary rectus abdominis myocutaneous flap for repair of perineal and vaginal defects after extended abdomino-perineal resection. *Br J Surg* 2005;92(4):482—6.
8. Knol AC, Hage JJ. The infragluteal skin flap: a new option for reconstruction in the perineogenital area. *Plast Reconstr Surg* 1997;99(7):1954—9.
9. Leclère FM, Mordon S, Ramboaniaina S, Schoofs M. Breast reconstruction with a free DIEP flap complicated by spontaneous rupture of internal mammary artery. *Ann Chir Plast Esthet* 2010;55(6):593—6.  
<http://dx.doi.org/10.1016/j.anplas.2009.11.018> [Epub 2010 Jun 18].
10. Smith HO, Genesen MC, Runowicz CD, Goldberg GL. The rectus abdominis myocutaneous flap: modifications, complications, and sexual function. *Cancer* 1998;83(3):510—20.