

Original Research Article

Effect of feeding diabetic rats on breadsticks fortified with psyllium seed powder

Abstract

This study aimed to investigate the effect of breadsticks produced from wheat flour (72% extract) and psyllium seeds Powder (PSP) on chemical, and sensory properties. Effect of feeding diabetic rats with breadsticks on blood glucose, biochemical parameters, organs weight lipid profile, kidney, and liver functions were also assessed. Breadsticks were prepared by substituting wheat flour by 3, 6, and 9% PSP. Results revealed that PSP contained high values 15.82% of crude protein, 4.03% ash, 5.56% ether extract 26% crude fiber, 6.50% soluble fiber and 20.02% insoluble fiber. Biological results showed a significant improvement ($p < 0.05$) in values of blood glucose, lipid profile (HDL- LDL- VLDL, triglycerides and total cholesterol in serum for rats fed on bread sticks supplemented with PSP at extent 3,6,9% comparing to positive group. Also, feeding diabetic rats on PSP clarified a remarkable improvement in ALT, AST, urea, uric acid, and creatinine comparing to normal rats. It could be concluded that bread sticks prepared using PSP might help for the prevention and treatment of diabetic rats, also, could be beneficial for people with diabetes. This research recommends using PSP with wheat flour on a commercial scale for preparation of backing products to combat diabetic in Egypt.

Key words: *Diabetes, Plantago ovate seeds, bread sticks.*

1. Introduction

Type 2 diabetes mellitus is considered as a metabolic disorder which is characterized by β -cell dysfunction and high blood glucose level in the context of the relative insulin insufficiency

(Kumar *et al.*, 2005). Among the metabolic conditions that type 2 diabetes is accompanied by oxidative stress (excess free radicals and a malfunction in the body's antioxidant defense system), poor lipid and lipoprotein metabolism, hypertension, vascular endothelial dysfunction, and subclinical inflammation (Bekyarova *et al.*, 2007). Metabolic diseases caused microvascular and macro vascular consequences, (Constantino *et al.*, 2013). According to (Modi, 2007). The main modalities of treatment for type 2 diabetes are dietary and activity changes, insulin replacement therapy, insulin secretagogues, insulin sensitizers, and alpha-glucosidase inhibitors. To regulate their blood sugar and avoid complications from diabetes, patients must take numerous medicine combinations. But some negative effects of these medications include hypoglycemia, gastrointestinal distress, and lactic acid poisoning. Therefore, it is preferable for patients to utilize a plant medicine that is effective at reducing the symptoms of diabetes. (Grant *et al.*, 2013).

Plantagoovata (psyllium) belongs to the Plantaginaceae family. The plant's husk and seeds have significant therapeutic and economic value. Psyllium seeds are used in traditional medicine for a variety of medical purposes (Tewari *et al.*, 2014). Psyllium has a characteristic fiber got from psyllium seed. It is a highly fanned arabinoxylan polysaccharide with an extremely high water retention and gelling limit. The soluble base extractable polysaccharides of psyllium's gel-framing part are constituted of arabinose, xylose, and traces of other sugars (Fischer *et al.*, 2017).

Psyllium seeds are categorized as a mucilaginous fiber because of their exceptional capacity to gel in water. The plantagopsyllium is a significant medicinal plant that contains a variety of chemicals, including flavonoids, alkaloids, terpenoids, phenolic compounds (derivatives of caffeine), and vitamin C. Plantago plants contain significant levels of phenols, and it's possible that these phenolic substances are what give the plant its strong antioxidant properties. Although phenolic chemicals, particularly flavonoids, have antioxidant properties, they cannot completely prevent all mutations brought on by mutagenic agents. (Haddadian *et al.*, 2014). Psyllium is a viable option for usage in the nutraceutical business since it includes nutritional antioxidants, flavonoids, polyunsaturated fatty acids (PUFAs), including essential fatty acids (Omega-3 and Omega-6 fatty acids), sulfur-containing amino acids, and metabolites with bioactivities. Psyllium may help people with diabetes better control their blood sugar and cholesterol levels, according to several studies. In investigations on both animals and people, psyllium consumption has been demonstrated to enhance glucose and insulin responses (Singh, 2007). Due to the binding of bile acids in the intestinal lumen, psyllium has a hypolipidemic effect that decreases serum cholesterol levels and lowers the risk of coronary heart disease. (Van Rosendaal *et al.*, 2004). This work confirms the great importance of applied science in bakeries foods (Elhadidy, 2009; Elhadidy, 2014; El-Dreny and El-Hadidy, 2018; El-Hadidy, 2020; El-Hadidy *et al.*, 2020; El Hadidy and Rizk, 2020; El-Hadidy and El-Dreny, 2020; El-Hadidy *et al.*, 2022).

The goal of the current study was to determine whether psyllium seeds have any anti-diabetic effects in diabetic rats that have been fed a high-fat diet with streptozotocin.

2. Materials and Methods

2.1. Materials

Psyllium seed, wheat flour, and other ingredients necessary to make breadsticks, including salt, sugar, dry yeast, and fat were purchased from the market in Menoufia, Egypt. Analytical grade chemicals and solvents from the Sigma Company were employed throughout the study, rats were purchased from animal house of food technology research Institute.

2.2. Methods

2.2.1. Breadsticks making:

Making sticks with psyllium seed powder using the listed ingredients in Table 1. Psyllium seed powder was used in place of base flour at varying levels (3, 6 and 9 percent). The control dough was made with only wheat flour (72 percent extraction).

Table (1): The basic (control) formula used in the preparation of breadsticks

Ingredients(g)	Blend (1)	Blend (2)	Blend (3)	Blend (4)
Wheat flour 72%	100	97	94	91
psyllium seeds%	-	3	6	9
Sugar(g)	8	8	8	8
Fat (oil) (g)	10	10	10	10
Salt(g)	0.5	0.5	0.5	0.5
Dry yeast(g)	0.6	0.6	0.6	0.6

Dough preparation. Each type of flour was combined with warm water, oil, sugar, fat, salt, and dry yeast. The materials were then well mixed by hand. The dough was allowed to ferment for 30 minutes at $(30 \pm 2 \text{ }^\circ\text{C})$. After cutting the dough into pieces, it was given a 10 minute rest. After shaping the fermented pieces into their final form, they fermented for half an hour at $30 \text{ }^\circ\text{C}$ and 90% relative humidity. The fermented dough baked for 30 minutes at $170 \text{ }^\circ\text{C}$ (Ballester-Sánchez, 2019).

2.2.2. Chemical analysis:

Psyllium seed flour was analyzed for crude protein, ash, crude ether extract and crude fiber according to the methods of AOAC (2005). Available carbohydrates were calculated by difference.

% Available carbohydrates = $100 - (\text{protein} + \text{ash} + \text{ether extract} + \text{crude fiber})$ Calorie value were calculated agreeing to the Atwater system FAO (2000).

Calorie value (kcal/100g) = $(\% \text{ carbohydrate} \times 4) + (\% \text{ protein} \times 4) + (\% \text{ fat} \times 9)$.

2.2.3. Determination of soluble and insoluble dietary fiber:

Soluble and insoluble fiber were determined according to the method described by ASP *et al.*, (1983).

2.2.4. Sensory evaluation:

Twenty panelists conducted an organoleptic examination of the investigated breadsticks with and without psyllium seed flour to assess flavor, texture, color, taste, and acceptability in general. According to Renzo (1975), evaluations were done using scores ranging from 1 to 10, with exceptional (10-9), very good (8-6), fair (5-4) and not acceptable (3-2) being used.

2.3. Rats feeding:

The current experiment in the animal house of food technology used a total of thirty adult male albino rats that were weighted (145-150%). Under normal, healthy circumstances for 10 days, the Research Institute, Agric, Research, Center, Giza, Egypt, subjects were fed a typical diet (base diet), as stated in Table 2. The rats were fed a consistent diet and given free access to tap water.

Table (2): composition of the basal diet.

Ingredient	g/100g diet	Ingredient	g/100g diet
Corn starch	60	Cellulose	5
Corn oil	10	Salt mixture	4
Casein	20	Vitamin mixture	1

(Lane-Peter and Pearson, 1971)

2.3.1. Design of experiment:

Following ten days of only feeding on the standard diet as an adjustment period, a negative control group (G1) of six rats was kept and given the standard diet throughout the trial. According to **Dawson *et al.* (1986)**, 24 rats were given the intraperitoneal injection of streptozotocin (STZ) at a dose of 60 mg/kg body weight after fasting overnight. Blood samples from each rat were collected for the measurement of serum glucose after four days of injection (zero time) to ensure the development of diabetes. Diabetic rats were defined as having blood glucose levels of 250–550 mg/100 ml.

2.3.2. Hyperglycemia experiment of psyllium seed:

G1: Fed on the 50% blend1+50% standard diet. (Negative control).

G2: Fed on the 50% blend1+50% standard diet. (Positive diabetic control).

G3: Fed on the basic diet with 50% blend2+50% diet.

G4: Fed on the basic diet with 50% blend3+50% diet.

G5: Fed on the basic diet with 50% blend4+50% diet.

Each animal had ethyl ether anesthesia at the end of the experiment, which lasted six weeks. Blood was drawn from the venous plexus's eye, centrifuged for 10 minutes at 3000 rpm to extract the serum, and then frozen at 20 °C in dry, clean plastic until analysis.

2.4. Biochemical analysis:-

Kits relate to serum glucose, cholesterol (CHL), low density lipoprotein (LDL), high density lipoprotein (HDL), triglyceride (TG), serum alanine amino-transferase (ALT), serum aspartate amino-transferase (AST), Albumin, urea, creatinine, uric acid, purchased from SPINREACT Co, SPAIN 2021.

2.5. Statistical evaluation

The obtained results were analyzed statistically using Analysis of Variance (One way ANOVA) using SPSS 16.0.

3. Results

3.1. Proximate composition of psyllium seeds powder.

The chemical composition of psyllium seeds shown in Table 3. The results obtained were based on dry weight. From the tabular data, it can be seen that the crude protein content of **psyllium seeds Powder** was 15.82 %. In addition to ether extract 5.56%, crude fiber content 26.00%, ash 4.03%, available carbohydrates 47.52% , total carbohydrates 74.59% and Caloric value 303.4 (kcal/100g).(El-Hadidy,2020) reported that psyllium seeds powder comprise 16.40% crude protein,5.40% fat, 3.70%ash,48% carbohydrates,26.50 crude fiber, 6 %soluble fiber, and 20.55%insoluble fiber

Table (3): Chemical composition of psyllium seeds powder raw materials (g / 100g on dry weight basis)

Contents	psyllium seeds powder
Moisture	86.34±2.52
Crude protein %	15.82±1.22
Crude ether extract %	5.56±0.72
Ash %	4.03±0.82
Total carbohydrates %	74.59±3.32
Available carbohydrates %	47.52±2.62
Crude fiber %	26.00±1.23
Soluble fiber %	6.50±0.62
Insoluble fiber %	20.02±0.92
Caloric value (kcal/100g)	303.4±3.78

3.2. Effect of feeding bread sticks with different levels of psyllium seeds powder on body weight

Data in Table 4 reveal that initial body weight at the beginning of the experiment, ranged from 145 to 150g of all groups. While, the most effective of supplementation of psyllium seeds Powder in rats fed breadsticks on change in body weight of rats. Psyllium seeds powder supplementation improved final body weight gain (%) compared with rats fed with standard diet after 6 weeks of the experimental period ($p \leq 0.05$).

Table 4. Effect of feeding bread sticks different psyllium seeds powder on body weight gain of diabetic rats:

Groups of rats	Initial weight (g)	Final weight (g)	Change in body weight (%)
G1	148.43a ±1.87	170.36a ±1.95	14.77
G2	149.43a ±1.95	139.48d ±1.85	-6.66
G3	146.95a ±1.96	166.86c ±2.16	13.55
G4	147.10a ±1.35	167.35c ±1.49	15.02
G5	147.43a ±1.68	169.57b ±1.51	22.14a ±1.35

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

3.3. Effect of feeding bread sticks with psyllium seeds powder on organs weight:

The results in Table 5 indicate that there is a decrease in the weight of the organs of diabetic rats fed breadsticks with psyllium seeds powder compared to the control, which indicates that the diet of breadsticks psyllium seeds powder has an effective role on the organs of the body.

Table (5): Effect of feeding bread sticks with psyllium seeds powder on organs weight:

Rat Groups	Liver (g)	Kidney (g)	Heart (g)	Spleen (g)
G1	6.73bc ±0.42	1.39d ±0.05	0.58c ±0.04	0.69c ±0.07
G2	9.06a ±0.53	1.90a ±0.03	0.77a ±0.03	0.87a ±0.06
G3	7.99ab ±0.56	1.87a ±0.04	0.67ab ±0.07	0.86ab ±0.08
G4	7.45abc ±0.72	1.77b±0.05	0.65b ±0.05	0.78b±0.09
G5	6.79bc ±0.62	1.47c ±0.05	0.61b±0.06	0.61cd ±0.06

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

3.4. Blood serum glucose levels of rats fed on prepared bread sticks.

Table 6 and illustrates the blood glucose levels of negative and diabetic groups through the experimental period (6 weeks). Blood glucose levels of diabetic groups (G2) were markedly higher than that of normal negative control (G1). The results also indicate that, bread sticks contained psyllium seeds powder caused a significant decrease in blood glucose levels of the diabetic rats (G3 to G5) comparing with negative control rats (G1) and positive group fed on bread steaks contained wheat flour only (G2).

Table 6. Effect of feeding bread sticks with different psyllium seeds powder on blood glucose level of diabetic rats.

Rat groups	After 3 days mg/dl	After 2weeks mg/dl	After 4 weeks mg/dl	After 6 weeks mg/dl
G1	105.33^b ±1.54	105.44^e ±1.68	103.78^e ±1.17	100.78^e ±1.57
G2	304.66^a ±1.41	313.35^a ±1.87	317.45^a ±1.86	319.64^a ±1.67
G3	306.23^a ±1.57	287.47^b ±1.65	239.12^b ±1.85	216.61^b ±1.35
G4	305.12^a ±1.37	273.55^c ±1.68	204.13^c ±1.55	164.42^c ±2.03
G5	305.93^a ±2.03	264.46^d ±2.03	189.15^d ±1.46	147.33^d ±2.13

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

3.5. Effect of feeding bread sticks enriched with psyllium seeds powder on lipids serum parameters.

Data in Table 7, also show that treatments with psyllium seeds powder led to a gradual increase in serum HDL. Raising both treatments of the experiment caused an increase in serum HDL. While total cholesterol, LDL, VLDL and triglycerides decreased.

Table 7. Effect of feeding bread sticks with psyllium seeds powder on lipids serum parameters.

Rat groups	TC mg/dl(M±SD)	HDL-C mg/dl (M±SD)	LDL-C mg/dl (M±SD)	VLDL mg/dl (M±SD)	Triglycerides mg/dl (M±SD)
G1	110.53 ^e ±1.62	67.93 ^a ±1.45	20.77 ^e ±1.34	21.83 ^e ±2.01	109.15 ^e ±1.66
G2	153.51 ^a ±1.62	33.39 ^e ±1.37	84.3 ^a ±1.58	35.82 ^a ±2.14	179.1 ^a ±2.42
G3	133.85 ^b ±1.43	42.20 ^d ±2.56	63.34 ^b ±1.97	28.31 ^b ±2.54	141.55 ^b ±2.83
G4	123.54 ^c ±2.26	46.13 ^c ±2.51	51.59 ^d ±2.73	25.82 ^c ±2.41	129.10 ^c ±2.35
G5	121.86 ^d ±1.45	49.69 ^b ±1.98	48.16 ^c ±2.12	24.01 ^d ±2.61	120.05 ^d ±2.63

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

TC = Total cholesterol, HDL-C = High density lipoprotein cholesterol, LDL-C = Low density lipoprotein cholesterol

3.6. Effect of feeding bread sticks supplemented with psyllium seeds powder on liver functions.

Alanine amino transferase (ALT) and aspartate amino transferase (AST) enzymes activities are known as cytosolic marker enzymes reflecting hepatocellular necrosis as they are released

into the blood after the damage of the cell membrane. Therefore, both enzymes are used as indicators for hepatic damage (Andallu and Vardacharyulu, 2001). Data in Table 8 showed that the liver functions of normal control and diabetic groups through the experimental period (6 weeks). Liver functions of diabetic groups (G2) were markedly higher than that of normal control rats (G1). The results also indicated that, breadsticks contained psyllium seeds powder caused a significant decrease in liver functions of the diabetic rats (G3 to G5) comparing with diabetic control rats (G1) and diabetic rats fed on breadsticks contained wheat flour only (G2).

Table 8. Effect of feeding bread sticks supplemented with psyllium seeds powder on liver functions.

Rats groups	(AST) IU/L (10 - 37 IU/L) *	(ALT) IU/L (10 - 40 IU/L) *	Albumin (2 - 6 mg/dl) *
G1	35.62 ^b ±1.78	32.95 ^b ±1.66	3.55 ^d ±0.31
G2	46.42 ^a ±1.94	41.93 ^a ±1.78	6.38 ^a ±0.25
G3	35.65 ^b ±1.61	38.17 ^{ab} ±1.84	4.69 ^b ±0.36
G4	34.65 ^b ±2.06	36.63 ^{ab} ±1.52	4.10 ^b ±0.43
G5	32.32 ^b ±1.98	34.93 ^b ±1.78	3.95 ^c ±0.34

* Normal range (Ministry of Health, 2006) .

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

3.7. Effect of feeding bread sticks supplemented with psyllium seeds powder.

The results of kidney functions of negative and positive control at the end of experimental period are shown in Table 9 and the kidney functions of diabetic groups (G2) were markedly higher than that of negative control rats (G1). The results also indicated that, bread sticks contained psyllium seeds powder caused a significant decrease in kidney functions of the experimental groups (G3 to G5) comparing with negative control rats (G1) and diabetic rats (G2).

Table 9. Effect of feeding bread sticks at different psyllium seeds powder on kidney functions of diabetic rats.

Rat groups	Urea (10 - 50 mg/dL)*	Uric acid (2 - 6 mg/dL)*	Creatinine (0.5 - 1.5 mg/dL)*
G1	33.57 ^b ±2.25	2.82 ^{ab} ±0.23	0.65 ^c ±0.09
G2	43.91 ^a ±2.43	4.16 ^a ±0.38	1.38 ^a ±0.12
G3	35.25 ^b ±2.63	3.49 ^b ±0.27	0.99 ^b ±0.13
G4	34.25 ^b ±2.43	3.16 ^{ab} ±0.37	0.92 ^b ±0.12

G5	31.60 ^b ±1.97	3.00 ^{ab} ± 0.16	0.75 ^{ab} ±0.13
----	--------------------------	---------------------------	--------------------------

* Normal range (Ministry of Health, 2006). .

Each value is an average of five determinations.

Values followed by the same letter in column are not significantly different at $p \leq 0.05$.

3.8. Organoleptic properties of bread sticks supplemented with psyllium seeds powder.

Data in Table 10 shows the organoleptic properties of bread sticks supplemented with 3, 6 and 9% psyllium seeds powder. The obtained data indicated that, no differences were found between bread sticks produced by using 100% wheat flour and even those 3% psyllium seeds powder except for color, but differences were found in case of others levels of supplementation (6 and 9%). However, the results revealed that, the produced bread sticks by using 3% psyllium seeds powder replacement (instead of 100% wheat flour) had properties better than those of 6% and 9%. So, it can be concluded that, the best level of supplemented with psyllium seeds powder was 3% followed by 6% for making bread sticks with a very good acceptability. The results are harmony with the results of **EL Hadidy (2020)** which showed that adding psyllium seeds powder to pan bread was suitable for panelists.

Table 10. Organoleptic properties of bread sticks produced from wheat flour supplemented with different levels of psyllium seeds powder.

Blends	Organoleptic properties				
	Color (10)	Taste (10)	Flavor (10)	Texture (10)	Overall Acceptability (10)
Blend (1) 100 % wf	9.43 ^a ±0.35	8.97 ^a ±0.37	9.23 ^a ±0.28	8.67 ^a ±0.52	9.16 ^a ±0.46
Blend (2) + 3 % psp	7.56 ^b ±0.54	8.36 ^b ±0.26	8.98 ^a ±0.43	8.17 ^a ±0.44	8.72 ^a ±0.36
Blend (3) + 6 % psp	7.32 ^b ±0.26	7.78 ^b ±0.53	7.75 ^b ±0.32	7.45 ^b ±0.62	7.42 ^b ±0.28
Blend (4) + 9 % psp	6.63 ^c ±0.45	6.83 ^c ±0.46	7.52 ^b ±0.19	7.23 ^b ±0.26	7.37 ^b ±0.34

- Values followed by the same letter in columns are not significantly different at LSD at ($p \leq 0.05$).

- Each value was an average of twenty determinations ± standard deviation.

4. Discussion

The results of the present study show that injection with streptozotocin STZ produced hyperglycemia and hypoinsulinemia in rats. On the other hand, psyllium seeds reversed STZ induced change in glucose and insulin levels. STZ enter β cells via a glucose transporter (GLUT2) using a variety of intracellular toxic mechanisms such as production of oxygen free radicals that causes degeneration of pancreatic β -cells leading to hypoinsulinemia and subsequent hyperglycemia (**Guang-Kai *et al.*, 2017**).

Also, in agreement with our results (**Karhunen *et al.*, 2010**) found that psyllium fiber enriched meals improve glucose and insulin level significantly than non-fiber enriched meals. The creation of a thick gel in an aqueous solution is thought to be the cause of psyllium's glucose-lowering effects. This gel might prevent glucose from reaching the small intestine's absorptive epithelium, which would dampen postprandial glucose peaks. Additionally, soluble fiber may postpone stomach emptying, which would reduce the absorption of carbohydrates. Another mechanism that may contribute to the postprandial effect of psyllium is the sequestration of carbohydrates ingested with the meal, thus retarding their access to digestive enzymes (**Pal *et al.*, 2014**). Also, the hypoglycemic activity of psyllium seeds may be due to the inhibition of liver gluconeogenesis. Furthermore, the increased level of serum insulin by increase insulin secretion from the remnant of β -cells and enhancement of peripheral metabolism of glucose. Hypoglycemic activity of psyllium seeds may be due to modulating effects on insulin sensitization and/or insulin secretion and a regulating action on digestion and intestinal absorption.

Also, another mechanism by which psyllium promotes weight loss may be due to increase the energy expenditure in adipose tissue by stimulating lipolysis and thermogenesis by stimulation of adrenaline secretion from adrenal gland. This increase in serum adrenaline leads to activation of hormone-sensitive lipase (HSL), the key enzyme in the regulation of lipid stores. Moreover, HDL-C protects against or reverse atherosclerosis by their ability to serve as acceptor particles for macrophage cholesterol efflux, prevention of endothelial dysfunction, and maintenance of endothelial integrity (**Linsel-Nitschke and Tall, 2005**).

Conclusion

The findings suggested that psyllium seeds may be used as a new source of bioactive and functional food as well as an apotention agent for the treatment of diabetes rats. Comparing diabetic rats to healthy rats, daily administration of psyllium seed powder improved weight loss and decreased serum glucose, TC, TG, LDL and VLDL, urea, and uric acid while increasing HDL and improving liver and kidney function. Finally, it could make some bakery foods like bread sticks using PSP, and wheat flour (72%extr) with high quality that are suitable for diabetic patients.

References

- AOAC, (2005).** Official Methods of Analysis of the Association of Official Analytical Chemists. 18th edition, Washington DC.
- ASP, N.G., G.G. Johansson, H. Haller and M. Siljestrom, (1983).** Rapid enzymatic assay of insoluble and dietary fiber. *Journal of Agricultural and Food Chemistry*, 31: 476-482.
- Ballester-Sánchez, J.; Gil, J.V.; Haros, C.M.; Fernández-Espinar, M.T.(2019).** Effect of incorporating white, red or black quinoa flours on the total polyphenol content, antioxidant activity and colour bread. *Plant Food Hum. Nutr.* 74: 185–191.
- Bekyarova, G.Y., Ivanova, D.G. and Madjova, V.H.(2007).** Molecular mechanisms associating oxidative stress with endothelial dysfunction in the development of various vascular complications in diabetes mellitus. *Folia Med.*, 49, 13-19.
- Constantino, M.I., Molyneaux, L., Limacher-Gisler, F., Al-Saeed, A., Luo, C., Wu, T., Twigg, S.M., Yue, D.K. and Wong, J,(2013).** Longterm complications and mortality in young-onset diabetes: type 2 diabetes is more hazardous and lethal than type 1 diabetes. *Diabetes Care*, 36, 3863-3869.
- Dawson, R.M.C., Elliott, D.C., Elliott, W. H. and Jones, K.M. (1986)** Data for Biochemical Research, 3rd edition, Oxford: Calderon Press.
- El Hadidy, G. S. and Rizk, E. A. (2020).** Influence of Coriander Seeds on Baking Balady Bread. *J. Food and Dairy Sci., Mansoura Univ.*, 9 (2), 69 – 72.
- El-Dreny, E. G., and El-Hadidy, G. S. (2018).** Utilization of young green barley as a potential source of some nutrition substances. *Zagazig J. Agric. Res.*, 45(4),1333-1344. DOI: [10.21608/ZJAR.2018.48580](https://doi.org/10.21608/ZJAR.2018.48580).
- El-Dreny, E. G., and El-Hadidy, G. S. (2020).** Preparation of Functional Foods Free of Gluten for Celiac Disease Patients. *J. Sus. Agric. Sci.* 46, (1), 13- 24.
- Elhadidy, G. s. (2014).** Chemical, technological and biological studies on mulberry leaves and purslane in Egypt, Ph.D. Thesis, Food Indus. Dept., Fact. Agric. Mansoura Univ., Egypt.
- Elhadidy, G. s. (2009).** Chemical and Biological Studies on Some Hypoglycemic Foods. Ms c. Thesis, Food Techno. Dept., Fact. Agric. Kafrelsheikh Univ., Egypt.
- El-Hadidy, G. S. (2020).** Preparation and Evaluation of Pan Bread Made with Wheat flour and Psyllium Seeds for Obese Patients, *European Journal of Nutrition & Food Safety* **12(8): 1-13**. DOI: [10.9734/EJNFS/2020/v12i830256](https://doi.org/10.9734/EJNFS/2020/v12i830256)

El-Hadidy, G. S., and El-Dreny, E. G. (2020). Effect of Addition of Doum Fruits Powder on Chemical, Rheological and Nutritional Properties of Toast Bread, *Asian Food Sci, J*, 16(2),22-31. DOI: 10.9734/AFSJ/2020/v16i230169.

El-Hadidy, G. S., Eman, A. Yousef., and Abd El-Sattar, A. S. (2020). Effect of Fortification Breadsticks with Milk Thistle Seeds Powder on Chemical and Nutritional Properties, *Asian Food Sci, J*,17(2), 1-9. DOI: 10.9734/AFSJ/2020/v17i230187.

El-Hadidy, G. S., Rizk., E.A., and El-Dreny, E. G. (2020).Improvement of Nutritional Value, Physical and Sensory Properties of Biscuits Using Quinoa, Naked Barley and Carrot. *Egypt. J. Food. Sci*, 48, (1),147-157.

El-Hadidy, G., Nassef, S., and El-Sattar, A. (2022). Preparation of some functional bakeries for celiac patients. *Current Chemistry Letters*, 11(4), 393-402.

Fischer, H.M., Nanxiong, Y., Ralph, R.G.J., Anderson, L. and Marletta, J.A,(2017). The gel forming polysaccharide of psyllium husk (*Plantagoovata*Forsk). *Carbohydrate Res.*, 339.

Grant, S.J., Chang, D.H.T., Liu, J.X., Wong, V., Kiat, H. and Bensoussan, A, (2013).Chinese herbal medicine for impaired glucose tolerance: a randomized placebo controlled trial. *BMC Complement Altern. Med.*, 104, 1-8.

Guang-Kai .X.U. , Xiao-Ying Q.I.N., Guo-Kai W.A.N.G, Guo-Yong. X.I.E, Xu-Sen .L.I., Chen-Yu S.U.N., Bao-Lin L.I.U. and Min-Jian Q.I.N (2017). Antihyperglycemic, antihyperlipidemic and antioxidant effects of standard ethanol extract of *Bombaxceiba* leaves in high-fat-diet- and streptozotocin-induced Type 2 diabetic rats. *Chin. J. Natu. Med.* 15(3): 168-177.

Haddadian K, Haddadian K, and Zahmatkash M. A (2014).A review of plantago plant. *Ind. J. Tra. Knowl. ;* 13 (4):681-685.

Karhunen L.J., Juvonen K.R., and Flander S.M (2010). A psylliumfiberenriched meal strongly attenuates postprandial gastrointestinal peptide release in healthy young adults. *J. Nutr.* 40:737–744.

Kumar, V., Abbas, A.K., Fausto, N., Robbins, S.L. and Cotran, R.S,(2005) Robbins, and Cotran pathologic basis of disease, 7th edn, Elsevier Saunders, Philadelphia, USA, 1194-1195.

Lane-peter, W. and Pearson, A. E. (1971).Dietary requirement in "the laboratory animal Principals and practice" p142, Academic press, London and New York.

Linsel-Nitschke P.I., and Tall A.R. (2005). HDL as a target in the treatment of atherosclerotic cardiovascular disease. *Nat. Rev. Drug Discov.* 4:193–205.

Modi, P,(2007). Diabetes beyond insulin: review of new drugs for treatment of diabetes mellitus. *Curr. Drug Discov. Technol.*, 4, 39-47.

Pal S., Radavelli-Bagatini S., Ho S., McKay J. and Jane M. (2014). Using psyllium to prevent and treat obesity comorbidities. In *nutrition in the prevention and treatment of abdominal obesity*. Elsevier Inc. 505-514.

Renzo, D. (1975). Bakery products yeast leavened. Noyes Data Corporation. London, England.

Singh I.M., Shishehbor M.H., and Ansell B.J. (2007). High-density lipoprotein as a therapeutic target: a systematic review *J.A.M.A*; 298: 786-798.

Tewari, D. Anjum, N. and Tripathi, Y.C,(2014). Phytochemistry and pharmacology of *Plantago ovate*: a natural source of laxative medicine. *World J. Pharmaceutical Res.*, 3(9), 361-372.

Van Rosendaal G.M., Shaffer E.A., Alun L.E, and Brant R (2004). Effect of administration on cholesterol-lowering by psyllium: a randomized cross-over study in normocholesterolemic or slightly hypercholesterolemic subjects. *J. Nutr*; 28: 3– 17.