

Original Research Article

Relationship between family planning and malnutrition in children aged 0-24 months admitted to the nutritional centers of the Hospital of the Liberty and the Mother and Child University Hospital Center of N'Djamena (Chad).

ABSTRACT

This study was conducted with the aim of analyzing the relationship between the age of first pregnancy, the birth interval, family size and severe acute malnutrition in children aged 0 to 24 months living in Ndjamen admitted to the nutritional centers of the Hospital of the Liberty and the Mother and Child University Hospital Center of N'Djamena. It is a descriptive cross-sectional study, based on a field survey. It allowed us to collect data from 103 mother-child during the period from March 14 to May 15, 2018. The sampling was random according to the malnourished children received in the two hospitals during the survey.

A Slater scale, a UNICEF measuring rod, a survey form, a 2006 WHO Z-Score table were used to collect anthropometric data from children and those relating to family planning. The data collected was processed and analyzed manually and then on a computer using Word, Excel, Sphinx and Stata software. A univariate and bivariate analysis of the data was done.

The results obtained revealed that severe acute malnutrition affects 74.76% of the children surveyed. Of this population, 58.25% have an interbirth interval ≤ 24 months, 61.03% come from households with at least 6 individuals and 73.79% were born to mothers aged ≤ 25 years. From this work, the statistical tests using the Chi-square showed a clear significance between the dependent variable, the nutritional status of the children and each of the independent variables, which underlie the hypotheses of the study ($P < 0.01$). This made it possible to verify and confirm the main hypothesis of the study according to which "severe acute malnutrition in children aged 0 to 24 months is closely linked to short intervals between births, age at first birth and height of the family".

Keywords: Family planning, malnutrition, birth interval, Liberty Hospital, mother and child hospital, Ndjamen.

1. INTRODUCTION

Family planning "allows people to achieve the desired number of children and to determine the spacing of births". Family planning (FP) helps couples avoid unwanted pregnancies, reduces the spread of sexually transmitted infections (STIs), and decreases morbidity and mortality related to unwanted pregnancies. For this, she relies mainly on contraception. (WHO, 2016). The policy of family planning was mentioned by the

economist Malthus for whom the increase in population is a danger for the survival of humanity when as resources do not follow the same rhythm (anonymous, 2009) .

Over the years, the term has evolved involving several aspects, the one that interests us is the nutritional health of mother and child. This aspect, which seems to depend on food, the socio-economic level of households, Water-Hygiene and Sanitation actions and medical coverage also depends, according to several studies, on the spacing of births, the mother's age at first birth or even household size. Nutrition and family planning (FP) programs and services are often seen as separate, yet integrating these interventions can be mutually beneficial for mothers and children.

Central and West African countries have some of the highest fertility rates in the world, averaging 5.7 children per woman. In Chad, where the population has almost tripled over the past thirty years, this rate is 7 children per woman. The use of modern methods of contraception remains very limited throughout the country (Ministry of Health, 2012). Depending on the region, it concerns between 10% and 20% of all women aged 15 to 49, and the prevalence rate is even lower among 15 to 19 year olds. Teenagers who are forced into marriage represent 20, 30 and up to 60% of their age group. Maternal mortality rates are very high, with an average of 510 deaths per 100,000 live births (EDST-MICS, 2015).

In Chad, the prevalence of chronic malnutrition is 30.5% nationally. It appears that about a third of children under 5 suffer from growth retardation, of which 12% suffer from the severe form. The prevalence of GAM is 14.3% including 3.8% SAM in children aged 6 to 23 months against a prevalence of GAM of 7.6% including 1.2% SAM in children aged 24 to 59 months. These results show that children aged 6 to 23 months are significantly more affected by malnutrition than children aged 24 to 59 months. (National Nutrition Survey, 2020)

Faced with these high rates of malnutrition, the following study question imposed itself on us: what links can be established between the components of family planning and the occurrence of severe acute malnutrition in children from 0 to 24 months? living in Ndjamena? It is to answer this question that this study, which has as its title: " Relationship between family planning and malnutrition in children aged 0-24 months admitted to the nutritional centers of the Hospital of the Liberty and the Mother and Child University Hospital Center of N'Djamena" was initiated.

The overall objective of the study is: "to analyze the possible relationships between family planning/age of first pregnancy, birth interval, family size and severe acute malnutrition in children under 24 months living in N 'Djamena in Chad'. This will specifically be:

- Assess the relationship between the age of first pregnancy and severe acute malnutrition in children under 24 months;
- Assess the links between interbirth intervals and severe acute malnutrition in children under 24 months;
- To assess the links between family size and severe acute malnutrition in children under 24 months.

2. MATERIAL AND METHODS

2.1 Type of study

This is a descriptive cross-sectional study, based on a field survey. It allowed us to collect data for the period from March 14 to May 15, 2018 from dyads of mothers and children aged 0 to 2 admitted to the Mother and Child University Hospital Center as well as to that of the Hospital of the Liberty.

2.2 Study period

This research work was carried out over a period of six (06) months from February to July 2018. The month of February served us for the submission, correction and validation of the research protocol used to carry out this study. as well as obtaining authorization to collect data from the administrative managers of the two hospitals concerned. The months of March to May were devoted to the collection of survey data while those of June and July for the processing and analysis of data from the surveys, all this under the supervision of our thesis director.

2.3 Study population

The population used for this study consists mainly of:

- Children from 0 to 2 years old living in the province of N'Djamena and hospitalized in one of the nutrition centers of the Hospital of the Liberty and the Mother and Child University Hospital Center during the data collection period;
- Mothers of these children who met the study eligibility criteria.

2.3.1 Eligibility criteria

The following criteria were defined in the research protocol to allow us to define the sample of the population concerned by the study:

- For Children : these were children with an age less than or equal to two (0-2) years of age on the date of the survey; be malnourished and hospitalized in one of the nutrition centers of the Hospital of the Liberty and the Mother and Child University Hospital Center.
- For mothers: to be the mother of a child who meets the eligibility criteria and enrolled in the study, after informed consent of said mother;
- For the mother-child couple : come from a family that lives in the province of N'Djamena.

2.3.2 Exclusion criteria

Not considered by the study, all children and mothers who do not meet the eligibility criteria established above.

2.4 Sampling methods

This is a purposive, non-representative sample. It was obtained by reasoning based on the eligibility criteria over the data collection period, established by the research protocol. This sample is distributed as follows:

- 49 mother-child dyads (couples) surveyed at the Chad-China Liberty Hospital;
- 54 mother-child dyads (couples) surveyed at the Mother and Child Hospital.

2.5 Data collection and analysis tools

For the study, we used a survey data collection sheet which consists of in its first part with data relating to malnourished children and in its second part with data relating to mothers;

The data collected was then processed and analyzed manually and then on a computer using World, Excel, and Sphinx software.

The World software was used for entering textual data. Excel software was used for calculations, graphs and tables. The Sphinx software in its version v 5.1.0.4 was used for the design of the input mask, automatic input and data processing.

The main titles developed in the data collection sheet include:

- Identification of malnourished children;
- Anthropometric parameters of malnourished children;
- The family situation;
- Family planning and the gender relationship.

For the calculation of these anthropometric indicators in children, the following instruments were used:

- A Salter scale was used for weight gain;
- A UNICEF measuring rod for height determination;
- Survey props printed on A4 paper for note taking.

Before the operational phase of the collection itself, a test of the instruments was done in the two nutrition centers of the two hospitals. This test was intended to reassure us of the good condition of the equipment that will be used for the anthropometric measurements of children. The results obtained from this pre-test enabled us to definitively validate the survey sheet. The pre-test is the stage that followed the recruitment of survey agents.

3. RESULTS AND DISCUSSION

3.1 RESULTS

At the end of the surveys carried out with 103 mother and child dyads, having met the eligibility criteria and agreed to participate in the said study during the period indicated, several results were obtained.

3.1.1 Distribution of children by sex and age group

In accordance with table 1 of the results of the study which was carried out from March to May, the children who participated in the study are made up in more or less equal proportion of girls and boys. The age group most affected by severe acute malnutrition (SAM) at this time of year is that of 0 to 12 months, which constitutes 63% of the study sample.

Table 1. Distribution of children by gender and age group

		Children's age range				TOTAL
		0 to 6 months	6 to 12 months	12 to 18 months	18 to 24 months	
Sex	Male	16	17	11	7	51
	Female	16	16	8	12	52
TOTAL		32	33	19	19	103

3.1.2 Distribution of children according to their rank in the siblings

As for the rank of the children in the study among the siblings, the first born are the most affected by malnutrition. They make up 28%.The first three born to mothers surveyed together make up 65% of children affected by malnutrition (Fig. 1).



Fig.1. Distribution of children according to their rank in the siblings

3.1.3 Distribution of malnourished children according to their nutritional status

Fig. 2 reveals that, according to nutritional status, severely acute malnourished children ($-3Z$ -Score) are the ones who dominate in the study, with 74.8% of the study population.




Fig. 2. Proportion of malnourished children according to their nutritional status

3.1.4 Relationship between birth interval and nutritional status of children

The results in Table 2 show us that at the 1% threshold ($p = 0.000$), the birth interval is significantly associated with the nutritional status of the children in the study. In this table, children with a birth interval of 12 to 24 months represent more than half (60%) of children affected by acute malnutrition.

Table 2. Interbirth interval (IBI) and nutritional status of children

	Nutritional status of children							TOTAL
	No response	< -2 Z-Score	< -3 Z-Score	> -3 Z-Score and < -2 Z-Score	> -2 Z-Score	> -3 Z-Score	Special case	
No response	1	1	5	1	0	0	4	12
Under 12	0	0	13	0	0	0	1	14
From 12 to 24 months	0	2	48	6	0	0	4	60
25 to 36 months	1	1	2	2	0	0	2	8
From 37 to 48 months	0	0	3	0	0	0	0	3
48 months and over	0	0	6	0	0	0	0	6
TOTAL	2	4	77	9	0	0	11	103

Table 3 shows the results showing a significant correlation between household size and malnutrition, at the 1% threshold (P = 0.000).

Table 3. Household size and nutritional status of children

		Nutritional status of children							
		No response	< -2 Z-Score	< -3 Z-Score	> -3 Z-Score and < -2 Z-Score	> -2 Z-Score	> -3 Z-Score	Special case	TOTAL (%)
Number of individuals in households	No response	0.00	0.00	0.97	0.00	0.00	0.00	0.00	0.97
	≤ 5 individuals	0.00	0.97	29.13	2.91	0.00	0.00	2.91	35.92
	≥ 6 individuals	1.94	2.91	44.66	5.83	0.00	0.00	7.77	63.11
	TOTAL(%)	1.94	3.88	74.76	8.74	0.00	0.00	10.68	100.00

In accordance with the results presented in this table 3, assuming that families with no more than 5 people as being the least numerous and those with more than 6 people as the most numerous, the children of large families **more** exposed to malnutrition in a way general (63.11%) and severe in particular (44.66%). The study also revealed that the mother-child dyads surveyed come mostly from families whose size is between 5 to 10 people (54.90%).

The study revealed other interesting correlations, **which include:**

- The mothers who took part in this study largely come from monogamous households (67.96%). Even if the sample is not representative, we can affirm, for this study, that malnutrition is not associated with polygamy;
- Two monotheistic religions are practiced by the mothers and there is a clear predominance of mothers of Muslim faith (84.47%) who participated in the present study over those of Christian faith;
- The mothers surveyed are mostly young. The age group under 20 is the most represented with 37.88%. The accumulation up to 25 years constitutes 59.23%. The average age of these mothers is 23;
- Three-quarters (75%) of the first-born children in the study come from 37.88% of mothers whose age is less than or equal to 20 years;
- According to the mothers surveyed, the estimated average age for a girl to have her first **childbirth** is 16 years old. Among these mothers; 79.61% believe that a girl can conceive her first child before the age of 18;
- **Rare are the mothers surveyed who have a substantial income-generating activity. They are mostly housewives (72.82%). However, the few activities practiced and which generate income are trade, hairdressing, sewing, crafts, agriculture and livestock breeding.**
- Almost half of the mothers surveyed (49.51%) **have no** education. The section of literate mothers constitutes 7.77% of the study sample. The mothers of children who have received primary and secondary education form 26.21% of the sample, while mothers who know how to read the Koran constitute 14.56%.
- In the study population, 51.45% of mothers would like to have a number of children between 6-10 children against 28.15% of mothers who would like to have at most 5 children. The average of the study is 8 children per mother;
- Almost all of the mothers surveyed (94%) say they have not received training in family planning (FP) from health centres, hospitals and other structures in N'Djamena;
- With regard to the means used by mothers to delay or space pregnancies; 33.8% of mothers surveyed claim to practice abstinence while 48.54% claim to use modern methods of contraception including, in descending order: pills (44.66%), implants and injections (41.74%), condoms (33%). Mothers

who say they know nothing about these methods or do nothing at all to delay or space pregnancies make up 15.54%;

- The majority of women surveyed (87.37%) claim to recognize that family planning has advantages. It leads to better health for mothers (81.55%) and for children (70.87%); it offers more working time to mothers (33%), therefore a better economic situation (27.18%) and, finally, planning allows mothers to devote themselves more to the education of children.

- More than half (56.1%) of the mothers surveyed believe that a child should be breastfed until the age of 19 to 24 months and 51% of them affirm that from the age of 6 months, a mother can introduce complementary foods and water to infants;

- Most of the women surveyed affirm that there is a lack of communication in the home between the man and the woman, on family planning; some husbands impose their point of view.

3.2 Discussion

The main hypothesis (H0) of this study is "severe acute malnutrition in children aged 0 to 24 months is closely linked to short birth intervals, mothers' age at first pregnancy and family size".

The verification of the secondary hypotheses allowed us to confirm our basic hypothesis.

H₁: "The shorter the interbirth interval, the more children are exposed to severe acute malnutrition".

The processing of the data of our present study reveals that the children surveyed with a short birth interval suffer more from severe acute malnutrition compared to the rest. This confirms hypothesis H1. Out of 103 children who participated in our study, 85.71% suffered from severe acute malnutrition and were born or had a younger child after only 6 and 24 months. These short interbirth intervals, according to the results of our study, find their explanation in the low rate of use of contraceptive methods by the mothers surveyed.

Akoto and Hill (1988) established in their study that the birth interval is inversely associated with malnutrition: the shorter the birth interval (less than 2 years), the more precarious the nutritional status of children. While studies by Penders et al. (1997), showed that too short a birth spacing increases the risk of mortality and **retarded** growth in children as well as in their brothers and sisters, through prenatal growth retardation and early cessation of breastfeeding. maternal.

This situation can also be explained by the lack of communication from couples about family planning decisions.

H₂: "The larger the household size, the more the children are exposed to severe acute malnutrition".

Assuming that families of more than 6 people are the most numerous, 63.11% of these children in the study are exposed to malnutrition in general and severe malnutrition in particular (44.66%). The study also revealed that the mother-child dyads surveyed come mostly from families whose size is between 5 to 10 people (54.90%), which allows us to confirm hypothesis H2.

The average number of individuals per household from which our dyads come is 9. This average is therefore much higher than the national average for the fertility index of women, which is 7 children. This could find its explanation in the very fact of the conception of the child in Chadian society. The child is seen as a wealth of the family. In many households in our dyads, the study revealed that there is no decision on when to have children and many mothers are prevented from using contraceptives by their husbands. The results also revealed the lack of communication in the couples of our dyads with regard to family planning.

H₃: "The age of first childbearing, below 25 years, most often exposes children to severe acute malnutrition".

Of the 77 children in the study out of the 103 who suffer from Severe Acute Malnutrition (SAM), 28 are the first born of the mothers surveyed. Of these 28 first-born mothers, 21 come from mothers whose age is less than or equal to 20 years and among them, 20 suffer from SAM, or 71.42%. This number increases to 25 children if we consider the age of the mothers in the range of 14 to 25 years (89.28%). This allows us to say that the young age of the mother at her first maternity (first rank of the child in the siblings) exposes the child to SAM. This confirms research hypothesis H3.

AIDELF (2002) concluded in his work that children born to adolescent mothers are significantly more likely to experience stunted growth than those born to other women.. The young age of the mothers is a factor in severe acute malnutrition due to the inexperience of these mothers to properly take care of their child nutritionally. Another explanation is that of the early marriage of girls in Chad. In fact, 28% of women aged 15 to 49 were married before the age of 15 and 69% of women aged 20 to 49 were married before the

age of 18 (DHS, 2010). This rate varies according to the regions and oscillates between 60 and 84% in the other regions against 52% in N'djamena.

H4: Few women have knowledge of family planning and its links to their health and that of their children.

The mothers surveyed as part of this research have an overall knowledge of FP methods and its benefits for them and their children, but generally do not use its methods to capitalize on its benefits. With regard to the means used by mothers to delay or space pregnancies; 33.8% of mothers surveyed claim to practice abstinence while 48.54% claim to use modern methods of contraception including, in descending order: pills (44.66%), implants and injections (41.74%), condoms (33%). Mothers who say they know nothing about these methods or do nothing at all to delay or space pregnancies constitute 15.54%. This result allows us to invalidate our hypothesis H4.

Finally, the verification of the secondary hypotheses allows us to confirm the main hypothesis H0 of this study: "severe acute malnutrition in children aged 0 to 24 months is closely linked to short intervals between births, age at first pregnancy mothers and family size"

4. CONCLUSION

At the end of this study, conducted among 103 mother-child dyads hospitalized during the data collection phase both at the Hospital of the Liberty and the Mother and Child University Hospital Center, led us to the following conclusion: family planning is closely linked to the occurrence of severe acute malnutrition in children aged 0 to 24 months, through the short spacing between two consecutive births, the young age of mothers at their first pregnancy and the large size of families or households. This study also revealed that knowledge of contraceptive methods of family planning by mothers is satisfactory but the use of the latter remains a problem and this, by the opposition of husbands, religious prohibitions,

In light of the foregoing, it would be necessary to effectively involve family planning in nutrition programs or to create synergy between them in order to effectively guide the fight against child malnutrition in a country like Chad where households are generally numerous and where the fertility index is 7 children per woman. Women marry early. This integration of family planning would be beneficial for families, for the health of mothers and children if it can be made effective and sustainable.

REFERENCES

Alfred Sauvy, 1956. General Theory of Population, University Press of France, Paris, 2(01), Pages 1-19.

ACF-France, 2012. Conceptual models of child malnutrition, 76 pages.

ACF-Mali, 2015. Estudy on the link between family planning and acute malnutrition in Mali, 3 pages.

Akoto EM, 1993. Sociocultural determinants of child mortality in black Africa. Hypothesis and search for explanation, Louvain-la neuf, Academia, 269 pages.

Elisabeth Wollast, Marcel Vekemans, 1993. Practice and management of family planning in developing countries, Brussels, Ed de BOCK, 734 pages.

FAO, IFAD, WHO, WFP and UNICEF, 2017. The State of Food Security and Nutrition in the World- Building Resilience to Foster Peace and Food Security Rome, FAO.

KARIN RINGHEIM, JAMES GRIBBLE and MIA FOREMAN, 2011. Integrating Family Planning and Maternal and Child Health Care: Saving Lives, Money and Time, accessed online 11/29/2017, 6 pages.

Michael C. Latham, 1997. Nutrition in Developing Countries, News York, USA, FAO, 508 Pages.

Ministry of Public Health and National Solidarity of Chad, 2004. Policy and Service Standards for Family Planning in Chad, 35 pages.

Ministry of Public Health and National Solidarity of Chad, 2020. National Nutrition Survey, 68 pages.

Mwamba Odon Kadewe, 2009. The problem of family planning in the Kilobelobe district in the province of Katanga in the DRC, dissertation, University of Lubumbashi, Page 1-20.

PATH and UNFPA, 2006. Meeting a Need: Toward Strengthening Family Planning Programs, 56 pages.

RGPH 2-2009, March 2012. Final overall results, National Institute of Statistics, Economic and Demographic Studies, N'djamena-Chad, 44 pages.

Smith Rhonda et al., 2009. Family planning saves lives, online at www.hsrc.ac.za/Ed_Media/2009.htm, consulted on 05/11/2017.

The Lancet, 2008. Maternal and Child Nutrition, accessed online at www.lancet.com 10/28/2017, 15 pages.

Thomas Robert Malthus (1798), 1963. Essay on the principle of population, Paris-France, Gonthier, 236 pages.

UNDER PEER REVIEW