

# Kaposi Julius-berg syndrome on congenital ichthyosis, about a case

## Summary :

**Introduction: Kaposi-Julius berg syndrome** is a rare complication of infection with the virus of the herpes, often occurring on a pre-existing skin disease. It mainly affects children, The management of this syndrome is based on antiviral medication with antibiotic therapy. In the absence of diagnosis and early treatment, its prognosis is severe.

**Observation :** We present the first case of kaposi juliusberg syndrome on congenital ichthyosis described. This is a 5-year-old boy followed for congenital ichthyosis since birth, under keratolytic treatment and emollients. He was hospitalized for febrile rash made of lesions vesicular, pustular and crusty. He also presented with bilateral keratitis. The clinical examination showed a fever of 39 °C, an altered general condition with submandibular lymphadenopathy. The skin examination revealed a dozen vesicles and pustules in a bouquet on the face and scalp, meliceric crusts. antibiotic therapy with amoxicillin-clavulanic acid was instituted. The evolution was rapidly favorable. The clinical examination of the mother revealed a recurrence of oral herpes at the origin of the contagion, motivating the introduction of a preventive treatment for recurrences in her.

**Conclusion** This observation shows that it is essential to inform the family and friends of patients with ichthyosis of the risk of severe herpes infection and to avoid all contact with the herpes virus as far as possible.

**Key words:** congenital ichthyosis; HSV1; Kaposi-Julius berg, acyclovir, child

## Introduction :

Hereditary ichthyosis is a disease due to gene mutations involved in the development of the skin barrier, the latter is therefore very deficient. Hereditary ichthyosis begins, in the majority of cases, at birth or in childhood. As with all pre-existing skin diseases, the occurrence of herpes simplex virus type 1 (HSV1) infection responsible for Kaposi-Juliusberg herpetiform pustulosis (PHKJ) is classic; the same is true in atopic dermatitis (AD) as in other dermatological pathologies such as Darier's disease, pemphigus or epidermolysis bullosa (EB) of the Dowling-Meara type and seborrheic dermatitis (3).

Kaposi-Juliusberg syndrome begins with a fever of 40°C, accompanied by severe fatigue and chills. Numerous skin lesions then appear, predominantly on the face (sometimes haemorrhagic pustules, edema and redness of the skin).

These lesions can lead to peeling of the skin, necrosis of skin and corneal damage. They have the particularity of spreading very quickly over the body, and they are associated with inflammation of the lymph nodes. Bacterial superinfections are common.

Treatment with aciclovir or its precursor remains very effective, but in congenital ichthyosis there seems to be a need to use these antivirals at an appropriate dosage and rigorous follow-up in order to avoid relapses and complications.

### **Clinical case :**

We report the singular observation of KAPOSI JULIESBERG in a patient followed for ichthyosis after the free and informed consent of his parents. This is a 5-year-old male child, followed for congenital ichthyosis diagnosed at birth, treated with emollients and moisturizing creams, with a history of herpetic keratitis 3 years ago. He was hospitalized for a vesiculo-pustular febrile eruption, the clinical examination objectified an altered general condition made of anorexia and lethargy, a fever of 39°C, 40°C, the skin examination objectified pustules and vesicles in clusters, crusty, oozing, painful lesions of the face, scalp and pinnae of the ears with a limitation of the opening of the eyes, very evocative of kaposi Julius berg syndrome. There were infra-centimetric submandibular and retro-auricular adenopathies. Clinical symptoms revealed Kaposi Julius berg syndrome.

Fundus examination showed a dendritic-looking corneal ulcer that is pathognomonic for herpetic keratitis.



## **Figure 1**

**A): Vesiculo-pustular lesions grouped in a bouquet on the eczematous skin of the face, pavilion eyelids**

**B): vesiculo-pustular lesion of the face with ocular pus**

**C): vesicular-papular lesions on the pinna of the ears and scalp with ulceration and secondary impetiginization.**

Laboratory tests including complete blood count, CRP, kidney and liver function were done to watch for bacteremia or secondary visceral involvement and were found to be normal.

Injectable treatment with aciclovir at 10 mg/kg/8 h allowed an improvement in the general condition with gain of afebrile in 48 hours, disappearance of vesicular and bullous skin lesions in 22 days, it was necessary to continue the treatment long-term preventive based on emollients and keratolytics to preserve the skin barrier.

Emollients and keratolytics were applied once daily every day of the week + No topical corticosteroids were applied during the active period of the disease



**Figure 2: Healing of facial lesions after a 22-day follow-up**

### **Discussion :**

To our knowledge, no case of Kaposi Julius-berg syndrome on congenital ichthyosis was mentioned in the literature.

Superinfection with HSV1 or VZV is a classic complication of pre-existing skin diseases, cases are currently rarely reported except for a recent publication trying to correlate the intensity of MD (DARIER disease) to the frequency of occurrence of PVKJ (kaposi juliusberg varicella pustulosis) (1). Most of the publications relate to the occurrence of PVKJ during AD (atopic dermatitis), thus trying to correlate the occurrence of PVKJ with signs of severity of AD or with specific genetic factors, the dosage of aciclovir and valaciclovir are more important than usual. (1)

Atopic dermatitis (AD) is the most common inflammatory skin disease in children. The worldwide prevalence of AD is estimated to be 8 to 20%. It is characterized by recurrent chronic pruritic lesions with a different distribution according to age. Eczema herpeticum (EH) (Kaposi's varicelliform eruption) is an acute disseminated viral infection, usually Herpes simplexvirus (HSV) type 1, which develops in the presence of an existing skin lesion, often on the ground of the AD. Eczema herpeticum was first described in children and occurs with higher prevalence in childhood, (2)

Psoriasis is fertile ground for a herpetic transplant, however there are only a few cases in the literature, it is a very rare pathology in children (5)

There is a single described case of Dowling-Meara type EB complicated by Kaposi-Juliusberg syndrome (3).

Eczema herpetiformis is associated with a flu-like syndrome, fever, malaise, lymphadenopathy and may be complicated by keratoconjunctivitis, meningitis and encephalitis(2) which testifies to the severity of the prognosis.

These cases show the importance of preventing herpes superinfections linked either to HSV1, VZV or coxsackievirus in children with congenital ichthyosis, in the same way as in children with atopic dermatitis (2). Although there are few reported cases, the prevalence is probably higher. The information of the parents concerning the risk of contagion and the need for eviction must be systematic because there can exist serious clinical forms. This complication should be considered in any child with ichthyosis presenting with a febrile attack of the disease without an infectious point of infection found. In addition, watch out for any ocular symptoms that could be the cause of a complication, such as herpetic keratitis and blepharitis (4)

In our observation, the treatment of herpetic keratitis consisted of ocular washing with isotonic saline serum, intravenous aciclovir and the use of artificial tears without ocular corticosteroid therapy (4)

Morbidity and mortality from Kaposi Julius berg syndrome can be minimized by starting antiviral therapy as early as possible in diagnosis. The main treatment is acyclovir. (2) Systemic antivirals and hospitalization are recommended for severe illness and immunocompromised patients. (2) Secondary bacterial infection (mainly due to Staphylococcus aureus, Streptococcus pyogenes and Pseudomonas which may mask the underlying viral pathogenesis and sepsis are the most important complications, and therefore systemic antibiotic therapy should be added if necessary.

### **Conclusion :**

A careful history taken with awareness of risk factors and characteristic skin signs pointing to HE will be helpful for an accurate diagnosis. Laboratory evaluation is not diagnostic

But useful to support the clinic. Early diagnosis and timely antiviral treatment can prevent major complications and death.

**Conflict of interest:** The authors declare no conflicts of interest.

### **REFERENCES**

- 1) Ducournau<sup>1</sup>, A. Valois<sup>1</sup>, C. Brouillard<sup>1</sup> et al Kaposi-Juliusbergvarioliiformpustulosis in Darier'sdisease: should the dosage of valaciclovirbeadjusted? JP dermatology 2014
- 2) A.Ertuğrul, Z.ŞengülEmeksiz, E.UlusoySevercan, S. Özmen, I. Bostanc, Istanbul. Herpeticeczemaemergingduringatopicdermatitis in infants,TheTurkish Journal of Pediatrics 2020; 62: 514-519 <https://doi.org/10.24953/turkjped.2020.03.025>
- 3) J. Huguen<sup>1</sup>, P. La Selve<sup>2</sup>, S. Fraitag<sup>3</sup>, L.Misery<sup>1</sup>, C. Abasq-Thomas<sup>1</sup> Kaposi-Juliusberg syndrome complicatingDowling-Mearaepidermolysisbullosa, Dermatology, eslevier2014 <https://doi.org/10.1016/j.annder.2014.09.266>
- 4) p. thygeson, j. hogan,sjkinira, mdvorarv, pilaniap, jivani nb, kota rk. the unfavorableeffect of topicalsteroidtherapy on herpetickeratitis

- Kaposi's varicelliform rash. *Indian Dermatology Online* j 2015 ; 6: 364-366.
- 5) Saraswat1, RKRatho2, B. Kumar1 Two Unusual Cases of Kaposi's Varicelliform Eruption, DOI : 10.1080/00015550252948211
  - 6) Nath AK, Sori T, Thappa DM. A case series of kaposi's varicelliform eruption in dermatology in-patients in a tertiary care centre. *Indian J Dermatol.* 2011 Jan;56(1):110–5. 13
  - 7) KülçüÇakmak S, Alli N, Yilmaz E, Artüz F. A Case of Kaposi's Varicelliform Eruption in a Patient with Psoriasis Receiving Cyclosporine Therapy. *Ann Dermatol.* 2015 Jun;27(3):345–56. 14
  - 8) Garg G, Thami GP. Psoriasis herpeticum due to varicellazoster virus: a Kaposi's varicelliform eruption in erythrodermic psoriasis. *Indian J Dermatol.* 2012 May;57(3):213–4. 15
  - 9) Wakkee M, de Vries E, van den Haak P, Nijsten T. Increased risk of infectious disease requiring hospitalization among patients with psoriasis: a population-based cohort. *J Am Acad Dermatol.* 2011 Dec;65(6):1135–44. 16
  - 10) Demitsu T, Kakurai M, Azuma R, Hiratsuka Y, Yamada T, Yoneda K. Recalcitrant pemphigus foliaceus with Kaposi's varicelliform eruption: report of a fatal case. *Clin Exp Dermatol.* 2008 Aug;33(5):681–2. 17
  - 11) Sais G, Jucglà A, Curcó N, Peyrí J. Kaposi's varicelliform eruption with ocular involvement. *Arch Dermatol.* 1994 Sep;130(9):1209–10