

## Effect of the Golden Liquid from Honeybees and Refined Sugar on the Blood Glucose and Serum Iron Levels of Albino Rats

### ABSTRACT

This study investigated the effect of the golden liquid from honeybees (natural honey) and refined granulated sugar on the blood glucose and serum iron levels of Wistar albino rats. The experimental animals used in this study were grouped into five treatments based on the dose of natural honey and granulated sugar administered namely: T1 (1.02g of honey /kg BW), T2 (1.40g of honey /kg BW), T3 (1.02g of granulated sugar /kg BW), T4 (1.40g of granulated sugar /kg BW) while rats in T5 were not administered with honey and granulated sugar hence, served as the control. The results of the weekly glucose level of albino rat, orally administered natural honey and refined granulated sugar showed that the mean glucose level was highest in T3 (112.95mg/dl), followed by T5 (92.20mg/dl) while the least value was recorded in T2 (74.86mg/dl). The blood glucose concentration of the albino rats was measured using the glucose strips with glucometer while the serum iron analysis was conducted using Atomic Absorption Spectrophotometer. The result showed that the highest serum iron level was recorded in T2 ( $1.22^a \pm 0.115$ ) while the least glucose concentration was recorded in T2 ( $74.86^a \pm 24.457$ ) though the differences were not statistically significant ( $P > 0.05$ ). The use of natural honey is recommended since, albino rats orally administered honey at varying doses had lower blood glucose level and also the consumption of natural honey increases the serum iron level in the blood.

**Keywords:** Blood Glucose, Serum Iron Levels Albino Rats, Natural Honey, Refined Sugar

### INTRODUCTION

Honey is a sweet natural food made by bees (an insect) using water, pollen, and nectar from flowers (El Sohaimy *et al.*, 2015). Honey is produced by bees, In Nigeria; honey is commonly produced by honey bees from the Genus *Apis mellifera*. The colour, aroma and consistency of honey all depend upon which flowers the bees have been foraging. Honey is collected by most beekeepers and consumed by people (de Sousa *et al.*, 2016; Famuyide *et al.*, 2014). Nectar is reduced to honey containing predominantly carbohydrates with very little protein, vitamins, minerals, enzymes, amino acids, and as well several other compounds like phenol compounds thought to function as antioxidants (Wright *et al.*, 2018).

These chemical components are of great importance as they contribute to the quality, granulation, texture, nutritional and medicinal properties of honey (Kadirvelu and Gurtu, 2013). The major constituents of honey are nearly the same in all honey samples; however, the biochemical composition and physical properties of natural kinds of honey vary significantly according to the plant species on which the bees forage (Formickiet *al.*, 2013). Furthermore, the properties of natural kinds of honey also vary depending on the differences in climatic conditions and vegetation of the areas (Ononye and Akunne, 2019).

**Comment [MM(1):** The reference could be cited as [1] and in the reference section according to serial number.

**Comment [MM(2):** These could be cited as [2, 3] and in the reference section according to serial number

**Comment [MM(3):** The reference could be cited as [4] ..... and applicable for all references.

Natural honey is one of the most widely sought products due to its unique nutritional and medicinal properties, which are attributed to the influence of the different groups of substances it contains (Khan *et al.*, 2018). Honey as an antioxidant has been used by physicians throughout history and is utilized in both disease and health conditions, because of its anti-microbial and antioxidant properties (Adenekan *et al.*, 2012). Honey is full of enzymatic and non-enzymatic antioxidants such as catalase, alkaloids, ascorbate, and flavonoids that act against oxidative agents to prevent the body from combining with non-saturated fatty acids and lipoprotein oxidation (Okpokiri *et al.*, 2015). Honey is rich in fructose, glucose, and minerals such as magnesium, potassium, calcium, phosphate, sulphur, ferrous, and sodium chloride as well as vitamins such as B1, B2, B5, B6, and C (Adekanmbi *et al.*, 2019). However, natural honey is used as a replacement for many other sweeteners such as refined sugar/granulated sugar.

Refined granulated sugar commonly is widely used as a sweetener in meals in Nigeria such as breakfast cereals, beverages, and snacks, and many other food products (National Sugar Development Council (NSDC), 2020). Refined granulated sugar typically refers to a category of simple carbohydrates that includes monosaccharides like fructose and glucose, and disaccharides, like sucrose and lactose, which have different effects on the body (Green *et al.*, 2019). Notwithstanding the vast use of this product, there has been marked growing awareness of the metabolic and associated health effects of excessive sugar consumption in the past several years (Ohilebo, and Odudu, 2020). As of 2019, the World Health Organization recommends reducing the level of sugar consumption to less than 5% of daily caloric intake to lower the risk of unhealthy weight gain and obesity (Green *et al.*, 2019). The American Academy of Paediatrics recommended that parents should not feed fruit juice containing high amount of refined granulated sugar to infants younger than one year because of its high content (Fotereket *et al.*, 2016). There are continued awareness and a growing body of research investigating the adverse effect of refined granulated sugar as one of the causative agents of obesity and metabolic syndrome (a combination of risk factors like high blood pressure, high triglycerides, high fasting blood glucose, etc.) that increase the likelihood of cardiovascular disease, type 2 diabetes mellitus, and also non-alcoholic fatty liver disease (Stanhope, 2016).

The lifestyle of modern-day persons involves a high intake of refined granulated sugar either directly or indirectly from foodstuffs (Green *et al.*, 2019). The consumption of large amount of refined granulated sugar alters haematological and physiological changes in human body system (Pries *et al.*, 2017). Although, both honey and refined sugar are food components containing mainly fructose as sugars. Their dietary effects have been extensively investigated in adult animal models such as albino Wistar rats. Several scientific findings have revealed the use of honey as an alternative to the dangers and health risks associated with the consumption of refined granulated sugar (Samat *et al.*, 2017; Atangwhoet *et al.*, 2020). Many reports encourage consumption and the use of natural honey in food due to its nutritive and medicinal properties in human body (Oyeyemi *et al.*, 2017; Solayman *et al.*, 2016). The use of commercial sugar has been long documented (Rhee *et al.*, 2012). When food materials are ingested, the blood level changes from its normal state depending on the food substance consumed. Many micro nutrients in honey such as iron, calcium and potassium enhance the circulation of blood in human body. It is known that iron found in honey is a precursor for haemoglobin hence consumption of natural honey increases the red blood cell count and serum iron level in the body (Kadri *et al.*, 2017).

According to Erejuwa, (2014), administration of natural honey to albino rats was found to increase serum levels of insulin while it reduced serum concentrations of glucose and fructosamine in diabetic rats. Nevertheless, consumption of honey and refined granulated sugar also increases the blood glucose level at varying levels. In a study conducted by Atangwhoet *al.* (2020) noted that in comparison with refined sugar, honey may reduce weight gain and adiposity, presumably due to lower food intake and also promote lower serum triglycerides (Nemosecket *al.*, 2011). Hence, this study was carried out to determine the effect of natural honey and refined granulated sugar on blood glucose and serum iron level on albino rats in Awka, Anambra State.

Refined sugar is one of the well-known sweeteners which have become a favourite sweetener for centuries. It is in the form of crystallized sucrose (a combination of fructose and glucose) which is extracted from sugar beets or sugarcane. Refined granulated sugar has been a part of the daily diet for literally hundreds of years. Research is now suggesting that sugar intake can be detrimental to our health (Moseley, 2012). A growing body of scientific evidence now links long-term overconsumption of added sugars to diabetes, cavities, liver disease, and heart disease. Much of this evidence focuses on a cluster of metabolic issues, known collectively as metabolic syndrome (MetS), that raises people's risk of developing chronic diseases (Ekawati *et al.*, 2019). Atangwhoet *al.* (2020), revealed that replacing sugar with honey could lower blood sugar levels, improve the growth performance and prevent weight gain or aid in weight loss.

There has been an increasing demand of natural honey and refined granulated sugar by consumers in both developed and developing countries, as well as renewed interest in prospecting for sweeteners with high nutritional quality and pharmaceutical relevance. Interestingly, the outcomes of many of these research exercises on health effect of sweeteners such as diabetic, obesity, increased weight gain in the human body. Addition of excess sugar ultimately affects arteries and circulation and may increase the risk of coronary diseases and also negatively affects the body's organs in human body (Moeller *et al.*, 2017). Hence, natural honey as a sweetener has immense therapeutic, nutrition value and economic importance, which could extend beyond local boundaries of healthcare systems and national markets (Solayman *et al.*, 2016). It is also been recognized as having several beneficial health properties, including slower uptake into the bloodstream, pharmacological action of reducing blood glucose levels, and a high level of bioavailable antioxidants, all of which may mean that honey could be less harmful to health than refined granulated sugar in the human body (Atangwhoet *al.*, 2020). The aim of this study was to determine the effect of the golden liquid from honeybees and refined sugar on the blood glucose and serum iron levels of albino rats.

## **MATERIALS AND METHODS**

### **Study Area**

The study was carried out in the Animal House of the Department of Biochemistry, Nnamdi Azikiwe University, Awka, Nigeria. The experimental site lies between latitude 6° 15' 18.06" N and longitude 7° 06' 41.37" E. Awka is town is located on latitudes 6°9'19"N 7°07'12"E and stretches 8 km in an east-west direction along the Enugu-Onitsha expressway and about 5 km in a North-south orientation. The dimension of Awka is 1,207,800m<sup>2</sup> or 12,007 ha. Ecologically, Awka lies in the Guinea Savanna and has experienced 1798.52mm of rain annually between 1977 and 2019. It experiences two seasons: the dry and the wet seasons with a spell of harmattan between December and January (Abajue and Ewuim, 2020).

### **Source of Experimental Materials**

Twenty-five (25) healthy albino rats with mean weights ranging from  $41.21 \pm 1.247$ g to  $47.27 \pm 1.769$ g were used in the study which was purchased from Mr. Onyewuchi farms in Awka, Anambra State. The albino rats were carefully transferred to the Animal House of the Department of Biochemistry, Nnamdi Azikiwe University, Awka, where the experimental animal was housed for the whole period of the experiments. The refined sugar (Dangote granulated sugar) used for this study was used for the experiment was purchased from Eke-Awka market, Awka Anambra State while the freshly processed natural honey (Benbeela natural honey) was purchased from Uzoben Integrated Services in Awka Anambra State.

### **Management of experimental Animals**

The albino rats were intensively managed in an improvised metallic cage. Each cage consists of length 1m, width 0.5m and height 1m was used to house four albino rats. The top of the cages was covered with wire to allow proper ventilation and the floor was made of tin metal covered with saw dust. During the period of the experiment, the albino rats were subjected to similar sanitary condition so that the only source of variation is the dietary inclusions. Also, clean water was provided for the albino rats daily from 9am to 12pm for the period of 1 month.

### **Experimental Design**

For easy identification, the experimental rats were marked at the head, abdomen, centre, neck and tail then labelled H, Ab, C, N and T respectively. The albino rats were grouped into five treatment groups based on the dose of natural honey and granulated sugar namely: T1(1.02g of honey /kg BW), T2(1.40g of honey /kg BW), T3(1.02g of granulated sugar /kg BW), T4(1.40g of granulated sugar /kg BW) while rats in T5 was not administered with honey and granulated sugar hence was served as control. The experiment was set up in a completely randomized design. Each treatment comprised of five albino rats.

### **Glucose Content Determination**

The glucose content of the albino rats was measured using the glucose strips with glucometer. A puncture was made on the tale to collect the blood sample. A drop of blood was applied to a chemically active disposable test strip on which the chemical reaction occurred due to the action of glucose deoxyreductase that resulted in color change. Glucometer verified this colour change and showed glucose level. At the same time, a drop of capillary blood, (drawn from the heel) was sampled and applied to the test strip to measure level of glucose. Before sampling, the heel was warmed up by hand massage followed by disinfecting the spot, and then blood sample was taken (Flavigny, 2014).

## Serum Iron Analysis:

### Method for the analysis of Serum iron

Serum iron analysis was conducted using Varian AA240 Atomic Absorption Spectrophotometer according to the method of APHA 1995 (American Public Health Association) as reported by Lopes *et al.* (2019).

### Sample Dilution

1ml of sample (serum) was measured followed by a 5 dilution with distilled water. This was made up to 10ml for analysis. The absorbance was then measured using FS2400AA Agilent atomic absorption spectroscopy.

**Working principle:** Atomic absorption spectrometer's working principle is based on the sample being aspirated into the flame and atomized when the AAS's light beam is directed through the flame into the monochromator, and onto the detector that measures the amount of light absorbed by the atomized element in the flame. Since micro nutrients have their own characteristic absorption wavelength, a source lamp composed of that element is used, making the method relatively free from spectral or radiational interferences. The amount of energy of the characteristic wavelength absorbed in the flame is proportional to the concentration of the element in the sample.

### Preparation of reference solutions

A series of standard micronutrient solutions in the optimum concentration range are prepared, the reference solutions were prepared daily by diluting the single stock element solutions with water containing 1.5 mL concentrated nitric acid/litre. A calibration blank was prepared using all the reagents except for the micronutrient stock solutions. Calibration curve for each micronutrient was prepared by plotting the absorbance of standards versus their concentrations.

### Statistical Analysis

The data obtained from the study was recorded and captured Microsoft excels 2019. The data on the concentrations of blood glucose and serum iron levels in the five treatments was subjected to analysis of variance (ANOVA. Turkey honest significant difference (HSD) was used to separate means using IBM SPSS statistics version 23 at 5% significance ( $P < 0.05$ ).

## RESULTS

The result from Table 1 showed the weekly blood glucose level of albino rat orally administered natural honey and refined granulated sugar. The result showed that the mean weekly blood glucose level in week 1 was highest in T3 (112.95mg/dl), followed by T5 (92.200mg/dl), followed by T4 (93.700mg/dl) while the least value was recorded in T2 (74.866mg/dl). Statistical analysis showed that there were no significant differences in the albino rats orally administered natural honey and refined granulated sugar at varying levels ( $P > 0.05$ ).

**Table 1: Blood glucose concentration in albino rats orally administered natural honey and refined granulated sugar for a period of three weeks.**

Treatment	Weekly mean concentration of blood glucose level $\pm$ SD			
	Week 1	Week 2	Week 3	Mean blood glucose level $\pm$ SD
T1	78.2 <sup>a</sup> $\pm$ 8.871	84.7 <sup>a</sup> $\pm$ 19.709	104.7 <sup>b</sup> $\pm$ 23.905	89.20 <sup>a</sup> $\pm$ 20.808
T2	70.95 <sup>a</sup> $\pm$ 36.502	85.2 <sup>a</sup> $\pm$ 22.51	68.45 <sup>a</sup> $\pm$ 7.098	74.86 <sup>a</sup> $\pm$ 24.457
T3	97.7 <sup>a</sup> $\pm$ 22.726	144.95 <sup>b</sup> $\pm$ 34.956	96.2 <sup>a</sup> $\pm$ 21.993	112.95 <sup>a</sup> $\pm$ 34.407
T4	93.7 <sup>a</sup> $\pm$ 21.058	94.45 <sup>a</sup> $\pm$ 18.341	87.45 <sup>a</sup> $\pm$ 16.908	91.86 <sup>a</sup> $\pm$ 17.749
T5	92.45 <sup>a</sup> $\pm$ 22.19	94.95 <sup>a</sup> $\pm$ 21.045	89.2 <sup>a</sup> $\pm$ 3.962	92.20 <sup>a</sup> $\pm$ 16.663

*Columns with different superscript are significantly different (P<0.05)*

**Effect of Natural Honey and Refined granulated sugar on the serum iron levels of albino rats at different dosages.**

The result from Table 2 showed that the highest serum iron level was recorded in T2 (1.22 $\pm$ 0.115ppm) followed by T3, while the least serum iron level was recorded in T1 (0.88 $\pm$ 1.319ppm). It was observed that there were no significant differences (p>0.05) between the treatment groups orally administered natural honey and refined granulated sugar.

**Table 2 serum iron concentration of albino rats fed with natural honey and refined granulated sugar**

Treatment	Mean Serum iron level $\pm$ SD
T1	0.88 <sup>a</sup> $\pm$ 1.319
T2	1.22 <sup>a</sup> $\pm$ 0.115
T3	1.04 <sup>a</sup> $\pm$ 0.277
T4	0.99 <sup>a</sup> $\pm$ 0.299
T5	1.31 <sup>a</sup> $\pm$ 0.395

*Columns with different superscript are significantly different (P<0.05)*

**Discussion**

Findings of the study indicates that natural honey does not significantly increase the blood glucose level of albino rat as granulated sugar does. This finding is similar to that reported by Sohaimy *et al.* (2015) who noted that Dextrose (glucose) and laevulose (fructose) are the main sugars in honey. Similarly, Keket *et al.* (2017). These are the building blocks for the more complex honey sugars and account for about 85 percent of the solids present in honey. Furthermore, Adekanmbi *et al.*, (2019) reported that honey is rich in fructose, glucose, and minerals such as magnesium, potassium, calcium, phosphate, sulphur, ferrous, and sodium chloride as well as vitamins such as B1, B2, B5, B6, and C.

The result from the weekly glucose level in albino rats orally administered natural honey and refined sugar showed that highest value was found in T3. The findings of this study were in line with previous studies (Tappy and Le, 2010; Bezborodkina *et al.*, 2014; Sun *et al.*, 2010) on the effect of natural honey on regulation of blood glucose level and it benefits in the treatment of type 2 diabetes. Thus, bee honey is useful for diabetic patients as a tasty substance and as a nutritious substance. Honey may have a role in diabetic complications, so more effort is needed to be exerted in this field (Pasupuleti *et al.*, 2020).

The effect of natural honey and refined sugar on the serum iron levels of albino rats at varying doses showed that the highest serum iron level was recorded in T2 (1.40g of honey /kg BW) ( $1.22 \pm 0.115$ ) while the least serum iron level was recorded in T1 ( $0.88 \pm 1.319$  kg BW). the result of this study showed that honey inclusion at 1.40g of honey /kg BW has high serum. This has been reported by Erejuwa *et al.*, (2012) who reported that honey administration was found to increase serum iron levels of while it reduced serum concentrations of glucose and fructosamine in diabetic rats. It was further concluded that natural honey could affect the levels of many blood (haematological) elements in the body.

## Conclusion

This study showed that honey possess many advantages over commercial sugar, since the natural honey in this study honey possess high fructose content it means that honey can be used as a good source of carbohydrate. Also, natural honey in this study was found to contain substantial amount of copper and manganese which is good for human body compared to refined sugar. It was further observed that natural honey can be used to manage blood glucose level in the body and also that the consumption of natural honey increases the serum iron level in the blood. It was therefore recommended that natural honey should be used over refined sugar especially to people suffering from diabetes.

## REFERENCES

Abajue, M. C., and Ewuim, S. C. (2020). Evaluation of activities of dipteran maggots on a poisoned pig cadaver at Nnamdi Azikiwe University Awka, Nigeria. *Egyptian Journal of Forensic Sciences*, 10(1), 1-8.

- Abdulrhman, M., El Hefnawy, M., Ali, R., Hamid, I. A., Abou El-Goud, A., and Refai, D. (2013). Effects of honey, sucrose and glucose on blood glucose and C-peptide in patients with type 1 diabetes mellitus. *Complementary therapies in clinical practice*, 19(1), 15-19.
- Abeshu, M. A., and Geleta, B. (2016). Medicinal uses of honey. *Biology and Medicine*, 8(2), 1-9.
- Abeshu, M. A., and Geleta, B. (2016). Medicinal uses of honey. *Biology and Medicine*, 8(2), 1-11.
- Adadi, P., and Obeng, A. K. (2017). Assessment of bacterial quality of honey produced in Tamale metropolis (Ghana). *Journal of Food and Drug Analysis*, 25(2), 369-373.
- Adekanmbi, O. H., Walter, O. J., and Ikegbunam, N. C. (2019). Pollen analysis and heavy metals detection in honey samples from southern Nigeria. *World News of Natural Sciences. An International Scientific Journal*, 26, 1-9.
- Adenekan, M. O., Amusa, N. A., Lawal, A. O., and Okpeze, V. E. (2010). Physico-chemical and microbiological properties of honey samples obtained from Ibadan. *Journal of Microbiology and Antimicrobials*, 2(8), 100-104.
- Adenekan, M. O., Amusa, N. A., Okpeze, V. E., and Owosibo, A. O. (2012). Nutritional and microbiological components of honey samples obtained from Ogun State, Southwestern Nigeria. *European Journal of Sustainable Development*, 1(2), 271-271.
- Adepoju, O. T. (2009). Proximate composition and micronutrient potentials of three locally available wild fruits in Nigeria. *African Journal of Agricultural Research*, 4(9), 887-892.
- Affek, A. N. (2018). Indicators of ecosystem potential for pollination and honey production. *Ecological Indicators*, 94, 33-45.
- Agbagwa, O. E., Otokunefor, T. V. and Frank-Peterside, N. (2011). Quality assessment of Nigerian honey and manuka honey. *Journal of Microbiology and Biotechnology Resources*, 1(3), 20-31.
- Ahmed, A. A., Fadel-Almawla, W. A., Gasm-El-Bari, A. A., and Osman, Z. (2015). Clinical interactions of honey with antidiabetic drugs among some Sudanese diabetic patients in Khartoum. *Sudan Medical Monitor*, 10(3), 81-83.
- Alvarez-Suarez, J. M., Tulipani, S., Romandini, S., Bertoli, E., and Battino, M. (2010). Contribution of honey in nutrition and human health: a review. *Mediterranean Journal of Nutrition and Metabolism*, 3(1), 15-23.
- Alvarez-Suarez, M. J., Giampieri, F., and Battino, M. (2013). Honey as a source of dietary antioxidants: structures, bioavailability and evidence of protective effects against human chronic diseases. *Current Medicinal Chemistry*, 20(5), 621-638.
- Assia, A., and Ali, L. (2015). Enzymes activities, hydroxymethylfurfural content and pollen spectrum of some Algerian honey. *African Journal of Agricultural Research*, 10(7), 613-622.
- Bahrami, M., Ataie-Jafari, A., Hosseini, S., Foruzanfar, M. H., Rahmani, M., and Pajouhi, M. (2009). Effects of natural honey consumption in diabetic patients: an 8-week randomized clinical trial. *International Journal of Food Sciences and Nutrition*, 60(7), 618-626.

- Baydar, E., and Dabak, M. (2014). Serum iron as an indicator of acute inflammation in cattle. *Journal of dairy science*, 97(1), 222-228.
- Bezborodkina, N. N., Chestnova, A. Y., Okovity, S. V., and Kudryavtsev, B. N. (2014). Activity of glycogen synthase and glycogen phosphorylase in normal and cirrhotic rat liver during glycogen synthesis from glucose or fructose. *Experimental and Toxicologic Pathology*, 66(2-3), 147-154.
- Bleich, S. N., and Vercammen, K. A. (2018). The negative impact of sugar-sweetened beverages on ' 'children's health: an update of the literature. *Bio-medical Centre of Obesity*, 5(1), 1-27.
- BtHjIdrus, R., Sainik, N. Q. A. V., Nordin, A., Saim, A. B., and Sulaiman, N. (2020). Cardioprotective Effects of Honey and Its Constituent: An Evidence-Based Review of Laboratory Studies and Clinical Trials. *International Journal of Environmental Research and Public Health*, 17(10), 3613.
- Buawangpong, N., and Burgett, M. (2019). Capped Honey Moisture Content from Four Honey Bee Species; *Apis dorsata* F., *Apis florea* F., *Apis cerana* F, and *Apis mellifera* L. (Hymenoptera: Apidae) in Northern Thailand. *Journal of Apiculture*, 34(2), 157-160.
- Čaušević, B., Haurdić, B., Jašić, M., and Bašić, M. (2017). Enzymatic activities in honey. Drug i Kongres O Pčelarstvu I Pčelinjim Proizvodimasameddunarodnimučేశćem, 20. kolovoz 2017, Gradačac, Bosnia i Hercegovina. *Zbornik radova isazetaka*, 90-97.
- Chepulis, L. M. (2008). An Investigation of the Health Benefits of Honey as a Replacement for Sugar in the Diet (Thesis, Doctor of Philosophy (PhD)). The University of Waikato, Hamilton, New Zealand. Retrieved from <https://hdl.handle.net/10289/2612>.
- Chepulis, L. M. (2008). *An Investigation of the Health Benefits of Honey as a Replacement for Sugar in the Diet* (Doctoral dissertation, The University of Waikato).
- Chepulis, L., and Starkey, N. (2008). The long-term effects of feeding honey compared with sucrose and a sugar-free diet on weight gain, lipid profiles, and DEXA measurements in rats. *Journal of Food Science*, 73(1), H1-H7.
- Clemens, R. A., Jones, J. M., Kern, M., Lee, S. Y., Mayhew, E. J., Slavin, J. L., and Zivanovic, S. (2016). Functionality of sugars in foods and health. *Comprehensive Reviews in Food Science and Food Safety*, 15(3), 433-470.
- Codex Alimentarius Commission Standards. (2002): Draft revised standard for honey. 9-11 February 2000. Joint FAO/WHO Food Standards Programme
- Codex Alimentarius Committee on Sugars. (2001). Codex standard 12: Revised codex standard for honey. *Standard Methods*, 11, 1-7.
- de Sousa, J. M. B., de Souza, E. L., Marques, G., de Toledo Benassi, M., Gullón, B., Pintado, M. M., and Magnani, M. (2016). Sugar profile, physicochemical and sensory aspects of monofloral honeys produced by different stingless bee species in Brazilian semi-arid region. *LWT-Food Science and Technology*, 65, 645-651.

- Degani, G., Altomare, A. A., Colzani, M., Martino, C., Mazzolari, A., Fritz, G., ... and Aldini, G. (2017). A capture method based on the VC1 domain reveals new binding properties of the human receptor for advanced glycation end products (RAGE). *Redox Biology*, 11, 275-285.
- Edewor-Kuponiya, T. I. (2013). Determination of Fructose Content in *Anacardium occidentale*. *International Journal of Agricultural and Food Research*, 2(3), 2-6.
- Edwards, C. H., Rossi, M., Corpe, C. P., Butterworth, P. J., and Ellis, P. R. (2016). The role of sugars and sweeteners in food, diet and health: Alternatives for the future. *Trends in Food Science and Technology*, 56, 158-166.
- El Sohaimy, S. A., Masry, S. H. D., and Shehata, M. G. (2015). Physicochemical characteristics of honey from different origins. *Annals of Agricultural Sciences*, 60(2), 279-287.
- Erejuwa, O.O.; Sulaiman, S.A.; Wahab, M.S.; Salam, S.K.; Salleh, M.S. and Gurtu, S. (2011) Hepatoprotective effect of tualang honey supplementation in streptozotocin-induced diabetic rats. *International Journal of Applied Resources and National Production*, 4: 37-41.
- Erejuwa, O.O. (2014). Effect of honey in diabetes mellitus: matters arising. *Journal of Diabetes and Metabolic Disorders*, 13(1), 1-4.
- Erejuwa, O.O., Sulaiman, S.A., and Ab-Wahab, M.S. (2012). Honey: a novel antioxidant. *Molecules*, 17(4), 4400-4423.
- Erejuwa, O. O., Sulaiman, S. A., and Wahab, M. S. A. (2012). Fructose might contribute to the hypoglycemic effect of honey. *Molecules*, 17(2), 1900-1915.
- Eteraf-Oskouei, T., and Najafi, M. (2013). Traditional and modern uses of natural honey in human diseases: a review. *Iranian Journal of Basic Medical Sciences*, 16(6), 731.
- Eyer, M., Neumann, P., and Dietemann, V. (2016). A look into the cell: honey storage in honey bees, *Apis mellifera*. *Public Library of Science*, 11(8), e0161059.
- Famuyide, O. O., Adebayo, O., Owese, T., Azeez, F. A., Arabomen, O., Olugbire, O. O., and Ojo, D. (2014). Economic contributions of honey production as a means of livelihood strategy in Oyo State. *International Journal of Science and Technology*, 3(1), 7-11.
- Flavigny, R. (2014). Diabetic glucose meter for the determination of glucose in microbial cultures. *Journal of Microbiological Methods*, 100, 91-92.
- Folayan, J. A., and Bifarin, J. O. (2013). Profitability analysis of honey production in Edo North local government area of Edo state, Nigeria. *Journal of Agricultural Economics and Development*, 2(2), 60-64.
- Formicki, G., Greń, A., Stawarz, R., Zyśk, B., and Gał, A. (2013). Metal Content in Honey, Propolis, Wax, and Bee Pollen and Implications for Metal Pollution Monitoring. *Polish Journal of Environmental Studies*, 22(1). 2-9.
- Foterek, K., Buyken, A. E., Bolzenius, K., Hilbig, A., Nöthlings, U., and Alexy, U. (2016). Commercial complementary food consumption is prospectively associated with added sugar intake in childhood. *British Journal of Nutrition*, 115(11), 2067-2074.

- García, N.L. (2018). The current situation on the international honey market. *Bee World*, 95(3), 89-94.
- Getu, A., and Birhan, M. (2014). Chemical analysis of honey and major honey production challenges in and around Gondar, Ethiopia. *Academic Journal of Nutrition*, 3(1), 06-14.
- Goldman, R. D. (2014). Honey for treatment of cough in children. *Canadian Family Physician*, 60(12), 1107-1110.
- Green, M., Hadihardjono, D. N., Pries, A. M., Izwardy, D., Zehner, E., and Huffman, S. L. (2019). High proportions of children under 3 years of age consume commercially produced snack foods and sugar-sweetened beverages in Bandung City, Indonesia. *Maternal and Child Nutrition*, 15, e12764.
- Hilliard, M. E., Wu, Y. P., Rausch, J., Dolan, L. M., and Hood, K. K. (2013). Predictors of deteriorations in diabetes management and control in adolescents with type 1 diabetes. *Journal of Adolescent Health*, 52(1), 28-34.
- Hu, F. B., and Malik, V. S. (2010). Sugar-sweetened beverages and risk of obesity and type 2 diabetes: epidemiologic evidence. *Physiology and behaviour*, 100(1), 47-54.
- Huang, Z. (2010). Honey bee nutrition. *American Bee Journal*, 150(8), 773-6.
- Islam, M. N., Khalil, M. I., Islam, M. A., and Gan, S. H. (2014). Toxic compounds in honey. *Journal of Applied Toxicology*, 34(7), 733-742.
- Itan, Y., Powell, A., Beaumont, M. A., Burger, J., and Thomas, M. G. (2009). The origins of lactase persistence in Europe. *Public Library Computed Biology*, 5(8), 1-91.
- Kadirvelu, A., and Gurtu, S. (2013). Potential benefits of honey in type 2 diabetes mellitus: a review. *International Journal of Collaborative Research on Internal Medicine and Public Health*, 5(4), 199.
- Kadri, S. M., Zaluski, R., and de Oliveira Orsi, R. (2017). Nutritional and mineral contents of honey extracted by centrifugation and pressed processes. *Food Chemistry*, (218), 237-241.
- Kaškonienė, V., and Venskutonis, P. R. (2010). Floral markers in honey of various botanical and geographic origins: a review. *Comprehensive Reviews in Food Science and Food Safety*, 9(6), 620-634.
- Kebede, H., and Tadesse, G. (2014). Survey on honey production system, challenges and opportunities in selected areas of Hadya Zone, Ethiopia. *Journal of Agricultural Biotechnology and Sustainable Development*, 6(6), 60-66.
- Kek, S. P., Chin, N. L., Tan, S. W., Yusof, Y. A., & Chua, L. S. (2017). Classification of honey from its bee origin via chemical profiles and mineral content. *Food Analytical Methods*, 10(1), 19-30.
- Khan, S. U., Anjum, S. I., Rahman, K., Ansari, M. J., Khan, W. U., Kamal, S., ... and Khan, H. U. (2018). Honey: Single food stuff comprises many drugs. *Saudi Journal of Biological Sciences*, 25(2), 320-325.

- Kolayli, S., Boukraâ, L., Şahin, H., and Abdellah, F. (2012). Sugars in honey. Dietary sugars: Chemistry, analysis, function and effects. *Journal of Nutrition*, 3-15.
- Kumar, K. S., Bhowmik, D., Biswajit, C., and Chandira, M. R. (2010). Medicinal uses and health benefits of honey: an overview. *Journal of Chemistry and Pharmaceutical Resources*, 2(1), 385-395.
- Linkon, M. R., Prodhan, U. K., Elahi, T., Talukdar, J., Alim, M. A., and Hakim, M. A. (2015). Comparative analysis of the physico-chemical and antioxidant properties of honey available in Tangail, Bangladesh. *Universal Journal of Food and Nutrition Science*, 3(1), 19-22.
- Lopes, A., Alves, M., Pontes, J., Dias, N., Figueiredo, J., Santos, R., ...&Caseiro, A. (2019). Association between serum levels of C-reactive protein and lipid profile. *European Journal of Public Health*, 29(1), 35-39.
- Machado-De-Melo, A. A., Almeida-Muradian, L. B. D., Sancho, M. T., and Pascual-Maté, A. (2018). Composition and properties of *Apismellifera* honey: A review. *Journal of Apicultural Research*, 57(1), 5-37.
- Mahato, D. K., Keast, R., Liem, D. G., Russell, C. G., Cicerale, S., and Gamlath, S. (2021). Optimisation of natural sweeteners for sugar reduction in chocolate flavoured milk and their impact on sensory attributes. *International Dairy Journal*, 115, 104922.
- Majid, M., Younis, M. A., Naveed, A. K., Shah, M. U., Azeem, Z., and Tirmizi, S. H. (2013). Effects of natural honey on blood glucose and lipid profile in young healthy Pakistani males. *Journal of Ayub Medical College Abbottabad*, 25(3-4), 44-47.
- Malik, V. S., Popkin, B. M., Bray, G. A., Després, J. P., and Hu, F. B. (2010). Sugar-sweetened beverages, obesity, type 2 diabetes mellitus, and cardiovascular disease risk. *Circulation*, 121(11), 1356-1364.
- Manley, D. (2011). Sugars and syrups as biscuit ingredients. In ' *Manley's technology of biscuits, crackers and cookies*. Woodhead Publishing (pp. 143-159).
- Manyi-Loh, C. E., Clarke, A. M., and Ndip, N. (2011). An overview of honey: Therapeutic properties and contribution in nutrition and human health. *African Journal of Microbiology Research*, 5(8), 844-852.
- Manyi-Loh, C. E., Ndip, R. N., and Clarke, A. M. (2011). Volatile compounds in honey: a review on their involvement in aroma, botanical origin determination and potential biomedical activities. *International Journal of Molecular Sciences*, 12(12), 9514-9532.
- Meixner, M. D. (2010). A historical review of managed honey bee populations in Europe and the United States and the factors that may affect them. *Journal of Invertebrate Pathology*, 103, S80-S95.
- Mesele, T. L. (2021). Review on physico-chemical properties of honey in Eastern Africa. *Journal of Apicultural Research*, 60(1), 33-45.
- Minhas, S., and Dhaliwal, Y. S. (2018). Effect of processing on proximate composition of honey during storage. *International Journal of Food Science and Nutrition*, 3(2), 160-165.

- Mohammed, M. E. A. (2020). Factors Affecting the Physicochemical Properties and Chemical Composition of Bee's Honey. *Food Reviews International*, 1-12.
- Nayik, G. A., and Nanda, V. (2016). A chemometric approach to evaluate the phenolic compounds, antioxidant activity and mineral content of different unifloral honey types from Kashmir, India. *LWT*, 74, 504-513.
- Nazir, L., Samad, F., Haroon, W., Kidwai, S. S., Siddiqi, S., and Zehravi, M. (2014). Comparison of glycaemic response to honey and glucose in type 2 diabetes. *J. Pakistan Medical Association*, 64(1), 69-71.
- Ndife J., Abioye L. and Dandago M. (2014). Quality assessment of Nigerian honey sourced from different floral locations. *Nigerian Food Journal*, 32(2):48-55.
- Nigussie, K., Subramanian, P. A., and Mebrahtu, G. (2012). Physicochemical analysis of Tigray honey: An attempt to determine major quality markers of honey. *Bulletin of the Chemical Society of Ethiopia*, 26(1).1-7.
- O'Connell, S. (2012). *Sugar: the grass that changed the world*. Random House Publisher, pp6.
- Okpokiri, C. I., Nwachukwu, I. N., and Onwusiribe, C. N. (2015). Determinants and profitability of honey production in Ikwuano Local Government Area, Abia State, Nigeria. *Scientific Papers Series Management, Economic Engineering in Agriculture and Rural Development*, 15(3), 211-216.
- Omojokun, J. (2013). Regulation and enforcement of legislation on food safety in Nigeria. *Mycotoxin and Food Safety in Developing Countries*, 251-268.
- Ononye, B. U., and Akunne, C. E. (2019). Microclimate, Colonization, Colony Strength and Absconding of *Apis mellifera adansonii* Reared in Hives with Modified Entrances. *Journal of Apiculture*, 34(2), 117-123.
- Oyeyemi S. D. (2017). Quality Assessment of Honey Sourced from Natural and Artificial Apiaries in Ekiti State, Nigeria. *Turkish Journal of Agriculture - Food Science and Technology*, 5(10): 1125-1129.
- Pasupuleti, V. R., Arigela, C. S., Gan, S. H., Salam, S. K. N., Krishnan, K. T., Rahman, N. A., and Jeffree, M. S. (2020). A Review on Oxidative Stress, Diabetic Complications, and the Roles of Honey Polyphenols. *Oxidative medicine and cellular longevity*, 4(1), 1-8.
- Popov-Raljić, J., Arsić, N., Zlatković, B., Basarin, B., Mladenović, M., Laličić-Petronijević, J., ... and Popov, V. (2015). Evaluation of colour, mineral substances and sensory uniqueness of meadow and acacia honey from Serbia. *Romanian Biotechnological Letters*, 20(5), 10-84.
- Prica, N., Baloš, M. Ž., Jakšić, S., Mihaljev, Ž., Kartalović, B., Babić, J., and Savić, S. (2014). Moisture and acidity as indicators of the quality of honey originating from Vojvodina region. *Archives of Veterinary Medicine*, 7(2), 99-109.
- Pries, A. M., Huffman, S. L., Champeny, M., Adhikary, I., Benjamin, M., Coly, A. N., and Zehner, E. (2017). Consumption of commercially produced snack foods and sugar-sweetened beverages during the complementary feeding period in four African and Asian urban contexts. *Maternal and Child Nutrition*, 13, e12412.

- Ramli, N. Z., Chin, K. Y., Zarkasi, K. A., and Ahmad, F. (2018). A review on the protective effects of honey against metabolic syndrome. *Nutrients*, *10*(8), 1009.
- Rhee, J. J., Mattei, J., and Campos, H. (2012). Association between commercial and traditional sugar-sweetened beverages and measures of adiposity in Costa Rica. *Public Health Nutrition*, *15*(8), 1347-1354.
- Sakač, M. B., Jovanov, P. T., Marić, A. Z., Pezo, L. L., Kevrešan, Ž. S., Novaković, A. R., and Nedeljković, N. M. (2019). Physicochemical properties and mineral content of honey samples from Vojvodina (Republic of Serbia). *Food Chemistry*, *276*, 15-21.
- Saklayen, M. G. (2018). The global epidemic of the metabolic syndrome. *Current Hypertension Reports*, *20*(2), 1-8.
- Seguí, L., Calabuig-Jiménez, L., Betoret, N., and Fito, P. (2015). Physicochemical and antioxidant properties of non-refined sugarcane alternatives to white sugar. *International Journal of Food Science and Technology*, *50*(12), 2579-2588.
- Sheiham, A., and James, W. P. T. (2014). A new understanding of the relationship between sugars, dental caries and fluoride use: implications for limits on sugars consumption. *Public Health Nutrition*, *17*(10), 2176-2184.
- Solayman, M., Islam, M. A., Paul, S., Ali, Y., Khalil, M. I., Alam, N., and Gan, S. H. (2016). Physicochemical properties, minerals, trace elements, and heavy metals in honey of different origins: a comprehensive review. *Comprehensive Reviews in Food Science and Food Safety*, *15*(1), 219-233.
- Solayman, M., Islam, M. A., Paul, S., Ali, Y., Khalil, M. I., Alam, N., and Gan, S. H. (2016). Physicochemical properties, minerals, trace elements, and heavy metals in honey of different origins: a comprehensive review. *Comprehensive Reviews in Food Science and Food Safety*, *15*(1), 219-233.
- Stanhope, K. L. (2016). Sugar consumption, metabolic disease and obesity: The state of the controversy. *Critical reviews in Clinical Laboratory Sciences*, *53*(1), 52-67.
- Stanner, S. A., and Spiro, A. (2020). Public health rationale for reducing sugar: Strategies and challenges. *Nutrition Bulletin*, *45*(3), 253-270.
- Steinhauer, N., and Saegerman, C. (2021). Prioritizing changes in management practices associated with reduced winter honey bee colony losses for US beekeepers. *Science of The Total Environment*, *753*, 141629.
- Sun, S. Z., Flickinger, B. D., Williamson-Hughes, P. S., and Empie, M. W. (2010). Lack of association between dietary fructose and hyperuricemia risk in adults. *Nutrition and Metabolism*, *7*(1), 1-12.
- Tappy, L., and Lê, K. A. (2010). Metabolic effects of fructose and the worldwide increase in obesity. *Physiological reviews*. *3*(1), 1-4)
- Uluozlu, O. D., Tuzen, M., Mendil, D., and Soylak, M. (2007). Trace metal content in nine species of fish from the Black and Aegean Seas, Turkey. *Food Chemistry*, *104*(2), 835-840.

- Wang, Y., Kaftanoglu, O., Brent, C. S., Page, R. E., and Amdam, G. V. (2016). Starvation stress during larval development facilitates an adaptive response in adult worker honey bees (*Apis mellifera* L.). *Journal of Experimental Biology*, 219(7), 949-959.
- White, J. S. (2014). Sucrose, HFCS, and fructose: history, manufacture, composition, applications, and production. In *fructose, high fructose corn syrup, sucrose and health*. Humana Press, New York, NY (pp. 13-33).
- Wright, G. A., Nicolson, S. W., and Shafir, S. (2018). Nutritional physiology and ecology of honey bees. *Annual Review of Entomology*, 63, 327-344.
- Yetimwork, G. (2015). *Characterization of beekeeping systems and honey value chain, and effects of storage containers and durations on physico-chemical properties of honey in kilteAwlaelo district, eastern Tigray, Ethiopia* (Doctoral dissertation, Addis Ababa University). Pp1-68.

UNDER PEER REVIEW