

Basis for Enhancement of Nursing Concept and Skills: Assessment of Knowledge, Attitude and Practices of Nurses on Essential Newborn Care

ABSTRACT

Background: The development and implementation of the Department of Health's *Unang Yakap* or Essential Newborn Care Protocol in 2009 encouraged the health facilities, both government and private, to standardize newborn care practices as well as its proper sequencing based on current evidences that show reduction in neonatal morbidity and mortality. (DOH PPDCPM, 2016). Attitudes of healthcare workers also contribute to how one execute the Essential care.

Purpose: Assess the level of knowledge, attitude and practice of nurses on District hospitals on Essential Newborn Care to contribute to the nursing service in participating to updates and trainings of Essential Newborn Care.

Methods: The descriptive correlation study was utilized in the study to determine the assessment on knowledge, attitude and practices of nurses regarding Essential Newborn Care protocol. A total of 146 participants of the study.

Result: The result shows that 60.2% of Filipino nurses age range around 20-29 years old. This age group are considered to be at their peak of searching for clinical experience and career enhancement opportunities. The lowest age group was 51 years old and above, which is considered almost as close to retirement age.

Conclusion: The level of district nurses' Knowledge on Essential Newborn care and some of the vital aspects on it is low due to lack of proper training of the involved nurses.

Recommendation: some unnecessary and deficient procedures need to be re-taught to achieve excellent execution of Essential New-born Care protocol.

Keywords: Assessment of knowledge; attitude; practices of nurse; essential newborn care.

1. INTRODUCTION

Nursing shortage is still an existing concern in the Philippine healthcare industry. District Hospitals is not an exemption to this. For example, one district hospital only have 27 nurses, with impending 2 resigning nurses, during the time of data collection; the proposed plantilla is to have 34 nurses for their 30 bed capacity. This nursing staff shortage had made it difficult for the chief nurses to allot time for all the necessary trainings for nurses. Attitudes of healthcare workers also contribute to how one execute the Essential Newborn care[6-9]. Liao & Manalon [1] found out that not all steps in the Essential Newborn Care protocol were practiced because some staffs on the hospital believed that in the olden days without ENC, every newborn still survives (Noon naman walang

ENC, buhay naman lahat). This implied that training alone cannot change an old attitude or habit for some healthcare workers

1.1 Aim of the Study

The main purpose of this study is to assess the level of knowledge, attitude and practice of Bulacan nurses on District hospitals on Essential Newborn Care to contribute to the nursing service in participating to updates and trainings of Essential Care. The study intended to enumerate the identified gaps on knowledge and practice of ENC based on the DOH protocol AO 2009-0025, in order to give attention to these identified gaps for re-learning or re-training. This study shall be significantly beneficial to the concerned health institutions in reviewing current hospital policies on essential newborn care to

even the gap of the staff nurses handling such practices. This research shall also help medical personnel to understand each importance of the protocols for further improvement of the response they are taking in handling such cases. Lastly, the study intended to contribute to the nursing service division on the ease of decking their staff nurses for Essential Newborn Care training scheduling by considering the demographic profile of the nurses [17-20].

2. METHODS

2.1 Research Design

The descriptive correlation study was utilized in the study to determine the assessment on knowledge, attitude and practices of nurses regarding Essential Newborn Care protocol. Correlational research is a systematic investigation of the relationship present between two or more variables. The study made use of a quantitative research approach in analyzing and understanding the predictor and criterion variables.

2.2 Respondents of the Study

The main participants of the study were the nurses employed in District hospitals whose work involves the performance of ENC. There are five district hospitals in the province that was qualified to be part of the study. District Hospitals are Level I government hospitals that includes emergency room, operating room, recovery room, maternity facilities, isolation facilities, pharmacy, and clinical laboratory. A total of 146 respondents participated in the study. The researcher selected sample by purposive sampling based on the respondents' availability and willingness to participate. Purposive sampling is a non-probability sampling technique that make use of samples that are chosen by the judgement of the researcher. Due to the current health situation and strict health protocols imposed, the researcher had only gathered data from available district hospitals that had allowed the researcher during the time of data collection

2.3 Tool of the Study

To assess the nurses' Knowledge of Essential Newborn Care, the researcher adapted DOH guideline and standardized study tool of Bayisa Bereka Negussie and company [2] with reliability test done and Cronbach's alpha coefficient of 0.84. This is a 20 item multiple choice questions

To assess the nurses' attitude towards the Essential Newborn Care, a 9 item structured questions based on the theory of Planned Behavior will be used, a tool adopted from literature by Horiuchi, S., Rattana et al. [3]. This is a 5 point Likert scale with numbers corresponding as 5-Strongly Agree, 4-agree, 3-Undecided, 2-disagree and 1-strongly disagree .

To properly assess the Practice of ENC among the hospitals, the researcher adapted the tool of WHO and contextualized by the Department of Health to the study. This is a 48-item standardized questionnaire with a 5 point Likert scale with numbers corresponding as 5-Always, 4-Very often, 3-Often, 2-Seldom and 1-Never.

2.4 Data Gathering Procedure

The questionnaire method was the mode of data gathering. Each of the respondents was given a structured and standardized set of questions. In gathering the data, the researcher carried out the following procedure. (1) After the approval of Research Ethics Committee, the researcher secured a letter of permission from the hospital to conduct data gathering. (2)After the approval of the letter, the researcher waited for the data to be generated. (3) The study was conducted in district hospitals with approved permission letter on May 12 to 27, 2021. An approximate of 2 weeks' time was given by the researcher in order to achieve the desired number of respondents who will finish the questionnaires. (4)The answered questionnaires were collected by the chief nurses/head nurses and then handed to the researcher. Out of the targeted 146 respondents, a total of 125 completed questionnaires were returned to the researcher. After receiving the data, the researcher collated the data and handed it out to the research statistician. (5) The data were organized into tables and be removed of any confidential information, in accordance to the Data Privacy Act of 2012. (6) After the research, the researcher shall dispose of the data and inform the entities about the research outcome.

2.5 Statistical Analysis

The data collected was tabulated and processed with the help of Statistician and SPSS. To analyze and interpret the data gathered, the following statistical measures were used: (1) The profile of the nurses were described by means of frequency and percentage(2)The level of nurses' performance to render ENC in terms of Knowledge of Essential Newborn care were

described by means of frequency and percentage. And then, the computed mean score was also calculated.(3) Attitude towards Essential Newborn care were quantified using the following scale and were described by total average Mean.

3. RESULTS AND DISCUSSION

This chapter presents analyses and interprets the data collected in the study. For clarity of presentation and consistency in the discussion, the data are presented following the order and sequence of the questions raised in Chapter 1, to wit: (1) the demographic profile of the nurses in terms of age, area of specialization and number of ENC trainings attended (2) level of nurses' performance to render ENC in terms of Knowledge of Essential Newborn Care (3) level of nurses' performance to render ENC in terms of Attitude towards the Essential Newborn Care (4) level of nurses' performance to render ENC in terms of Practice of Essential Newborn Care (5) significant difference on the level of nurses' performance according to their demographic profile in terms of knowledge, attitude and Practices of Essential newborn Care.

3.1 Demographic Profile of Nurses

Table 1 present the frequency and percentage distribution regarding age profile ranging to 12 years up to 51 years and above. It can be deduced from the table that the highest frequency were those aged from 26 to 30 years with 52 nurse respondents or 41.6% of the population. It was followed by aged range of 31 to 35 years old with 28 nurse respondents or 22.4% of the population Third, was the aged range of 41 to 45 years old with 13 nurses or 10.4% of the total population. Moreover, no nurse respondents have been registered for aged less than 20 years old while the lowest

among the age range was those with 51 years and above having 4 nurses or 3.2 % of the population and the 2nd lowest were those 36 to 40 years old with 6 nurse respondents or 4.8 % of the total population.

In terms of the baby first bath, it can be deduced from the table almost half or 59 (47.2%) nurse respondents got the answer correctly that is should be at least 24 hours after birth while 40 (32%) chosen that bathing be done six hours after birth, and 26 (20.8%) nurse respondents tell that should be bathe immediately. In terms of knowledge on thermal protection of newborn, it is noted that 16 (12.8%) of the nurse respondents able to answered that it should immediately dry baby after birth while majority or 109 (87.2%) nurse respondents answered that it is skin to skin contact.

On item where should you placed the baby after birth, It is noted that 111 (88.8%) nurse respondents able to get a correct answer of putting on mother's abdomen or chest while 10 (8.0%) have answered it should be beside the mother, and 4 (3.2%) said that the baby should be placed on table/bed. The Department of Health, [4] had outlined in the implementation of Essential Newborn Care, that newborn should be placed prone on mother's abdomen or chest, skin-to-skin to facilitate bonding between mother and child and reduce likelihood of infection[26] and Current evidence on WHO, [5] indicated that skin-to-skin contact between mother and infant shortly after birth helps to initiate early breastfeeding and increases the likelihood of exclusive breastfeeding for one to four months of life as well as the overall duration of breastfeeding. Infants placed in early skin-to-skin contact with their mother also appear to interact more with their mothers and cry less.

Table 1. Frequency and percentage distribution regarding age profile

Age Range	Frequency	Percent
12 to 25 years old	12	9.6
26 to 30 years old	52	41.6
31 to 35 years old	28	22.4
36 to 40 years old	6	4.8
41 to 45 years old	13	10.4
46 to 50 years old	10	8
51 and above	4	3.2
Total	125	100

Table 2.1. Frequency and percentage distribution regarding the level of the nurse's performance to in rendering essential newborn care in terms of knowledge on immediate and thorough drying of newborn

Variables	Responses	Frequency	Percent
When should the baby be first bathe?	Immediately	26	20.8
	Within 6h of birth	40	32.0
	At least 24hours after birth	59	47.2
How do you practice thermal protection of newborn?	Immediately dry baby after birth	16	12.8
	Allow skin to skin contact	109	87.2
	Immediately bath the baby		

Table 2.2. Frequency and percentage distribution regarding the level of the nurse's performance to in rendering essential newborn care in terms of knowledge on early skin-to-skin contact between mother and newborn

Variables	Responses	Frequency	Percent
Where should you placed the baby after birth?	Beside the mother	10	8.0
	With someone else	-	-
	On mother's abdomen or chest	111	88.8
	On newborn table/bed	4	3.2

Table 3. Mean score & descriptive interpretation regarding the level of the nurse's performance in rendering essential newborn care in terms of attitude towards essential newborn care

Variables	Mean score	Descriptive interpretation
For me, to provide the EENC for every newborn I assist birth in this hospital on a regular basis is beneficial.	4.87	Strongly Agree
My providing EENC for every newborn at this hospital on regular basis will result in preventing newborn morbidities and mortalities.	4.40	Agree
For me, preventing newborn morbidities and mortalities is extremely desirable.	4.78	Strongly Agree
My providing EENC for every newborn at this hospital on regular basis will result in improving my clinical skills on birth assistance.	4.69	Strongly Agree
For me to keep up with my skills on birth assistance is extremely desirable	4.66	Strongly Agree
My providing EENC for every newborn at this hospital on regular basis will result in getting trust from patients.	4.07	Agree
I will want to provide full EENC for every newborn I assist birth on a regular basis	4.18	Agree
Whether or not I provide the EENC for every newborn I assist birth in this hospital on a regular basis is completely up to me	3.86	Agree
For me to provide the EENC for every newborn I assist birth in this hospital on a regular basis is extremely easy	4.02	Agree
Total Average Mean	4.40	Agree

*Strongly agree 5.0-4.5, Agree 4.49 – 3.50, Undecided 3.49 – 2.50, Disagree 2.49 – 1.50, Strongly Disagree 1.49- 1.0

Table 3 present the mean score and descriptive interpretation regarding the level of the nurses' performance in rendering essential newborn care in terms of attitude towards essential newborn care. It can be noted that there are four variables having a descriptive interpretation of "Strongly

Agree". This includes the highest mean score of 4.84 that deals with "for me, to provide the EENC for every newborn I assist birth in this hospital on a regular basis is beneficial" followed by mean score of 4.78 that deals with "For me, preventing newborn morbidities and mortalities is

extremely desirable". Then a mean score of 4.69 that deals with "My providing EENC for every newborn at this hospital on regular basis will result in improving my clinical skills on birth assistance" and finally a mean score of 4.66 that deals with "For me to keep up with my skills on birth assistance is extremely desirable".

thoroughly", "Breathing checked", and "Time of birth called out and recorded" and a mean score of 4.704 on the item "First wet towel discarded/ removed from the baby " which is both interpreted as Always practiced. The objective of the first 30 seconds of newborn life is to dry and provide warmth to the new-born to prevent hypothermia.

Table 4.1 shows Results of statistics manifested a mean score of 4.904 on the items "Baby dried

Table 4.1. Mean score and descriptive interpretation regarding the level of the nurse's performance in rendering essential newborn care in terms of practice of essential newborn care within the first 30 seconds of life

Variables	Mean score	Descriptive interpretation
Baby dried thoroughly	4.904	Always
Breathing checked	4.904	Always
First wet towel discarded/ removed from the baby	4.704	Always
Time of birth called out and recorded	4.904	Always

**Always 5.0-4.5, Very often 4.49 – 3.50, Often 3.49 – 2.50, Seldom 2.49 – 1.50, Never 1.49- 1.0*

Table 4.2. Mean score and descriptive interpretation regarding the level of the nurse's performance in rendering essential newborn care in terms of practice of essential newborn care in terms of the baby not breathing/crying at birth record

Variables	Mean score	Descriptive interpretation
Observe that the baby is not breathing/crying	4.84	Always
Baby dried thoroughly, still not crying	4.43	Very often
Changed to dry towel	4.88	Always
Check mouth/nose for secretions	4.88	Always
Mother's breast/chest NOT washed	4.73	Always
Suction mouth first then the nose	4.85	Always
Baby still not crying stimulate by rubbing the back	4.66	Always
.Baby still not breathing, cord cut (In 30 secs)	4.47	Very often
Explain to mother and move baby to resuscitation table	4.52	Always
Call for help	4.50	Always
Cord cut in between both clamps	4.87	Always
Newborn head placed in a slightly extended position	4.64	Always
Checks for secretions, clears the airway (mouth first then nose)	4.77	Always
Places correct size mask over the face of the newborn properly	4.74	Always
Checks the seal by ventilating twice, observes for chest rise	4.48	Very often
If chest still not rising, repositions mask to improve seal	4.69	Always
Ventilates at 40 breaths/ minute	4.64	Always
Ventilates for 1 minute, observes for spontaneous breathing	4.79	Always
If spontaneously breathing, move to routine care while observing breathing	4.69	Always
Baby still not breathing spontaneously, continue ventilation	4.66	Always
Check HR: if <100/min, ventilation continued and advanced care sought	4.76	Always
Total Average Mean	4.69	Always

**Always 5.0-4.5, Very often 4.49 – 3.50, Often 3.49 – 2.50, Seldom 2.49 – 1.50, Never 1.49- 1.0*

Table 4.2 presents the mean score and descriptive interpretation on the practice of Airway Management on Essential Newborn care. Data showed that the total average mean score

was 4.69 which is interpreted as Always practiced. The least mean computed were the variables "Baby dried thoroughly, still not crying" 4.43, "Baby still not breathing, cord cut (In 30 secs)" 4.47, and "Checks the seal by ventilating twice, observes for chest rise" 4.48 with descriptive interpretation of Very Often.

Untoward events may happen during birth, hence, the DOH Guideline for Essential Newborn Care also incorporated the Management of Airway/ Resuscitation for Newborn which is critical for the neonatal survival. In positioning the newborn to optimize the airway, the shoulders should be raised with posterior support and the head shifted to neutral or slightly extended. Mask ventilation is a potentially life-saving skill that is the cornerstone of effective airway management[10-12]. The mask should be sized appropriately to minimize air leak and in general should rest such that the inflatable cushioned portion superiorly encompasses the nose while not covering the eyes, and the inferior border rests on the chin. If the baby is vigorous (strong respiratory effort i.e., cry, good muscle tone, heart rate > 100 bpm) at birth, clear the airway by suctioning mouth first and then the nose with a bulb syringe or suction catheter. In every delivery room, an area should be allotted for neonatal resuscitation, with all the necessary equipment and drugs stored nearby. During every delivery there should be at least one person whose primary responsibility is the newborn. This person must be capable of initiating resuscitation, including administration of positive-pressure ventilation and chest compression.

Table 5.1 presents the composite table on the significance difference on the assessment of the respondents on the level of nurses' performance and knowledge of essential newborn care when grouped according to age profile using Analysis of Variance (ANOVA). It is clearly manifested from the table that the knowledge of essential newborn care when assessed by different ages have significant difference in terms of knowledge on the essential newborn care having a f-value of 2.638 with df of 115 with p-value of 0.027, practice on the essential newborn care having a f-value of 4.176 with df of 115 with p-value of 0.002 and baby not breathing having a f-value of 5.353 with df of 115 with p-value of 0.000 that the three variables have less than the alpha value 0.05 which means that the evidences gathered must reject the null hypothesis. This implies that there is a statistical difference between the two variables of the

study. Data denotes that knowledge on the essential newborn care, practice on the essential newborn care and baby not breathing have difference assessment when grouped in terms of age which means that age 20 years old have difference assessment with 30, 40 and 50 years old[21-23].

Table 5.2 presents the composite table on the significance difference on the assessment of the respondents on the level of nurses' performance when grouped according to newborn care training using Analysis of Variance (ANOVA). It is clearly manifested from the table that the knowledge of essential newborn care when assessed by different newborn care training do not have significant difference in terms of knowledge on the essential newborn care have an f-value of 1.148 with df of 115 with p-value of 0.321, attitude on the essential newborn care have an f-value of 2.037 with df of 115 with p-value of 0.135 and practice on the essential newborn care have an f-value of 0.497 with df of 115 with p-value of 0.609. The three variables have more than the alpha value 0.05 which means that the evidences gathered must accept the null hypothesis. This implies that there is a no statistical difference between all variables of the study. Data denoted that knowledge on the essential newborn care, attitude towards essential newborn care, and practice on the essential newborn care have no difference in assessment when grouped in terms of newborn care training which means that those who assessed the three parameters without or with training on newborn care training have the same degree or no difference[24,25,27].

Table 5.3 presents the composite table on the significance difference on the assessment of the respondents on the level of nurses' when grouped according to area of specialization using Analysis of Variance (ANOVA). It is clearly manifested from the table that the knowledge of essential newborn care when assessed by area of specialization have significant difference having a f-value of 3.535 with df of 115 with p-value of 0.009, practice on the essential newborn care having a f-value of 7.809 with df of 115 with p-value of 0.000, baby not breathing having a f-value of 6.162 with df of 115 with p-value of 0.000 and before discharge care having a f-value of 5.359 with df of 115 with p-value of 0.001 that the four variables have less than the alpha value 0.05 which means that the evidences gathered must reject the null hypothesis. This implies that there is a statistical

difference between the two variables of the study. Data denotes that knowledge on the essential newborn care, practice on the essential newborn care, baby not breathing and before discharge have significant difference when grouped in terms of area of specialization. This means that a nurse assigned in general unit ward have different assessment with those assigned in emergency room and other areas[13-16].

In summary, data denotes that age profile, and the area of specialization of nurses has significant difference in the knowledge and practice of essential newborn care. Attitude on the essential newborn care revealed no significant difference to any of the demographic profile presented.

Table 5.1. Significant difference on the assessment of the respondents on the level of nurses' performance when grouped according to age profile using Analysis of Variance (ANOVA)

Essential New-born Care	f-value	Df	Significant value	Decision	Remarks
Knowledge on the essential newborn care	2.638	115	0.027	Significant	Reject the Null Hypothesis
Attitude on the essential newborn care	2.246	115	0.054	Not Significant	Accept the Null Hypothesis
Practice on the essential newborn care	4.176	115	0.002	Significant	Reject the Null Hypothesis
Baby not Breathing	5.353	115	0.000	Significant	Reject the Null Hypothesis
Before discharge	2.223	115	0.057	Not Significant	Accept the Null Hypothesis

Table 5.2. Significant difference on the assessment of the respondents on the level of nurses' performance when grouped according to new born care training profile using Analysis of Variance (ANOVA)

Essential New-born Care	f-value	Df	Significant value	Decision	Remarks
Knowledge on the essential newborn care	1.148	115	0.321	Not Significant	Accept the Null Hypothesis
Attitude on the essential newborn care	2.037	115	0.135	Not Significant	Accept the Null Hypothesis
Practice on the essential newborn care	0.497	115	0.609	Not Significant	Accept the Null Hypothesis
Baby not Breathing	0.559	115	0.573	Not Significant	Accept the Null Hypothesis
Before discharge	3.875	115	0.023	Significant	Reject the Null Hypothesis

Table 5.3. Significant difference on the assessment of the respondents on the level of nurses' performance when grouped according to area of specialization profile using Analysis of Variance (ANOVA)

Essential New-born Care	f-value	Df	Significant value	Decision	Remarks
Knowledge on the essential newborn care	3.535	115	0.009	Significant	Reject the Null Hypothesis
Attitude on the essential newborn care	1.879	115	0.119	Not Significant	Accept the Null Hypothesis
Practice on the essential newborn care	7.809	115	0.000	Significant	Reject the Null Hypothesis
Baby not Breathing	6.162	115	0.000	Significant	Reject the Null Hypothesis
Before discharge	5.359	115	0.001	Significant	Reject the Null Hypothesis

Further analysis showed that, when considering the age profile and area of specialization of nurses, Statistics showed that: Ages 31-35 years old gained the highest mean (13.7 out of 20) followed by 46-50 years old (13.3 out of 20) then 26-30 (13.09 out of 20), then 41-45 years old (12.9 out of 20), then 21-25 years old (12.5 out of 20), last are ages 36-40 years old and 51 years old above (12 out of 20). Highest mean score came from Emergency room nurse (13.63 out of 20), followed by Pediatric and DR/Labor room nurse (13 out of 20), next is the general unit nurse (12.9 out of 20), and last is the maternity ward nurse (12 out of 20).

4. CONCLUSION AND RECOMMENDATIONS

The level of district nurses' Knowledge on Essential Newborn care and some of the vital aspects on it is low due to lack of proper training of the involved nurses. Though the Practice of Essential Newborn Care resulted as *Always* or highly being practiced, some unnecessary and deficient procedures need to be re-taught to achieve excellent execution of ENC protocol. The age and the area of specialization of nurses have significant difference in the knowledge and practice of essential newborn care. Attitude on the essential newborn care revealed no significant difference to any of the demographic profile presented. A number of implications may be considered by the District Hospital nursing management in further improving the knowledge and practice of their nurses in ENC.

CONSENT

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

ACKNOWLEDGEMENTS

The authors would like to extend gratitude to our University for supporting us in our research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Liao AO, Manalon RC. Theory and practice: identifying the gaps in Essential newborn Care Practice of Nursing and Midwifery Students during their Clinical Practicum. Puerto Princesa: Faculty of the College of Nursing and Health Science; 2015.
2. Negussie BB, Hailu FB, Megenta AD. Knowledge and practice of essential newborn care and associated factors among nurses and midwives working at health centers in JimmaZone, Ethiopia, 2016. *J Nurs Care*. 2018;07(1). DOI: 10.4172/2167-1168.1000446
3. Horiuchi S, Rattana S, Saysanasongkham B, Kounnavongsa O, Kubota S, Cayrol J et al. Study protocol of a cluster randomized controlled trial to evaluate effectiveness of a system for maintaining high-quality early essential newborn care in Lao PDR. *BMC Health Serv Res*. 2018;18(1):489. DOI: 10.1186/s12913-018-3311-7, PMID 29940953
4. Department of Health. Essential newborn care new guideline 2015. PPD compendium of Philippine Medicine. 16th ed.; 2014-5.
5. World Health Organization. Protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services. Geneva; 2011. Available:<http://www.who.int/nutrition/publications/guidelines/breastfeeding-facilities-maternity-newborn/en/>.
6. Chichiabellu TY, Mekonnen B, Astawesegn FH, Demissie BW, Anjulo AA. Essential newborn care practices and associated factors among home delivered mothers in Damot pulasa Woreda, southern Ethiopia. *Reprod Health*. 2018;15(1):1-11. DOI: 10.1186/s12978-018-0609-1
7. Mohamed S, Faheim S, Farg D, Hassan H. The relation between trunk-to-head bathing and the traditional head-to-trunk bathing on newborns' outcome. *Am Res J Nurs*. 2018;4(1):1-12. DOI: 10.21694/2379-2922.18002
8. Hassan HE, EL-Kholy GA, Ateya AA, Hassan AA. Breast feeding knowledge and

- practices among primiparous women with Caesarean section: impact on breast engorgement in upper Egypt. *Commun Soc Media*. 2020;3(2):34-78.
DOI: 10.22158/csm.v3n2p34
9. Faheim SS, Hassan HE, Gamel WMA. Topical Application of Human Milk versus alcohol and Povidine-iodine on Clinical Outcomes of Umbilical Cord in Healthy Newborn: impact of an Educational Program of Mothers' Knowledge and Practice regarding Umbilical Cord Care. *Int J Stud Nurs*. 2019;4(2):35-51.
DOI: 10.20849/ijsn.v4i2.573
 10. Saaka M, Ali F, Vuu F. Prevalence and determinants of essential newborn care practices in the Lawra District of Ghana. *BMC Pediatr*. 2018;18(1):173.
DOI: 10.1186/s12887-018-1145-4, PMID 29793543
 11. Elzeblawy Hassan H. Evidence-based practice in midwifery and maternity nursing for excellent quality of care outcomes. *Am J Nurs Res*. 2020;8(6):606-7.
DOI: 10.12691/ajnr-8-6-3
 12. Gaber Zagloul M, Saied Hassan S, Ali Abd Elmoaty Sheha E, Elzeblawy Hassan H. Postnatal evidence based nursing practice: Effect of an educational guideline on internship Student's knowledge and skills. *Egypt J Health Care*. 2022;13(2):437-47.
DOI: 10.21608/EJHC.2022.230408
 13. Philippine Statistics Authority (PSA), ICF. Key findings from the Philippines national demographic and health survey 2017. Quezon City, Philippines, and Rockville, MD: PSA and IMB Community Foundation; 2018.
Available:https://psa.gov.ph/sites/default/files/2017%20PHILIPPINES%20NDHS%20KEY%20FINDINGS_092518.pdf
 14. Hassan H. The impact of evidence-based nursing as the foundation for professional maternity nursing practices. *Open Access J Reprod Syst Sex Disord. OAJRSD.MS*. 2019;2(2):195-7:ID.000135.
DOI: 10.32474/OAJRSD.2019.02.000135
 15. Tosif S, Jatobatu A, Maepioh A, Gray A, Sobel H, Mannava P et al. Healthcare worker knowledge and skills following coaching in WHO early essential newborn care program in the Solomon Islands: A prospective multi-site cohort study. *BMC Pregnancy Childbirth*. 2020;20(1):84.
DOI: 10.1186/s12884-020-2739-z, PMID 32033598
 16. Hassan HE, Nasr EH. Improving nurses' knowledge and skills regarding tocolytics for inhibiting preterm labor. *Clin Nurs Stud*. 2017;5(1):1-12.
DOI: 10.5430/cns.v5n1p1
 17. Salvage J, White J. Our future is global: nursing leadership and global health. *Rev Latino-Am. Enfermagem*. 2020;28:e3339.
DOI: 10.1590/1518-8345.4542.3339
 18. Alipour Z, Eskandari N, Abbasi M, Raisi M, Bakouei S. Structural challenges in the health domain of the health system reform: A qualitative study. *J Educ Health Promot*. 2019;8:55.
DOI: 10.4103/jehp.jehp_186_18, PMID 31008122
 19. Hassan H, Zahran K, Youness E, Nady F. Pregnant women's awareness, intention and compliance regarding folic acid usage for prevention of neural tube defects according to health belief model in Beni-Suef city. *Pyrex J Nurs Midwif*. 2015;1(3): 13-26.
 20. Farg D, Hassan H. Study hyperemesis Gravidarum requiring hospital admission during pregnancy: effect of nursing implication on its progress. *Am J Nurs Res*. 2019;7(3):328-41.
DOI: 10.12691/ajnr-7-3-14
 21. Sheha E, Hassan H, Genedy A, Hassanine Sh. Effect of educational program on mother's knowledge and practice regarding hepatitis C virus in rural areas. *Am J Nurs Res*. 2020;8(3): 303-10.
DOI: 10.12691/ajnr-8-3-1
 22. Hassan H, Nady F, Youns E, Zahran K. Call for change level of knowledge, awareness and attitude to follow A high folate diet among pregnant women. *IOSR JNHS*. 2016;5(1):93-100.
DOI: 10.9790/1959-051293100
 23. Mostafa H, Yousef F, Hassan H. Health related quality of life educational interventions: effect on chronic hepatitis C patients'. *Saudi J Nurs Health Care*. 2018;1(2):56-67.
 24. Zagloul MC, Naser EG, Hassan HE. Influence of hot compresses versus cabbage leaves on engorged breast in early puerperium. *Int J Stud Nurs*. 2020; 5(2):7-14.
DOI: 10.20849/ijsn.v5i2.740
 25. Hassan H, Said S, Hassanine Sh. Disparities of prevalence and causes of maternal antenatal anxiety among

- primigravida pregnant women in Egypt. Am Res J Nurs. 2017;3(1):1-15.
DOI: 10.9790/1959-051293100
26. Hassan H, Malk R, Abdelhamed A, Genedy A. Infection control knowledge and practices: program management in labor units according to standard infection control precautions in northern upper Egypt. Am J Nurs Res. 2020;8(4):412-25.
DOI: 10.12691/ajnr-8-4-1
27. Chadha IA. Neonatal resuscitation: current issues. Indian J Anaesth. 2010;54(5):428-38.
DOI: 10.4103/0019-5049.71042, PMID 21189881