

Study on Crop Diversification based on Long Term Rain fall analysis of Dry farming tracts in southern India

Abstract. This study was conducted based on 30 years of rainfall data from 1985 to 2015 from the regions of semiarid districts of Karnataka, Telangana and Rayalaseema region of Andhra Pradesh states to recommend the alternate crops. This is based on the rainfall variability and potential evapotranspiration (PET) in kharif and rabi seasons. Moisture-sensitive crops, especially rice, sugarcane and banana must be taken in high rainfall areas and low moisture-sensitive crops can go at the time of rainfall shortages and vice versa to avoid farm risk at farm level and optimizing production. The crops which are suggested based on rainfall and PET can thrive well, grow very efficiently and give reasonable yields to farmers under limited rainfall situations and these are of drought-tolerant and drought resistant and short-duration types. In the rabi season, most of these regions received less rainfall (<200mm), particularly in Telangana and the North interior Karnataka regions. Under this situation, avoiding more water demanded crops like cotton and sunflower crops. Instead of these crops, less water required short - duration and drought tolerant Bengal gram and Cowpea crops were suggested. These crops can survive well and give good optimum yields without crop failures.

Keywords-Diversification, Drought resistant, Potential evapotranspiration, Rainfall.

1. INTRODUCTION

“Crop diversification: It refers to the addition of new crops or cropping systems to agricultural production on a particular farm, taking into account the different returns from value-added crops with complementary marketing opportunities” (Khanam et al, 2018). “Crop Diversification refers to a shift from the regional dominance of one crop to regional production of a number of crops, to meet ever increasing demand of cereals, pulses, vegetables, fruits, oilseeds, fibres, fodder, grasses etc. It aims to improve soil health and to maintain dynamic equilibrium of the agro-ecosystem. In the instant case, crop diversification is intended to promote technological innovations for sustainable agriculture and enable farmers to choose crop alternatives for increased productivity and income”. (crop diversification programme 2013-14).

Traditionally, diversification was used more in the context of a subsistence kind of farming wherein farmers grow many crops on their farms. Farmers all over the world, especially in the developing countries, try to grow various crops in their holdings in an agricultural year. The level of crop diversification largely depends on the geo-climatic/socio-economic conditions and technological development in a region. Moreover, rich farmers prefer to specialize in agricultural enterprises, while the poor and subsistent farmers are generally more interested in diversification of crops.

“Under weather uncertainty and almost exclusive dependence of small holders on rainfall for productivity, several studies have investigated the nature and degree of crop riskiness in relation to the presence of production and market risks” (Fafchamps 1992; Haile 2007; Dercon 1996). It should be noted, however, that these earlier studies either rely on subjective measures of the riskiness of the crops or focus on selected major crops. An objective way of measuring the riskiness of individual crops and aggregating them (in a multiple-crop setting) allows a more accurate measurement of the contribution of individual crops to the riskiness, incorporating the mutual interdependence across crops at a farm-household level. It was also observed that costly risk-coping mechanisms were commonly adopted in response, leading to the recommendation that “livelihood diversification programmes be scaled up to reduce dependency on agriculture-based activities in view of climate shocks” (VAC, 2015).

“Diversification means taking on less profitable but less risky crops. It is an example of conservative crop production strategies and one such risk coping mechanism” (Benin *et al.*, 2004; Morduch, 2002). “Diversification is especially important to hedge against weather risk” (Kurukulasuriya and Mendelsohn 2006). “In a pioneering study of crop choices under multivariate risk, Fafchamps (1992) showed that crop diversification, which is a characteristic feature of small farmers in developing countries, is a response to high variance in food prices and other risks that they are poorly insured against. Similarly, using data from the Punjab area of Pakistan”, Kurosaki Fafchamps (1992) demonstrated that “farmer crop choices are dependent on price and yield risk. Crop diversity, crop sequencing or rotation (Amedeet al., 2001; Benin et al., 2004), and intercropping (Corbeels et al., 2000) are all traditional

methods for increasing soil productivity (renewing soil nutrients) and maximising return from cultivated land in uncertain conditions. Farmers also commonly plant varieties that mature earlier in the season (short-season crops) and protect crops from moisture shortages and yield loss". Di Falco and Chavas (2009) showed that "greater diversity can reduce the risk of crop failure". Seo and Mendelsohn (2008) and Kurukulasuriya and Mendelsohn (2006) also looked at "the climate sensitivity of crop choices. Using cross-country data from Latin America and Africa, understanding household-level crop choices can generate important information about how farm households change the riskiness of their crop composition in reaction to weather-related risk". Haile (2007) showed that "Ethiopian farmers choose crops most suited to a specific rainfall condition as a strategy for coping with unpredictable rainfall". "In times of low rainfall, farmers predominantly choose moisture- and stress-tolerant crops and not moisture-sensitive crops. All aspects of food security may be potentially threatened by the effects of changes in climate, which will include food availability, access, utilization, and stability" (e.g., Challinor et al., 2010; IPCC, 2014).

"The semi-arid tropics are characterised by low and highly variable rainfall in space and time, limiting potential crop yields in these areas" (Graef and Haigis, 2001). Very few studies have tried to quantify the spatial and temporal variability of rainfall in the semi-arid tropics. The high degree of rainfall variability, when combined with the relatively low asset base of most rural households, restricts household crop management strategies and overall crop water productivity.

KARNATAKA

"Agriculture plays an important role in the overall growth of Karnataka's economy, despite a fall in its share of the state's gross domestic product. It is characterised by wide crop diversification and remains highly dependent on the vagaries of the southwest monsoon. Rainfall plays an important role in crop production in Karnataka, where more than 70 percent of the cropped area is rainfed. Districts like Bijapur, Bellary, Bagalkot, Koppal, Gadag, Raichur, and Chitra Durga can be classified as low rainfall districts where the average annual rainfall (1998–2010) is less than 650 mm. The normal number of rainy days varied from less than 40 in low rainfall districts to more than 80 in high rainfall districts". (Economic survey of Karnataka 2011–12).

TELANGANA

Telangana is a semi-arid area and has a predominantly hot and dry climate. Over 80% of the original forest cover has been cleared for agriculture, timber harvesting, or cattle grazing. The more humid Eastern_Highlands' moist_deciduous_forests cover the Eastern_Ghats in the eastern part of the state. With improved agricultural technology, cropping system diversity has decreased from 23 to 10 crops in the kharif (monsoon) season. These changes are also associated with a shift from intercropping to monocrop-based systems. The results indicate that these changes in cropping systems have resulted in a decreased management of production risk with increased investments in agriculture and less flexibility in the decision-making process on crops. Because of the differences in agro-ecology, socioeconomic characteristics, and resource availability, these regions must be distinguished in order to determine their agricultural development strategies.

Objectives of the study

state the meaning of diversification in agriculture;

explain How rainfall trend varies in semi-arid regions of Karnataka, Telangana and Rayalaseema.

Discuss the effect of diversification on soil health and water resources, the sustainability of production.

Suggesting crops Based on Rainfall amount, Potential Evapotranspiration and water requirement values of different crops.

DIVERSIFICATION OPPORTUNITIES IN INDIAN AGRICULTURE

suitability of the agro-climatic situation

Availability of appropriate technology

Some remunerative avenues of diversification

1. Diversification through pulses and oilseeds
2. b) Diversification with fodder crops
3. Diversification with horticultural crops
4. d) Diversification with medicinal and aromatic plants

Climate change is projected to cause increased temperatures and altered precipitation in the near future. Moreover, the weather extremes are also projected to increase in the future, which can have a detrimental effect on agricultural yields (Malhi et al., 2021).

Crop diversification is recognised as an effective mitigation and adaptation strategy (Lakhran et al., 2017).

Climate shocks can be mitigated through crop diversification, and the benefits of adaptation are more visible in the long run (Birthal and Hazrana, 2019).

Monocropping cultivation, combined with increased fertiliser and pesticide use over the years, has resulted in deteriorating soil health, increased insect pest infestation, and yield stagnation in some crops. This necessitates the diversification of crops so as to ensure the income stability of the farmers through the reduction of production and market risk. Crop diversification has proven useful in maintaining soil health and biodiversity and ensuring sustainability in the arena of climate change.

2. Material and methods

The study was conducted in semi-arid regions of Telangana, Karnataka states, and the Rayalaseema region of Andhra Pradesh for the period 1985 to 2015. Telangana, a state on the south-eastern coast of India, has a net cropped area of 4.5 million hectares, of which 2.5 million hectares (56% of the net cropped area) are under rainfed agriculture (Directorate of Economics and Statistics 2011). Telangana is located between latitudes 18.1124°N and 79.0193°E longitudes; Rayalaseema is located between latitudes 15.7722°N and 78.0641°E longitudes; and Karnataka is located between latitudes 15.3173°N and 75.7139°E longitudes. **Quantitative Analysis:** A 30-year rainfall data set (1985–2015) is used to compute seasonal rainfall in Telangana and Karnataka states, as well as the Rayalaseema region of Andhra Pradesh, from the Hydrology Section at IMD Pune. Each district trend was taken from each region, and the tendency of the trend was taken using Microsoft Excel. Drought severity is conventionally assessed by drought indices. A powerful drought index, the Reconnaissance Drought Index (RDI), is gaining wide acceptance, mainly in the arid and semiarid climatic regions. It is achieved by potential evapotranspiration and precipitation. Potential evapotranspiration was calculated with the Penman and Monteith equations. Based on the values of precipitation and potential evapotranspiration, appropriate crops were suggested. Soil condition, moisture holding capacity, texture details, and the minimum water requirement of crops to complete their life cycle for a season are collected from CRIDA (the Central Research Institute for Dryland Agriculture).

3. Results and Discussion

Based on 30 years of rainfall data from the regions of semiarid districts of Karnataka, Telangana, and the Rayalaseema region of Andhra Pradesh states, the following trends were observed in corresponding regions: The rainfall trends of different districts of south interior Karnataka during the *kharif* season, i.e., Chickmangalore, Mysore, Kodagu, Hassan, and Chitradurga, showed an increasing trend, whereas Shivamogga, Bangalore urban, and Chamarajanagar districts showed a decreasing trend. Kolar, Tumkur, and Bangalore rural districts showed no trend. In the *Rabi* season, Kolar, Tumkur, and Mysore districts showed an increasing trend, whereas Shivamogga district showed a decreasing trend. There was no trend in the remaining districts of southern interior Karnataka. The rainfall trends of different districts of north interior Karnataka during the *kharif* season, i.e., Haveri, Gadag, Gulbarga, and Koppal districts, showed an increasing trend, whereas decreasing trends were observed in Raichur, Dharwad, Bidar, and Belgaum districts. Bagalkot and Bijapur districts showed no trend. During the *rabi* season, Koppal and Belgaum districts showed an increasing trend, but Haveri, Raichur, Gulbarga, Dharwad, Bidar, and Bijapur districts showed a decreasing trend. Gadag and Bagalkot districts showed no trend. In terms of rainfall trends in different districts of Rayalaseema during the *kharif* season, Anantapur district showed a decreasing trend and Kurnool district showed an increasing trend, whereas Chittoor and Kadapa districts showed no trend. In the case of *rabi* season rainfall, Anantapur district showed an increasing and Kurnool district showed a decreasing trend. where Chittoor and Kadapa districts showed no trend. In case of the rainfall trends in different districts of Telangana during the *kharif* season, Karimnagar, Adilabad, Medak, Ranga Reddy, Nalgonda, Nizamabad, and Medak showed a decreasing trend, whereas Hyderabad, Khammam, and Mahbubnagar districts showed an increasing trend, and Warangal district showed no trend. In case of the rainfall during the *rabi* season, Mahbubnagar, Adilabad, Karimnagar, Medak, Nalgonda, Nizamabad, Ranga Reddy, Warangal, and Hyderabad districts showed a decreasing trend, whereas Khammam district showed no trend.

In semiarid regions of Karnataka, Telangana, and Rayalaseema districts, agriculture is almost exclusively rainfed, so rainfall variability comprises an important source of uncertainty in agricultural production decisions. A better understanding of production risk and its management is important to help farmers make informed and critical decisions about their crops because their welfare depends on their ability to withstand the risk of crop loss. In line with this, the study demonstrates crop selection as a risk management mechanism when crop insurance is limited or non-existent. Our central premise is that in an

alternate-cropping system, the crops chosen are likely to be sensitive to weather risk, measured by *kharif* and *rabi* rainfall variability.

“The ability to adjust in response to a change in temperature is common in plants and includes both higher and lower values relative to the optimum temperature. Besides, the availability of liquid water depends not only on the amount of water present but also on temperature” (Tiziana *et al.*, 2000). Heat and drought are undoubtedly the two most important stresses, having a huge impact on the growth and productivity of the crops (Google.com). “The crop water stress index is a means of irrigation scheduling and crop water stress quantification based on canopy temperature measurements and prevailing meteorological conditions. Plant temperature is an indicator of plant water status because stomata close in response to soil water depletion, causing a decrease in water uptake and an increase in leaf temperature” (Payero *et al.*, 2006). “Inadequate rainfall and soil-bound water lead to water deficiency in crops” (Sekhonet *et al.*, 2010; Vadezet *et al.*, 2011). “Plant-water relations and water-stress tolerance at the scale of physiology and molecular biology can significantly improve plant productivity and environmental quality”. (Liu *et al.*, 2005; Shao *et al.*, 2008).

Drought severity is conventionally assessed by drought indices. Several drought indices with varying complexity have been used in many geographical areas. Recently, a powerful drought index, the Reconnaissance Drought Index (RDI), has gained wide acceptance, mainly in the arid and semiarid climatic regions. Since RDI is based both on precipitation and potential evapotranspiration (PET), it is interesting to assess the effect of the PET calculation method on the drought severity characterization obtained by RDI. The FAO Penman–Monteith method is used as a reference method.

In south and north interior Karnataka, Telangana, and Rayalaseema areas, PET was calculated based on the Penman and Monteith equation. In districts wherever rainfall is less than potential evapotranspiration, one or two irrigations should be given at critical stages of crop growth.

In the districts of south and north interior Karnataka, Telangana, and Rayalaseema regions, the existing crops under cultivation are shown in tables 1–8 in both the *kharif* and *rabi* seasons. According to rainfall patterns and potential evapotranspiration in different districts, suitable crops were suggested. The crops that are suggested based on rainfall and potential evapotranspiration can thrive well, grow very efficiently, and give reasonable yields to farmers in limited rainfall situations. Drought-tolerant and drought-resistant short-duration crops were mostly recommended.

4.SUMMARY AND CONCLUSIONS

Based on 30 years of average rainfall and potential evapotranspiration losses in regions of semiarid districts of Karnataka, Telangana, and the Rayalaseema region of Andhra Pradesh states, the results indicate that the rainfall variability both in the *kharif* and *rabi* seasons means that moisture-sensitive crops like rice, sugarcane, and banana have to be taken up in high rainfall areas and in areas with good irrigation facilities, and farmers should go for less moisture-sensitive crops at times of rainfall shortages and vice versa to avoid risk at farm level.

The timely arrival of southwest monsoon rains is crucial for seed-bed preparation for short- and long-duration *kharif* crops such as maize, sorghum, and rice. This implies that favourable southwest monsoon rains are important for crop production.

In semiarid regions of Karnataka, Telangana, and Rayalaseema districts of Andhra Pradesh, agriculture is almost entirely rainfed, so rainfall variability becomes an important factor of uncertainty in agricultural production decisions. A better understanding of production risk and its management is important to help farmers become informed and to take critical decisions about their crops because their welfare depends on their ability to withstand the risk of crop loss. If the monsoon rains stop early in the season, normal sowing of short-duration upland rice, blackgram, and sesame crops can be resumed. If the rain ceases very early, i.e., by the end of August or the first week of September, only fodder crops or grain legumes could be harvested. Depending upon the soil moisture condition, relay sowing of crops like chickpea, lentil, mustard, linseed, and barley could be done in the *rabi* season.

The crops that are suggested based on rainfall and potential evapotranspiration can thrive well, grow very efficiently, and give reasonable yields to farmers under limited rainfall situations. In districts where the rainfall is less than the potential evapotranspiration, one or two irrigations may be given at critical stages of crop growth.

5. Advantages of diversification

- More Income, Poverty Alleviation, and Social Upliftment
- Improved Soil Health
- More Employment Generation at the Farm
- Effective and Profitable Utilization of Farm Waste and By-Products
- Risk Reduction

ACKNOWLEDGEMENT

I gratefully acknowledge the India Meteorological Department for encouraging and supporting me to do this study. The contents and views expressed in this paper are those of the authors and do not necessarily reflect the views of the organisation they belong to.

CONFLICT OF INTEREST

None.

AUTHOR CONTRIBUTION

The first author contributed to the manuscript's collection and writing. 2nd and 3rd Authors: Data Collection and Preparation of Some Figures Data plotting by the fourth author

REFERENCES

- Amede, T., T. Belachew, and E. Geta. 2001. Reversing the Degradation of Arable Land in the And Statistics, Government of Karnataka, Bangalore.
- Benin, S., M. Smale, J. Pender, B. Gebremedhin, and S. Ehui. 2004. The Economic Census of India. 2011. Government of India.
- Birthal, P.S. and Hazrana, J. (2019). Crop diversification and resilience of agriculture to climate shocks. *Agricultural Systems*. 173: 345-354.
- Chaves MM, Pereira JS, Maroco J, Rodrigues ML, Ricardo CPP, Osorio ML, Carvalho I, Faria T, Pinheiro C (2002) How plants cope with waterstress in the field? Photosynthesis and growth. *Ann Bot* 89(7): 906–917.
- Corbeels, M., A. Shiferaw, and M. Haile. 2000. Farmers' Knowledge of Soil Fertility and Dercon, S. 1996. Risk, Crop Choice, and Savings: Evidence from Tanzania. *Economic Determinants of Cereal Crop Diversity on Farms in the Ethiopian Highlands. Agricultural Economics*31(2–3): 197–208.
- Di Falco, S., and J.-P. Chavas. 2009. On Crop Biodiversity, Risk Exposure, and Food Security Directorate of Economics and Statistics, Government of Karnataka, Bangalore.
- effects on farmers' management strategies. *Journal of Arid Environments* (2001) 48: 221–231
- Environmental Economics and Policy in Africa (CEEPA), University of Pretoria. Ethiopian Highlands. Managing Africa's Soil, no. 23. London: IIED.
- Fafchamps, M. 1992. Cash Crop Production, Food Price Volatility, and Rural Market
- Government of Karnataka (2012). "Economic survey 2011-12". Directorate of Economics And Statistics, Government of Karnataka.
- Government of Telangana (2012). "Economic survey 2011-12". Directorate of Economics and Statistics, Government of Telangana, Hyderabad.
- Graef F. and Haigis J. 2001. Spatial and temporal rainfall variability in the Sahel and its
- H.Vangelis D.TigkasG.TsakirisThe effect of PET method on Reconnaissance Drought Index RDI) calculation *Journal of Arid Environments* Volume 88, January 2013, Pages130-140
- Haile, N. 2007. An Economic Analysis of Farmers' Risk Attitudes and Farm Households'

<http://www.entrepreneur.com/tradejournals/article/203135307.html>. Accessed August 2011. In Africa. CEEPA Discussion Paper, no. 26. Pretoria, South Africa: Center for In the Highlands of Ethiopia. *American Journal of Agricultural Economics* 91(3).

Khanam, R., Bhaduri, D. and Nayak, A.K. 2018. Crop diversification: an important way-out for doubling farmers' income. *Indian farming*. 68(1): 31-32. Vol.1 Issue-8, APRIL 2021 (e-ISSN: 2582-8223) www.justagriculture.in Page 4

Kurukulasuriya, P., and R. Mendelsohn. 2006. Crop Selection: Adapting to Climate Change Integration in the Third World. *American Journal of Agricultural Economics* 74(1): 90–99.

IPCC. 2012. Managing the risks of extreme events and disasters to advance climate change adaptation. A special report of Working Groups I and II of the Intergovernmental Panel on Climate Change. C.B. Field, V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G-K. Plattner, S.K. Allen, M. Tignor & P.M. Midgley, eds., Cambridge, UK, and New York, USA, Cambridge University Press.

Liu *et al.*, 2005; Shao *et al.*, 2008. Water-deficit stress-induced anatomical changes in higher plants; Water-deficit stress-Induced anatomical changes in higher plants. *Comptes Rendus Biologies* 331(3):215-25.

Mahtab, S. B., (2000), Diversification of Agriculture for Human Nutrition. *Current Science*, Vol. 78(7):1-2.

Malhi, G.S., Kaur, M. and Kaushik, P. (2021). Impact of climate change on agriculture and its mitigation strategies: a review. *Sustainability*. 13: 1318. <https://doi.org/10.3390/su13031318>.

Odekunle T. O., Orinmoogunje I. O. O. and Ayanlade A. 2007. Application of GIS to assess rainfall variability impacts on crop yield in Guinean Savanna part of Nigeria. *African Journal of Biotechnology* 6 (18): 2100-2113.

Responses to Rainfall Risk in Tigray, Northern Ethiopia. PhD thesis, Mansholt Graduate School of Social Sciences, Wageningen University, the Netherlands.

Sekhon HS, Singh G, Sharma P, Bains TS (2010) Water use efficiency under stress environments. In: *Climate Change and Management of Cool Season Grain Legume Crops*, Yadav SS, McNeil DL, Redden R, Patil SA (Eds) Springer Press, Dordrecht-Heidelberg-London-New York.

Seo, N., and R. Mendelsohn. 2008. Measuring Impacts and Adaptations to Climate Change: A Structural Ricardian Model of African Livestock Management. *Agricultural Economics* 38: 1–15.

Shukla, A.K, Shukla, N.D. and Singh, V.K., (2002), Farming System Approach for Food Security and Sustained Rural Economy. *Fertiliser News*, Vol. 47(11): 55-62 & 65-66.

Vadez V, Kholova J, Choudary S, Krishnamurthy L, Kumar PR, Turner NC (2011) Whole plant response to drought under climate Change. *Crop adaptation to Climate Change*

UNDER PEER REVIEW

Table:1**SOUTH INTERIOR KARNATAKA Kharif**

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL (mm) | PET (mm) | RECOMMENDED CROPS |
|------|----------------|---|----------------|----------|--|
| 1 | CHICKMANGALUR | Ragi, Paddy, Sunflower, Jowar, Horse gram | 1307.4 | 416.1 | Ragi,Paddy, Sunflower, Jowar,Red gram |
| 2 | KOLAR | Ragi,Ground nut, Maize, Red gram and Pulses | 368.8 | 595.2 | Ragi, Red gram and Pulses |
| 3 | TUMKUR | Ragi, Maize, Redgram, Ground nut and Horsegram | 353.3 | 460.2 | Ragi, Redgram, and Horsegram |
| 4 | SHIMOGA | Paddy, Maize, Sugarcane,and Cotton | 2176.4 | 41.1 | Paddy, Maize, Sugarcane,and Cotton |
| 5 | MYSORE | Maize,Ragi, Cotton and pulses | 351.5 | 429 | Ragi and pulses |
| 6 | KODAGU | Horti crops | 2186.4 | 416.1 | Bajra and Sesame |
| 7 | BANGLORE URBAN | | 445.3 | 460.2 | |
| 8 | HASSAN | Paddy, Ragi, Maize and Cowpea | 805.8 | 435.9 | Ragi, Maize and Cowpea |
| 9 | CHITHRADURGA | Groundnut, maize, Ragi, Sunflower, Jowar and Redgram | 299.7 | 537.7 | Bajra, Bengal gram and Sesame |
| 10 | CHAMARAJNAGAR | Ragi, Paddy, Sunflower, Horsegram Sorghum and Bengal gram | 331.1 | 424.2 | Ragi, Soybean, Horsegram and Bengal gram |
| 11 | BANGLORE RURAL | Ragi, Maize, Redgram and Ground nut | 436.2 | 456.4 | Ragi, Redgram and Soybean |

Table:2**SOUTH INTERIOR KARNATAKA Rabi**

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL (mm) | PET (mm) | RECOMMENDED CROPS |
|------|---------------|----------------------------------|----------------|----------|-------------------------|
| 1 | CHICKMANGALUR | Black gram, Green gram and Bajra | 233 | 613.2 | Bengal gram |
| 2 | KOLAR | Horse gram | 212.9 | 609 | Bengal gram |
| 3 | TUMKUR | Bengal gram | 189.3 | 582.5 | Horsegram |
| 4 | SHIMOGA | Fallow | 203.1 | 613.2 | Bengal gram |
| 5 | MYSORE | Fallow | 210.4 | 593.5 | Bengal gram |
| 6 | KODAGU | Fallow | 305.8 | 602 | Bengal gram and Soybean |

| | | | | | |
|----|----------------|---|-------|-------|-------------------------------------|
| 7 | BANGLORE URBAN | Fallow | 219.3 | 582.5 | |
| 8 | HASSAN | Ragi, Maize and Cowpea | 236.6 | 596.6 | Blackgram Green grm and Bengal gram |
| 9 | CHITHRADURGA | | 162.5 | 609.3 | Bengal gram |
| 10 | CHAMARAJNAGAR | Sunflower, Sorghum, Bengal gram and Horsegram | 266 | 544.5 | Bengal gram and Horsegram |
| 11 | BANGLORERURAL | | 209.9 | 571.7 | Bengalgram |

*Water requirement of Bengal gram is 150mm, Black gram 250mm, Green gram 250mm Horsegram 200mm.

Table:3 NORTH INTERIOR KARNATAKA *Kharif*

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL | PET(mm) | RECOMMENDED CROPS |
|------|-----------|--|-----------|---------|---|
| 1 | GADAG | Greengram, Groundnut and Sorghum | 320.4 | 422.7 | Greengram and Bajra |
| 2 | HAVERI | Maize, Oilseed, Sorghum, Paddy and Cotton | 454.8 | 410.9 | Maize, Sunflower and Safflower and Sorghum |
| 3 | RAICHUR | Sunflower, Sorghum, Bajra, Groundnut, Cotton, Redgram | 402.4 | 633.1 | Sunflower and Redgram |
| 4 | KOPPAL | Sunflower, Sorghum, Bajra, Groundnut and Maize | 346.2 | 551.4 | Greengram and Bajra |
| 5 | GULBARGA | Redgram, sorghum, Sunflower and Bajra | 507 | 560.2 | Redgram, sorghum, Sunflower and Bajra |
| 6 | DHARWAD | Cotton, Chilli, Sorghum, Groundnut, Paddy, Soybean and Greengram | 460.6 | 453.1 | Redgram, Sorghum, Soybean and Greengram |
| 7 | BIDAR | Sorghum, Redgram, Soybean, Blackgram, Green gram | 616.6 | 753.7 | Cotton, Sorghum, Groundnut, Redgram and Soybean |
| 8 | BIJAPUR | Redgram, Sunflower and Bajra | 349.1 | 483.1 | Soybean and Bajra |
| 9 | BELGAUM | Soybean, Maize, Groundnut, Sorghum and Cotton | 780.4 | 453.1 | Maize, Groundnut, Sorghum and Cotton |
| 10 | BAGALKOTE | Gram, Sorghum and Groundnut | 330.6 | 540.1 | Gram, Greengram and Bajra |

Table:4 NORTH INTERIOR KARNATAKA *Rabi*

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL(mm) | PET(mm) | RECOMMENDED CROPS |
|------|----------|---------------|---------------|---------|-------------------|
|------|----------|---------------|---------------|---------|-------------------|

| | | | | | |
|----|-----------|--|-------|-------|------------------------|
| 1 | GADAG | Sorghum, Bengalgram | 138 | 585.2 | Bengalgram |
| 2 | HAVERI | Maize, Oilseed, Sorghum and cotton | 158.3 | 599.9 | Sorghum |
| 3 | RAICHUR | Sorghum, Bengalgram | 131.3 | 608 | Sorghum, Bengalgram |
| 4 | KOPPAL | Sunflower, Sorghum, Maize, Bengalgram and cotton | 144.3 | 632.7 | Bengalgram and Sesamum |
| 5 | GULBARGA | Sorghum, Bengalgram and Sunflower | 114.9 | 613.3 | Sesame |
| 6 | DHARWAD | Cotton, Bengalgram and Sorghum | 145.7 | 585.2 | Bengalgram and Sorghum |
| 7 | BIDAR | Bengalgram, Sorghum and Sunflower | 118.9 | 616.5 | Sesame |
| 8 | BIJAPUR | Sorghum, Bengal gram and Sunflower | 121.8 | 592.5 | Sesame |
| 9 | BELGAUM | Maize and cotton | 134.6 | 585.2 | Sesame |
| 10 | BAGALKOTE | Sorghum and Bengalgram | 133.6 | 592.5 | Sorghum and Bengalgram |

* Water requirement of Bengal gram is 150mm, Sesamum 150 mm and Sorghum 500mm

Table:5

RAYALASEEMA

KHARIF

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL(mm) | PET(mm) | RECOMMENDED CROPS |
|------|------------|---|---------------|---------|------------------------------------|
| 1 | ANANTHAPUR | Groundt, Sunflower, Redgram, Maize and Cotton | 296.3 | 789.6 | Redgram, Black gram and Green gram |
| 2 | CHITTOOR | Groundnut and Redgram | 433.6 | 726 | Redgram and sunflower |
| 3 | KADAPA | Groundnut, Sunflower, Cotton and Redgram | 385.8 | 689.2 | Red gram and Bajra |
| 4 | KARNOOL | Groundnut, Sunflower, Sorghum, Redgram, | 449.8 | 738.8 | Sunflower and Redgram, |

Table:6

RAYALASEEMA

RABI

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL(mm) | PET(mm) | RECOMMENDED CROPS |
|------|------------|---|---------------|---------|----------------------------|
| 1 | ANANTHAPUR | Bengalgram and Sorghum | 185.9 | 675.2 | Bengal gram and Vegetables |
| 2 | CHITTOOR | Groundnut, Paddy, Redgram and sunflower | 421.2 | 635 | Redgram |
| 3 | KADAPA | Sunflower, Bengalgram, Cotton and | 266 | 641.4 | Bengalgram, Black gram, |

| | | | | | |
|---|---------|-----------------------|-------|-----|-----------------------------------|
| | | Sesamum | | | Sesame and Green gram |
| 4 | KARNOOL | Bengalgram and Chilli | 152.9 | 643 | Bengal gram and Chilli, Lablab |

*Water requirement of Lablab is 160mm

Table:7

TELANGANA

KHARIF

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL(mm) | PET(mm) | RECOMMENDED CROPS |
|------|-------------|--|---------------|---------|--------------------------------------|
| 1 | ADILABAD | Cotton, Soybean, Redgram and Sorghum | 931.8 | 576.1 | Cotton, Redgram and Soybean |
| 2 | HYDERABAD | Redgram, Sorghum and Maize | 602.2 | 426.3 | Redgram, Sorghum and Maize |
| 3 | KARIMNAGAR | Cotton, MaizeandGreengram | 755.6 | 539.1 | Cotton, MaizeandGreengram |
| 4 | KHAMMAM | Cotton, Maize, Greengram and Redgram | 900.1 | 576.8 | Cotton, Maize, and Redgram |
| 5 | MAHBUBNAGAR | Maize, Cstor, Groundnut, Redgram, Sorghum and Cotton | 485.1 | 668.4 | Castor, Redgram and Sorghum |
| 6 | MEDAK | Maize, Sorghum Cotton Greengram, Redgram and Blackgram | 613.3 | 581.5 | Maize, Sorghum and Cotton |
| 7 | NALGONDA | Cotton, Castor, Redgram and Groundnut | 506.6 | 586.8 | Castor and Redgram |
| 8 | NIZAMABAD | Maize and Redgram | 824.7 | 581.5 | Maize and Redgram |
| 9 | WARANGAL | Cotton, Maize, Paddy, Groundnut, Green gram | 819.1 | 601.5 | Cotton, Maize, Groundnut, Green gram |
| 10 | RANGAREDDY | Redgram, Sorghum, Maize and Cotton | 606.5 | 426.3 | Redgram, Sorghum, Maize and Cotton |

*Water requirement of Castor crop 500mm

Table:8 TELANGANA

RABI

| S.NO | DISTRICT | EXISTINGCROPS | RAIN FALL(mm) | PET(mm) | RECOMMENDED CROPS |
|------|-------------|------------------------------------|---------------|---------|-----------------------------|
| 1 | ADILABAD | Sorghum, Redgram and Bengal gram | 108.8 | 549.1 | Bengal gram |
| 2 | HYDERABAD | Sorghum and Bengal gram | 151.5 | 544.5 | Bengal gram and Green gram |
| 3 | KARIMNAGAR | Green gram | 115.5 | 544 | Bengal gram and Fodder crop |
| 4 | KHAMMAM | Maize, Green gram and Redgram | 151.5 | 580.1 | Bengalgram and Greengram |
| 5 | MAHBUBNAGAR | Maize and Sorghum | 128.6 | 297.5 | Bengalgram and Sesame |
| 6 | MEDAK | Sorghum, Safflower and Bengalgram | 100 | 549.2 | Fodder crop(cowpea) |
| 7 | NALGONDA | Groundnut | 149.7 | 567.8 | Bengal gram |
| 8 | NIZAMABAD | Maize | 126.25 | 549.2 | Bengal gram and Sesame |
| 9 | WARANGAL | Maize and Chilli | 124.2 | 591.9 | Bengal gram and Sesame |
| 10 | RANGAREDDY | Sorghum, Bengal gram and Groundnut | 93.6 | 544.5 | Fodder crop(cowpea) |

*Most drought tolerant, as per the following ranking. Soybean < black gram < moong bean < ground nut < maize < sorghum < pearl millet < lab lab bean < cowpea.