

Original Research Article

Incidence and Intensity of Early Blight in Potato under different dates of Planting

ABSTRACT

Aim: The study aims to progression of early blight of potato in relation to weather parameters such as, maximum temperature, minimum temperature, maximum relative humidity, minimum relative humidity and rainfall.

Place and Duration of Study: Epidemiological investigation was conducted during rabi season of 2021-22 at the research field of All India Coordinated Research Project (AICRP) on potato, OUAT, Bhubaneswar.

Methodology: The trial was laid out in Split Plot Design with different dates of planting (26 November and 16 December), as the main plots and four varieties (Kufri Pukhraj, Kufri Khyati, Kufri Surya, and Kufri Jyoti) as the sub plots.

Results: The initial infection of early blight occurred during 52 Standard Meteorological Week (SMW). The highest percentage of disease intensity and incidence occurred in variety Kufri Pukhraj followed by Kufri Khyati, Kufri Jyoti and Kufri Surya under 26 November planting.

Conclusion: Minimum temperatures, maximum and minimum relative humidity had significantly positive correlation with the incidence and intensity of early blight. Stepwise multiple linear regression equations revealed that, maximum temperature, maximum and minimum relative humidity and rainfall were responsible for early blight in potato.

Keywords: Early blight, Epidemiology, Potato, incidence, intensity

1. INTRODUCTION

Potato (*Solanum tuberosum* L.) is a major food crop grown worldwide. It is used in the production of starch, foods like potato chips and stock feed. (Ahmadizadeh and Felenji 2011). Potatoes are a great source of carbohydrates and are utilized in both for table consumption and in processed foods. Potato tubers contain around 80% of water. 20% dry matter. More than 75 % of the dry Starch, protein, fibres, and a negligible quantity of fatty acids make up matter (Prokop and Albert, 2008). Additionally, it is abundant in minerals like potassium, phosphorus, magnesium with B1, B3, and B6 vitamins (Camire et al., 2009). The potato also includes Vitamin C and several phenolic compounds act as strong antioxidants (Brown, 2005).

Potato is the world's 4th important food crop after wheat, rice and maize because of its great yield potential and high nutritive value. China is the world's largest producer and consumer of potato, produce 78.24 Mt in 4.21 m ha⁻¹ with a productivity of 18.55 t ha⁻¹ followed by India (51.30 Mt production in 2.16 m ha⁻¹ with 23.78 t ha⁻¹ productivity) and Russia (19.61 Mt in 1.18 m ha⁻¹ with a productivity of 16.65 t ha⁻¹) (According to the estimates of 2020, FAO 2021). In India most of the potato production takes place in Uttar Pradesh (15892 thousand tonnes production in 622.50 thousand ha⁻¹ area) followed by West Bengal (12600 thousand tonnes in 447 thousand ha⁻¹) and Bihar (9125.80 thousand tonnes in 330 thousand ha⁻¹ area) according to the estimates of 2021, NHB2021.

Currently Odisha produce 308.22 thousand tonnes in 25.91 thousand ha⁻¹ area (NHB 2021). Potato is one of the major constituents of the daily diet in Odisha. It is cultivated all districts of the state in winter season and in kharif season it is cultivated at Phulbani and Koraput districts.

Early blight symptoms, which are characterized by dark brown to black lesions with concentric rings and produce a "target spot effect," are first noticed on older and senescing leaves (Van der Waals, et al., 2001). The lesions are frequently encircled by a small chlorotic halo because of the pathogen's toxins, which progress into healthy epidermal cells. Defoliation may occur completely under extreme circumstances, particularly in places with high temperatures (24–29°C), high humidity levels, and semi-arid condition's locations where frequent and protracted dew occur (Rotem and Reichert, 1964).

Considering the significance of these illnesses as the primary biotic the current study has been limited by factors in the region's potato yield taken to analyses the impact of various meteorological parameters (maximum temperature, minimum temperature, maximum RH, minimum RH, and rainfall etc.) on the onset and progression of early potato blight.

2. MATERIALS AND METHODS

2.1 Experimental Site

The field experiment was conducted during Rabi season of 2021-2022 at the experimental plots of All India Coordinated Research Project on Potato, Odisha University of Agriculture and Technology, Bhubaneswar, located at 20° North latitude, 86° East longitude and at about 45 m above MSL consecutively for three years. The soil for the study was sandy loam in texture, acidic (pH 5.56) in reaction, low in organic carbon (0.51 %) and available N (218.4 kg ha⁻¹), medium in both available P (20.8 kg ha⁻¹) and K (96.1 kg ha⁻¹). The treatments were

a combination of two planting dates (D) and four varieties (V). The two dates of planting followed in the experiment were 26 November, 16 December and the varieties were Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti in the medium duration groups. The seed tubers of these varieties were kept in cold store till planting. Well sprouted foundation seed tubers were planted in 3 m x 2.4 m sized plots at 60 x 20 cm spacing in a Split Plot design with five replications. The present investigations on the development of early blight of potato in relation to weather parameters (maximum temperature, minimum temperature, maximum relative humidity, minimum relative humidity and rainfall), The weekly data of weather parameters during the cropping period (26 November to 04 March, 2021-22) were collected from the central observatory of the department of Agricultural meteorology, University of Agriculture & Technology University in Bhubaneswar, Odisha.

2.2 Climate

The general climatic condition of Bhubaneswar is under hot and humid. The maximum temperature during crop growth period in 2021-2022 was 31.2°C and minimum temperature was 10°C. The experimental site is in the east coastal plain of India. The mean annual rainfall is approximately 1408mm and annual maximum temperature 42.2°C and annual minimum temperature 11.1°C.

2.3 Weather conditions during crop growth season

The weekly mean of maximum temperature, minimum temperature, relative humidity (RH) and bright sunshine hour (BSH) along with total weekly rainfall, number of rainy days during the crop growth season (26 November to 04 March).

The weekly maximum temperature during the crop growth period ranged from 25.7°C to 33.1°C, with a weekly average of 28.01°C, whereas the weekly minimum temperature ranged from 10°C to 19.8°C, with a weekly average of 16.03°C.

The mean morning relative humidity during the crop growth varied from 86.4 to 95%, while the mean afternoon relative humidity varied from 26 to 84.3%. The mean bright sunshine hour received during the crop growth period varied from 1.3 to 9 hours, with a weekly average of 5.0 hours. The total rainfall received during the crop growth period was 142.8 mm.

The weekly mean of maximum temperature, minimum temperature, relative humidity (RH) and bright sunshine hour (BSH) along with total weekly rainfall, number of rainy days during the crop growth season (26 November to 04 March).




Fig.1 Weekly mean of maximum temperature, minimum temperature, relative humidity (RH) and bright sunshine hour (BSH)

2.4 Pathological observations

Disease incidence and intensity was calculated from the five tagged plants in each plot of the experimental field. Based on the data, percentage of disease incidence and intensity at weekly intervals, from the date of disease initiation till the crop attained maturity were recorded.

2.4.1 Disease incidence

Percentage disease incidence was worked out as per the following formula given by James (1974).

$$\text{Percent disease incidence} = \frac{\text{No. of diseased leaves}}{\text{Total No. of leaves examined}} \times 100 \quad (\text{Eq. 1})$$

2.4.2 Disease intensity

According to the Mayer and Datar standard area diagram, Percentage intensity was computed (1986).

The Percentage of disease intensity was calculated by using the following formula: -

$$\text{Percentage of Disease Intensity} = \frac{\text{Sum of individual ratings}}{\text{No. of plants observed} \times \text{Maximum disease rating}} \times 100 \quad (\text{Eq.2})$$

Table 1 Disease intensity scale for early blight of potato

Scale	Description of the symptoms	Reaction
0	Leaves free from infection	Highly resistant
1	Small irregular spots covering <5% leaf area	Resistant
2	Small irregular brown spots with concentric rings covering 5.1-10% leaf area	Moderately resistant
3	Lesions enlarging, irregular brown with concentric rings covering 10.1-25% leaf area	Moderately susceptible
4	Lesions coalesce to form irregular and appears as a typical blight symptom covering 25.1-50% leaf area	Susceptible
5	Lesions coalesce to form irregular and appears as a typical blight symptom covering >50% leaf area	Highly susceptible

2.5 METHODS

2.5.1 Correlation analysis

Potato early blight Correlation analysis of the disease incidence and intensity (dependent variable) and (independent variable) with weather parameter i.e., maximum and minimum temperature⁰C, maximum relative humidity (present) before a week from the date of minimum relative humidity (present) and rainfall (mm)) surveillance for early potato blight in 20121–22.

Varieties (Kufri Pukhraj, Kufri Khyati, Kufri Surya, Kufri Jyoti) was conducted with an aim to measure the degree of association among the different predictor and response variables in causing the onset and the progress of the disease incidence and intensity of both the dates of planting. The Pearson correlation coefficients (r) were studied to find out effect of single as well as combination of different weather factors on the disease progression of early blight of potato.

2.5.2 Multiple regression

To predict the disease incidence and intensity early blight of potato in four varieties, different models were generated with the disease incidence and intensity of both the dates and weather factors by using multiple linear regression.

$$Y = a + b_1X_1 + b_2X_2 + b_3X_3 + b_4X_4 + b_5X_5$$

Where, Y is the predicted disease incidence and intensity of disease “ a ” is the intercept, “ b_1 ” to “ b_5 ” are the partial regression coefficients, X_1 is the maximum temperature ($^{\circ}\text{C}$), X_2 is the minimum temperature ($^{\circ}\text{C}$), X_3 is the maximum relative humidity (%), X_4 is the minimum relative humidity (%) and X_5 is the total rainfall (mm).

3. RESULT AND DISCUSSION

3.1 Incidence of early blight

3.1.1 Effect of Weather parameters on the incidence of early blight of potato in under different dates of planting

Incidence of early blight of potato varied from 2.2 to 23% in all the selected varieties of potato. The disease was first observed at 52 Standard Meteorological Week (30 days after planting) in Kufri Pukhraj (3.00%), Kufri Khyati (2.7%) Kufri Surya (2.2%) and Kufri Jyoti (2.5%). The weather parameters during the preceding week were maximum and minimum temperature of 27.3°C and 15.4°C , maximum and minimum relative humidity 93.1% and 79.1% and rainfall of 0.8mm. Singh, et al. (2017) reported similar observation that maximum temperature and relative humidity had significant effect on early blight incidence. The disease incidence progressed gradually and showed progression from 4.80 to 17.5%, 3.25 to 10.25%, 3.2 to 7.8% and 5.5 to 15.6% in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, respectively during 1st to 3rd SMW (37 to 53 DAP). During the period, maximum and minimum temperatures were 26.7°C and 12.8°C , maximum and minimum relative humidity 94% and 26% and rainfall 16.5 mm. At the maturity, (60 DAP) in 4th SMW, maximum disease incidence was recorded in Kufri Pukhraj (23%) followed by Kufri Jyoti (15.6%), Kufri Khyati (13.50%) and Kufri Surya (9.8%), when maximum temperature was 27.5°C , minimum temperature 16.9°C , maximum relative humidity 95% and minimum relative humidity 41%. In second date of planting (16 December) incidence of early blight of potato varied from 1.1 to 12.40% in all the selected varieties of potato (Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti). The disease was first observed at 2nd Standard Meteorological Week (30 days after planting) in Kufri Pukhraj (1.25%), Kufri Khyati (1.50%) Kufri Surya (1.4%) and Kufri Jyoti (1.1%). The weather parameters during the preceding week were maximum and minimum temperature of 26.7°C and 15.7°C , maximum and minimum relative humidity 94% and 39% and rainfall of 16.5mm. The findings of Alternating low and high humidity conditions have also been shown to favour disease development (Van der-Walls et al., 2001). The disease varies from 2.25 to 10.20%, 2.50 to 7.40%, 1.8 to 6.8% and 3.2 to 9.6 in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, respectively during 3rd to 5th SMW (37 to 53 DAP). During the period, maximum and minimum temperatures were 28.2°C

and 14.6°C, maximum and minimum relative humidity 95% and 26% and rainfall 12.6mm. At the maturity stage, (60 DAP) in 6th SMW, maximum disease incidence was recorded in Kufri Pukhraj (12.40%) followed by Kufri Jyoti (11%), Kufri Khyati (8.25%) and Kufri Surya (7.5%), when maximum temperature was 28.7°C, minimum temperature 15.8°C, maximum relative humidity 95% and minimum relative humidity 66 %.The highest disease incidence (23.0 and 12.40%) was reported in Kufri Pukhraj followed by 15.6 and 11.0% in Kufri Jyoti, whereas minimum disease incidence was observed in Kufri Surya and Kufri Khayati (9.8 and 7.5%), in first and second date of planting, respectively. The variation in disease incidence might be due to the response of different varieties against the disease and planting under different environmental conditions. Present study showed that maximum incidence of early blight of potato was recorded in all selected varieties under first date of planting as compared to second date of planting which might be due to the older and senescence leaves, more susceptible for development of early blight disease. Similar results were also found by Vander-Walls et al., (2003). Changes in weather variables and amount of initial inoculum of *A. solani* may be responsible for varying disease intensities at different locations.

Table 2 Effect of weather parameters on the incidence of early blight of potato under different dates of planting

SM W	Disease Incidence %								Weather parameters				
	1 st Date of planting (26.11.2021)				2 nd Date of planting (16.12.2021)				Max Tem .	Min. Tem p.	RH (%) 7 hrs	RH (%) 14 hrs	Rain fall (mm)
	K. Pukhra j	K. Khyat i	K. Sury a	K. Jyoti	K. Pukhr aj	K.Khy ati	K. Sury a	K. Jyoti					
52	3.00	2.70	2.2	2.5	0	0	0	0	27.3	15.4	93.1	79.1	0.8
1	4.80	3.25	3.2	5.5	0	0	0	0	26.4	12.8	92	77	0.0
2	9.20	6.75	5.4	9.2	1.25	1.50	1.4	1.1	26.7	15.7	94	39	16.5
3	17.50	10.25	7.8	11.5	2.25	2.50	1.8	3.2	25.7	14.6	95	26	0.0
4	23.00	13.50	9.8	15.6	3.80	5.20	3.8	5.8	27.5	16.9	95	41	12.6
5	-	-	-	-	10.20	7.40	6.8	9.6	28.2	14.7	95	68	0.0
6	-	-	-	-	12.40	8.25	7.5	11	28.7	15.8	95	66	0.0

3.1.2 Correlation of weather parameters on the incidence of early blight of potato in under different dates of planting

Maximum temperature was non-significant with the disease incidence of early blight of potato, in all the varieties under 26 November planting (Table 3). Maximum temperature was positively and significant correlation with the incidence of early blight of potato, having ($r = 0.885, 0.91, 0.925, \text{ and } 0.881$) in all the selected varieties under on 16 December planting, which showed that increase in temperature leads to increase in disease incidence. The results were similar to those of Tiwari et al. (2004) who reported that maximum temperature and relative humidity had significant effect on early blight, indicating that higher *Alternaria* spore concentration was recorded at higher temperature. Morning relative humidity had also significant positive correlation with disease incidence ($r = 0.899, \text{ and } 0.881$) in, Kufri Khyati, and Kufri Surya potato varieties on 26th November planting. And Morning relative humidity was non-significant with the disease incidence of early blight of potato in Kufri Pukhraj and Kufri Jyoti on 26th November planting. Under 16th December planting night relative humidity had significant ($r = 0.929, 0.887, 0.938 \text{ and } 0.881$) with all the selected varieties which showed that night relative humidity less than 80% exerted negative impact on disease incidence. Temperature and relative humidity played a major role in the dispersion of *Alternaria* spores (Burch and Levetin, 2002; Stennet and Beggs, 2004). Sabariego et al. (2000) reported negative correlation of early blight of potato with humidity and rainfall. SukruthaHerle and Kamanna (2014) also reported that early blight of potato was negatively

correlated with minimum temperature, and relative humidity (maximum and minimum), while positively correlated with maximum temperature. was also reported by Escuredo et al. (2011), indicating that higher *Alternaria* spore concentration were recorded at higher temperature.




Fig.2. Disease Incidence% (D1)




Fig.3. Disease Incidence% (D2)

Table.3 Correlation of weather parameters on the incidence of early blight of potato in under different dates of planting

Weather parameters	1 st Date of planting (26.11.2021)				2 nd Date of planting (16.12.2021)			
	Varieties				Varieties			
	Kufri Pukhraj	Kufri Khyati	Kufri Surya	Kufri Jyoti	Kufri Pukhraj	Kufri Khyati	Kufri Surya	Kufri Jyoti
T _{max.} (°C)	0.004 ^{NS}	0.035 ^{NS}	-0.018 ^{NS}	-0.004 ^{NS}	0.885*	0.911*	0.925*	0.881*
T _{Min.} (°C)	0.563 ^{NS}	0.617 ^{NS}	0.566 ^{NS}	0.543 ^{NS}	-0.111 ^{NS}	0.080 ^{NS}	-0.006 ^{NS}	-0.009 ^{NS}
RH _{7hrs} (%)	0.877 ^{NS}	0.899*	0.881*	0.829 ^{NS}	0.529 ^{NS}	0.657 ^{NS}	0.571 ^{NS}	0.675 ^{NS}
RH _{14hrs} (%)	-0.791 ^{NS}	-0.818 ^{NS}	-0.832 ^{NS}	-0.820 ^{NS}	0.929*	0.887*	0.938*	0.881*
Rainfall (mm)	-0.369 ^{NS}	0.448 ^{NS}	0.444 ^{NS}	0.526 ^{NS}	-0.652 ^{NS}	-0.569 ^{NS}	-0.585 ^{NS}	-0.647 ^{NS}

T_{Max.} = Temperature maximum; T_{Min.} = Temperature minimum; RH_{7hrs} = Maximum Relative humidity; RH_{14hrs} = Minimum Relative humidity

***Significant at 5% level **Significant at 1% level**

The multiple linear regression model (Table 4) indicated among different independent variables that morning relative humidity (X3) was responsible 76.38%, 80.89%, 77.62%, and 68.72% variations for early blight of potato in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, respectively under 26 November date of planting. And the multiple linear regression indicates the relationship between disease incidence and weather parameters of early blight of potato in under 16 December planting where maximum temperature (X1), maximum relative humidity (X3), minimum humidity (X4), rainfall (X5) had contributed 86.27%, 99.98%, 98.53% and 97.82% in the incidence of early blight of potato Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti respectively. Similar results were also found by Behera (2009) the study on regression coefficient in relation to disease incidence indicated that maximum temperature had a significant impact on early blight of potato. Among all the weather parameters the contribution of rainfall was maximum (39.1 %) on the disease incidence.

Table 4 Multiple linear regression weather parameters on the incidence of early blight of potato in under different dates of planting

Date of sowing	Varieties	Stepwise Regression Equation	R ²	P value
1 st Date of planting (26.11.2021)	Kufri Pukhraj	$Y = -861.59 + 9.34868 * X_3$	0.7638	0.0527
	Kufri Khyati	$Y = -294.45 + 3.21616 * X_3$	0.8089	0.0377
	Kufri Surya	$Y = -196.58 + 2.15584 * X_3$	0.7762	0.0484
	Kufri Jyoti	$Y = -299.13 + 3.28279 * X_3$	0.6872	0.0827
2 nd Date of planting (16.12.2021)	Kufri Pukhraj	$Y = -6.2119 + 0.254 * X_4$	0.8627	0.0226
	Kufri Khyati	$Y = -228.78 + 1.93065 * X_1 + 1.91231 * X_3 + 0.624 * X_5$	0.9998	0.0178
	Kufri Surya	$Y = -202.86 + 2.11935 * X_3 + 0.12939 * X_4$	0.9853	0.0147
	Kufri Jyoti	$Y = -416.18 + 4.36792 * X_3 + 0.17174 * X_4$	0.9782	0.0218

3.2 Intensity of early blight

3.2.1 Effect of Weather parameters on the intensity of early blight of potato in under different dates of planting

In early blight of potato, intensity of early blight of potato varied from 2.40 to 21.23% in all the selected varieties of potato (Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti) in under 26th November planting (Table 5). The result showed that disease initiation occurred under field condition on 52SWM, in Kufri Pukhraj (2.81%), Kufri Khyati (2.40%) Kufri Surya (2.50%) and Kufri Jyoti (3.50%). The weather parameters during the preceding week were maximum and minimum temperature of 27.3^oC and 15.4^oC, maximum and minimum relative humidity 93.1% and 79.1% and rainfall of 0.8mm. Abuley (2015) reported that temperature from 10 to 35^oC were more favourable for early blight epidemic. The disease intensity progressed gradually and showed progression varies from 4.31 to 15.5%, 3.12 to 9.82%, 3.4 to 8.0% and 6.5 to 12.5% in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, respectively during 1st to 3rd SMW (37 to 53 DAP). During the period, maximum and minimum temperatures were 26.7^oC and 12.8^oC, maximum and minimum relative humidity 94% and 26% and rainfall 16.5mm. At the maturity stage, (60 DAP) in 4th SMW, maximum disease intensity was recorded in Kufri Pukhraj (21.23%) followed by Kufri Jyoti (16.8%), Kufri Khyati (12.84%) and Kufri Surya (10.2%), when maximum temperature was

27.5⁰C minimum temperature 16.9 ⁰C, maximum relative humidity 95% and minimum relative humidity 4%. In second date of planting (16 December) disease intensity was first observed at 2nd Standard Meteorological Week (30 days after planting) in Kufri Pukhraj (1.25%), Kufri Khyati (1.30%) Kufri Surya (1.5%) and Kufri Jyoti (1.3). The weather parameters during the preceding week were maximum and minimum temperature of 26.7⁰C and 15.7⁰C, maximum and minimum relative humidity 94% and 39% and rainfall of 16.5mm. These findings agree with the findings of Mehboob et al. (2013) who observed that maximum disease intensity occurred at maximum temperature of 17 to 20⁰C and minimum temperature of 6 to 9⁰C. Rotem (2004) also reported that availability of water in the form of relative humidity, rainfall or dew increased conidial germination of *A. solani*. The disease intensity progressed gradually and showed progression from 2.14 to 10.7%, 2.30 to 6.84%, 2.0 to 7.0% and 3.6 to 10.2% in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, respectively during 3rd to 5th SMW (37 to 53 DAP). During the period, maximum and minimum temperatures were 28.2⁰C and 14.6⁰C, maximum and minimum relative humidity 95% and 26% and rainfall 12.6mm. At the maturity stage, (60 DAP) in 6th SMW, maximum disease intensity was recorded in Kufri Pukhraj (11.80%) followed by Kufri Jyoti (11.2%), Kufri Surya (8.5%), and Kufri Khyati (7.20%) when maximum temperature was 28.7⁰C, minimum temperature 15.8⁰C, maximum relative humidity 95% and minimum relative humidity 66%. The highest disease intensity (21.23 and 11.80%) was reported in Kufri Pukhraj followed by 16.8 and 11.2% in Kufri Jyoti, whereas minimum disease intensity was observed in Kufri Surya and Kufri Khayati (10.20 and 7.20%), in first and second date of planting, respectively. The variation in disease intensity might be due to the response of different varieties against the disease and planting under different environmental conditions. Our results show that maximum intensity of early blight of potato was recorded in all selected varieties under first date of planting as compared to second date of planting which may be due to that older and senescence leaves are more susceptible for development of early blight disease. Similar results were also found by R. Chaerani and Voorrips (2006) reported that initially progress of early blight was slow but accelerated as plants attain maturity showing sigmoid disease curve.




Fig.4 . Disease Intensity% (D1)




Fig.5. Disease Intensity% (D2)

Table 5 Effect of weather parameters on the intensity of early blight of potato in under different dates of planting

S M W	Disease intensity %								Weather parameters				
	1 st Date of planting (26.11.2021)				2 nd Date of planting (16.12.2021)				Max Temp.	Min. Temp.	RH (%) 7 hrs	RH (%) 14 hrs	Rain fall (mm)
	Kufri Pukhraj	Kufri Khyati	Kufri Surya	Kufri Jyoti	Kufri Pukhra j	Kufri Khyati	Kufri Surya	Kufri Jyoti					
52	2.81	2.40	2.5	3.5	0	0	0	0	27.3	15.4	93.1	79.1	0.8
1	4.31	3.12	3.4	6.5	0	0	0	0	26.4	12.8	92	77	0.0
2	9.57	6.24	5.8	9.5	1.25	1.30	1.5	1.3	26.7	15.7	94	39	16.5
3	15.50	9.82	8.0	12.5	2.14	2.30	2.0	3.6	25.7	14.6	95	26	0.0
4	21.23	12.84	10.2	16.8	3.56	5.06	4.0	6.5	27.5	16.9	95	41	12.6
5	-	-	-	-	10.70	6.84	7.0	10.2	28.2	14.7	95	68	0.0
6	-	-	-	-	11.80	7.20	8.5	11.2	28.7	15.8	95	66	0.0

3.2.2 Correlation and multiple linear regressions on the intensity of early blight of potato in under different dates of planting

Maximum temperature was non- significant with the disease intensity of early blight of potato, in all the selected varieties under 26November planting (Table 6). Maximum temperature was positively and significant correlated with the intensity of early blight of potato, having $r = 0.879, 0.901$ and 0.921 , in Kufri Pukhraj, Kufri Khyati, and Kufri Surya, respectively. On 16 December planting, which showed that increase in temperature leads to increase in disease intensity. Significantly positive correlation between temperature and the Alternaria spore concentration significant at 5% level also reported by Escuredo et al. (2019), indicating that higher Alternaria spore concentration was recorded at higher temperature.

Morning relative humidity had also significant positive correlation with disease intensity ($r = 0.884, 0.894, \text{ and } 0.884$) in Kufri Pukhraj, Kufri Khyati, and Kufri Surya potato varieties on 26 November planting. And Morning relative humidity was non-significant with the disease intensity of early blight of potato in all the selected varieties under on 16 December planting. Under 16 December planting night relative humidity had $r = 0.942, \text{ and } 0.924$ in Kufri Pukhraj, and Kufri Surya which showed that night relative humidity less than 80% exerted negative impact on disease intensity. Temperature and relative humidity played a major role in the dispersion of *Alternaria* spores (Burch and Levetin, 2002; Stennet and Beggs, 2004). Sabariego et al. (2000) reported negative correlation of early blight of potato with humidity and rainfall. SukruthaHerle and Kamanna (2014) also reported that early blight of potato was negatively correlated with minimum temperature, and relative humidity (maximum and minimum), while positively correlated with maximum temperature. was also reported by Escuredo et al. (2019), indicating that higher *Alternaria* spore concentration were recorded at higher temperature.

Table 6 Correlation of weather parameters on the intensity of early blight of potato in under different dates of planting

Weather parameters	1 st Date of planting (26.11.2021)				2 nd Date of planting (16.12.2021)			
	Varieties				Varieties			
	Kufri Pukhraj	Kufri Khyati	Kufri Surya	Kufri Jyoti	Kufri Pukhraj	Kufri Khyati	Kufri Surya	Kufri Jyoti
T _{max.} (°C)	-0.036 ^{NS}	0.021 ^{NS}	0.005 ^{NS}	0.008 ^{NS}	0.879*	0.901*	0.921*	0.875 ^{NS}
T _{Min.} (°C)	0.601 ^{NS}	0.600 ^{NS}	0.587 ^{NS}	0.534 ^{NS}	-0.154 ^{NS}	0.106 ^{NS}	0.011 ^{NS}	0.001 ^{NS}
RH _{7hrs} (%)	0.884*	0.894*	0.884*	0.823 ^{NS}	0.521 ^{NS}	0.683 ^{NS}	0.564 ^{NS}	0.697 ^{NS}
RH _{14hrs} (%)	-0.805 ^{NS}	-0.816 ^{NS}	-0.830 ^{NS}	-0.795 ^{NS}	0.942*	0.869 ^{NS}	0.924*	0.872 ^{NS}
Rainfall (mm)	0.441 ^{NS}	0.431 ^{NS}	0.467 ^{NS}	0.483 ^{NS}	-0.658 ^{NS}	-0.547 ^{NS}	-0.591 ^{NS}	-0.641 ^{NS}

T_{Max.} = Temperature maximum; T_{Min.} = Temperature minimum; RH_{7hrs} = Maximum Relative humidity; RH_{14hrs} = Minimum Relative humidity

*Significant at 5% level **Significant at 1% level

The multiple linear regression model indicated among different independent variables, morning relative humidity (X3) was responsible that 78.17%, 79.89%, 78.07%, and 67.66% variations for early blight of potato in Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti, under 26 November date of planting, respectively. And the multiple linear regression indicated the relationship between disease intensity and weather parameters of early blight of potato in under 16 December planting that the maximum temperature (X1), maximum relative humidity (X3), minimum humidity (X4), rainfall (X5) had contributed 98.21%, 99.28%, 85.36% and 99.94 % in the intensity of early blight of potato Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti respectively. Similar findings by Gupta et al., 2020 early blight of tomato appeared in the 12th SMW and had steep increase throughout the cropping period. Maximum and minimum temperatures had significantly positive correlation, whereas maximum and minimum relative humidity along with rainfall had negative correlation with the PDI of early blight of tomato. Stepwise regression model explained that 83 per cent variation in the PDI of the disease was due to maximum temperature.

Table 7 Multiple linear regression weather parameters on the intensity of early blight of potato in under different dates of planting

Date of sowing	Varieties Disease intensity	Stepwise Regression Equation	R ²	P value
1 st Date of sowing (26.11.2021)	Kufri Pukhraj	$Y = -816.07 + 8.8549 * X_3$	0.7817	0.0465
	Kufri Khyati	$Y = -281.8 + 3.07696 * X_3$	0.7989	0.0409
	Kufri Surya	$Y = -199.1 + 2.18592 * X_3$	0.7807	0.0468
	Kufri Jyoti	$Y = -299.78 + 3.29934 * X_3$	0.6766	0.0179
2 nd Date of sowing (16.12.2021)	Kufri Pukhraj	$Y = -3.4761 + 0.22007 * X_4 - 0.2057 * X_5$	0.9821	0.0072
	Kufri Khyati	$Y = -293.27 + 1.69386 * X_1 + 2.65257 * X_3$	0.9928	0.0193
	Kufri Surya	$Y = -2.844 + 0.15508 * X_4$	0.8536	0.0249
	Kufri Jyoti	$Y = -309.82 + 2.56774 * X_1 + 2.60484 * X_3 - 0.1387 * X_5$	0.9994	0.0307

4. CONCLUSION

The result of correlation analysis in Kufri Khyati, and Kufri Surya that maximum relative humidity had positive and highly significant correlation with disease under 26th November planting. Kufri Pukhraj, Kufri Khyati, Kufri Surya and Kufri Jyoti maximum temperature had positive and highly significant correlation with disease incidence. Whereas minimum relative humidity significantly positive correlation with the disease incidence, in all the varieties potato under 16th December planting. Stepwise multiple regression analysis among different independent variables that maximum relative humidity (X3) was highly significant in Kufri Pukhraj, Kufri Khyati, Kufri Surya, and Kufri Jyoti under 26th November planting. Under 16th December planting Minimum humidity(X4) was highly significant in Kufri Pukhraj followed by Kufri Khyati where maximum temperature (X1), maximum relative humidity (X3) and rainfall (X5) was highly significant and in Kufri Surya, and Kufri Jyoti maximum relative humidity (X3) and minimum humidity(X4) was highly significant.

The relationship between the intensity of early blight of potato and weather parameters produce result that Kufri Pukhraj, Kufri Khyati, and Kufri Surya varieties of potatoes disease intensity positive and highly significant with maximum relative humidity. In all the four potato varieties under 26November plantings disease intensity of Kufri Pukhraj, Kufri Khyati, and Kufri Surya had positive and highly significant with maximum temperature. Whereas Kufri Pukhraj, and Kufri Surya disease intensity was significantly positive correlation with minimum relative humidity under 16 December planting. Stepwise multiple regression analysis among different independent variables, maximum relative humidity (X3) was highly significant in Kufri Pukhraj, Kufri Khyati, Kufri Surya, and Kufri Jyoti varieties under on 26 November planting.

Minimum humidity(X4) and rainfall (X5) was highly significant in Kufri Pukhraj, maximum temperature (X1), and maximum relative humidity (X3) was highly significant in KufriKhyati, minimum relative humidity(X4) was highly significant in Kufri Surya, maximum temperature (X1), maximum relative humidity (X3), and rainfall (X5) was highly significant Kufri Jyoti under 16th December planting.

5. REFERENCES

1. Abuley IK. 2015. Decision support system in the control of potato early blight (*Alternaria solani* and *Alternaria alternata*). M.Sc. Agrobiolgy (Plant Nutrition and Health). Aarhus University.

2. Ahmadizadeh M, and Felenji H. 2011. Evaluating diversity among potato cultivars using agro-morphological and yield components in fall cultivation of Jiroft area. *American-Eurasian Journal of Agricultural and Environmental Science*, **11**(5): 655–66.
3. Behera B, Senapati S, Pattanaik KK, and Biswal G. (2009). Prediction of potato early blight in coastal tract of Orissa. *Journal of Plant Protection and Environment*, 6(1), 79-85.
4. Brown CR. 2005. Antioxidants in potato. *Annual Journal of Potato Research*, **82**:163-172.
5. Burch M. and Levetin E. 2002. Effects of meteorological conditions on spore plumes. *International Journal of Biometeorology*, **46**: 107-117.
6. Camire M E, Kubow S, and Donnelly DJ. 2009. Potatoes and human health. *Critical Reviews in Food Science and Nutrition*, **49**(10): 823-840.
7. Chaerani, R. and Voorrips, R. E. 2006. Tomato early blight (*Alternaria solani*): the pathogen, genetics and breeding for resistance. *Journal of General Plant Pathology*, **72**: 335–347.
8. Escuredo O, Seijo MC, Fernández-González M, and Iglesias I. (2011). Effects of meteorological factors on the levels of *Alternaria* spores on a potato crop. *International Journal of Biometeorology*, **55**(2), 243-252.
9. Gupta V, Razdan VK, Sharma S, and Fatima K. 2020. Progress and severity of early blight of tomato in relation to weather variables in Jammu province. *Journal of Agrometeorology*, **22**(2): 198-202.
10. James, W.. (1974). RESUME: Evaluation des pertes de récoltes dues au mildiou de la pomme de terre *Phytophthora infestans* (Mont.) de By. *Eppo Bulletin*. 4. 338-338. 10.1111/j.1365-2338.1974.tb02376.x.
11. Mayer C.D and Datar V.V. (1986) *Phytopathometry Technical Bulletin-1 1986 Marathwada Agriculture University Parbhani.*, p94
12. Mehboob S, Rehman A, Khan, MA, and Idrees M. 2013. Role of epidemiological and biochemical factors against early blight of potato. *Journal of Plant Pathology*, **2**(01): 8-13.
13. Prokop S, and Albert J. 2008. International year of the potato: Potatoes, nutrition, and diet. [online] Available from: <http://www.potato2008.org>
14. Rotem J, and Reichert L. 1964. Dew - A principal moisture factor enabling early blight epidemics in a semi-arid region of Israel. *Plant Diseases*, **48**: 211-215.
15. Rotem J. 2004. The genus *Alternaria*: biology, epidemiology, and pathogenicity. *American Phytopathological Society Press. St. Paul, MN, USA*.

16. Sabariego S, Diaz de la Guardia C, and Alba F. 2000. The effect of meteorological factors on the daily Variation of airborne fungal spores in Granada. *Int. J. Biometrol.* **44**: 1-5.
17. Singh SP, Bhatnagar A, Dua VK, Sharma SK, and Sadawarti MJ. (2017). Effect of planting windows on production of KufriKhyati: An early bulking potato cultivar for Central India. *International Journal of Chemical Studies*, **5**, 1798-1803.
18. Stennett, P. J. and Beggs, P. J. 2004. *Alternaria* spores in the atmosphere of Sydney, Australia, and relationship with meteorological factors. *International Journal of Biometeorology*, **49**: 98-105.
19. SukruthaHerle, G. and Kamanna, B. C. 2014. Effect of weather factors on early blight disease development in potato. *Plant Archives*, **14**(2): 1011-1014.
20. Tiwari RKS, Rajput ML, Singh A. Effect of sowing dates and spray schedule of mancozeb on early blight *Alternariasolani* [(Ell. & Mart.) Jones & Grout] of potato. *Indian Journal of Plant Protection* 2004; **32**(2):61-64.
21. Van der Waals, JE, Korsten L, and Aveling TAS. (2001). A review of early blight of potato. *African Plant Protection*, **7**(2), 91-102.
22. Van der Waals, JE, Korsten L, Aveling TAS, and Denner FDN. (2003). Influence of environmental factors on field concentrations of *Alternariasolani* conidia above a South African potato crop. *Phytoparasitica*, **31**(4), 353-364.