

Evaluation of recombinant inbred lines developed from the cross Rasi × Improved Samba Mahsuri for drought tolerance

Abstract:

Drought has been one of the most important limiting factors for rice productivity, particularly in the rainfed ecosystem. It is important to understand the genetic basis of drought tolerance in donor lines and develop drought tolerant rice cultivars, based on this information, we earlier identified the rice line, Rasi as tolerant to drought and developed a recombinant inbred line (RIL) mapping population consisting of 209 lines. These lines along with their parents were grown under well-watered and drought stress conditions in a two-year experiment (wet season 2020 and 2021) with drought stress imposed during reproductive stage. The study revealed high genetic variability for 12 key agro morphological traits associated with drought tolerance among the RILs. Of the 209 RILs, 59 showed superior performance over the checks and even the tolerant parent, Rasi under severe drought condition. Two RILs, RIL-33 and RIL-58 showed exceptional drought tolerance along with greater plot yield and grain weight under drought condition and they possessed medium slender grain type. These two lines can be a new novel genetic resource for drought tolerance in breeding programmes.

Keywords: Drought tolerance, Rasi, Improved Samba Mahsuri, Recombinant inbred lines

Introduction:

Rice (*Oryza sativa* L.) is the world's second most important cereal crop following corn. Traditionally, countries in Asia have the largest share in world rice production. Global consumption of rice has seen a slight increase over the last several years. In, 2021-22 crop year, about 509.87 million metric tons of rice was consumed worldwide. Rice is an important source of calories along with few proteins and essential nutrients, making it a very useful food for more than half of the world's population (Kilimo, 2018; Schenker, 2012).

Drought stress is one of the major abiotic constraints which causes remarkable losses in rice crop yield. The intensity of drought is increasing with changing environmental conditions and water scarcity. Rice production is heavily affected by drought which results in reduced germination, plant height, biomass, numbers of tillers, chlorophyll content, leaf number and size at various growth stages (Pandey and Shukla, 2015). Drought not only reduces rice yield but also affects the potential beneficial effects of improved crop management practices such as fertilizer application, pest and disease management. Plant responses to drought stress are very complex as the stress itself involves various climatic, soil and agronomic factors. Rice plants have a variety of strategies to combat drought; however tolerance to drought enables them to yield more, even in conditions of low water availability. Drought tolerance of rice is a complex trait and involves complex morpho-physiological mechanisms (Li and Xu, 2007). As the changing environment has a strong effect on drought tolerance response, different physio and morphological traits and the use of molecular markers helps to make a strategy for study of drought tolerance in rice under particular conditions. Thus, development of drought tolerant varieties is the most needed approach to tackle the problems imposed by a rapidly changing climate.

Selection of rice lines with improved grain yield under drought can be obtained by direct or indirect selection of traits that contribute for drought tolerance under water deficit condition (Dhawan et al., 2021; Sakran et al., 2022; Waheed et al., 2021). Traditional breeding methods for developing drought-tolerant rice varieties have become less effective due to low heritability and genotype selection being hampered by environmental and genetic interactions. This can be addressed by adopting cutting-edge high throughput genotyping and phenotyping methods (Sahebi et al., 2018).

In the current study, a rice cultivar, Rasi which was earlier identified and released as a drought tolerant rice variety was used to develop a recombinant inbred line (RIL) mapping population which was further screened under well-watered and drought stress conditions for the identification of RILs that are highly tolerant to drought conditions with desirable characters as the recurrent parent for their potential use in future rice breeding programs for drought tolerance.

Materials and methods:

Plant material:

Rasi is a drought-tolerant variety developed and released in the year 1977 by the ICAR-Indian Institute of Rice Research (ICAR-IIRR), Hyderabad, India. It has resistance to leaf blast, brown spot and tolerance to low soil phosphorus (V R Babu et al., 2016; Anila et al., 2022) and it can also grow well in aerobic conditions, saving 40-50% of the water consumption. Improved Samba Mahsuri (ISM) was developed through marker-assisted pyramiding of three bacterial blight resistance genes, *Xa21*, *xa13*, and *xa5* in the genetic background of the elite mega-variety of rice, Samba Mahsuri, and has broad spectrum resistance against bacterial blight disease (Sundaram et al., 2008). For drought tolerance screening under field conditions, the varieties, Sahbagidhan, DRR Dhan-42 and Vandana were used as positive checks, while Gangavathi Sona and BPT5204 were used as negative checks. When screened for drought at ICAR- IIRR, Hyderabad, ISM was found to be very sensitive while Rasi showed excellent tolerance to drought.

Experimental Site:

The experiments were conducted in upland irrigated conditions of ICAR-National Rice Research Institute-Central Rainfed Upland Rice Research Station (ICAR-NRRI-CRURRS), Hazaribagh, Jharkhand, India (23°56'34" N and 85°21'46" E) during Kharif 2020 (i.e. wet season 2020) and Kharif 2021 (i.e. wet season 2021). The RIL mapping population was subjected to drought stress and non-stress (control) conditions. Drought stress was artificially imposed by draining out water from the field during the reproductive stage which was referred to as stress trials.

Screening of the RIL population for Drought tolerance:

A cross was made between Rasi and Improved Samba Mashuri (ISM) and a RIL population was developed using a single seed descent method (Anila et al., 2022) These RILs (n=209) were used for phenotypic evaluation of drought tolerance at ICAR-NRRI-CRURRS, Hazaribagh. Wet bed nursery was being raised and 21 days old seedlings of the mapping population were transplanted into the drought screening plot along the parents. The

experiment was conducted in Alpha lattice design consisting of two replications with spacing of 15 × 20 cm (in three rows, constituting of 10 hills per row). The RILs were screened for reproductive stage drought tolerance by draining out water one month after transplanting (52 DAS), and the drain was kept open, till maturity. The crop was strictly maintained as rain fed and was not irrigated even once and the soil moisture was recorded constantly. The various agro-morphological parameters were recorded for the RILs along with the donor and recurrent parents. Days to 50% flowering, Plant Height (cm), Grain yield per plot (g/plot), Grain yield per hectare (Kg/ha), Harvest index (HI), Biomass (g)/0.5 m, Panicle length (cm), No. of Panicles per 0.5 m sample, Grain weight (g) of the sample for harvest index harvested from 0.5-meter length, No. of filled grains per panicle (average of 5 panicles), No. of unfilled grains per panicle (average of 5 panicles) and Sterility percentage (%). Three plants were selected randomly which represent the entire line and mean data was subjected to statistical analysis using open-source statistical software 'R' version 3.6.3 (R Core Team, 2016).

Results and Discussion:

Rice is the world's most important staple food crop, supplying more than half of the world's population (Narciso and Hossain, 2002). Drought is the single most significant constraint to rice productivity in both upland and lowland ecosystems. It affects 14 million ha of upland rice production and 19 million ha of lowland rice production worldwide (Pandey et al., 2007). A little effort has been put for developing drought rice cultivars. Many of the varieties grown in rainfed ecosystems were developed for irrigated ecosystems without drought tolerance screening. Farmers prefer these varieties due to their high yielding potential and good grain quality in rainfed regions of South and South East Asia (Vikram et al., 2011). However, because of their susceptibility to moisture stress, these varieties are subjected to significant yield loss due to unanticipated drought stress during the growing season. Traditional rice cultivars grown in rainfed areas, on the other hand, are highly drought tolerant but have low yield potential. Farmers need cultivars that combine high yield potential with improved drought tolerance to ensure high yield in non-drought years and acceptable yield in drought years, further development of drought tolerant varieties in genetic background (Fisher et al., 2003). Thus breeding of drought tolerant varieties with

high yield potential is highly challenging because of its complex physiological and molecular mechanisms, thus selecting genotypes which better available varieties is a major goal under water deficit condition. Thus, mapping populations are developed by combining parents with large differences in target traits, which enables the identification of QTLs for the trait of interest (Bai et al., 2010). In the present study, two varieties contrasting in their response to drought stress, viz., Rasi and ISM were crossed for the development of the RILs (Anila et al., 2017) at ICAR-IIRR, Hyderabad, India. Single seed descent (SSD) method was adopted for advancing the progenies in order to develop the RILs (Collard et al., 2005). A total of 209 RILs along with the parents, tolerant and sensitive checks were screened for drought tolerance under drought stress condition (NPK @ 80:40:40) at ICAR –NRRI – CRURRS, Hazaribagh and the weather factors were recorded for the period of drought screening (Table1).

Table.1: - Weather data recorded at drought screening plot across the two seasons.

Season	Month	Max temp (°C)	Min. temp (°C)	RH (%)	RAIN FALL (mm)	WIND VELOCITY (Km/hr)
WS 2020	June	36.8	24.3	65.5	100.1	7.3
	July	30.8	22.1	83.3	302.0	5.9
	August	29.9	21.9	82.8	169.3	4.9
	September	29.5	21.2	85.4	267.7	6.2
	October	27.8	17.5	83.4	174.2	2.3
	November	26.3	11.1	81.8	0.0	2.3
	December	21.8	7.2	86.1	14.2	3.0
WS 2021	June	31.23	21.87	81.03	380.8	2.59
	July	31.18	22.45	82.48	344.4	2.24
	August	30.61	21.84	85.65	313.8	1.78
	September	30.07	21.05	84.6	289.4	2.17
	October	29.02	17.82	77.35	146.4	1.21
	November	25.98	10.65	75.83	7.2	0.37
	December	22.45	7.64	77.74	48.4	0.99

Data was recorded and the average of two seasons is considered for a total of 12 agro-morphological traits *i.e.*, Days to 50% flowering, Plant Height, Plot yield, Grain yield per hectare, Harvest index, Biomass, Panicle length, No. of Panicles per 0.5 m sample, Grain

weight, Grain number per panicle, Chaffs number per panicle, Sterility percentage. Descriptive statistics of RILs along with parents under stress condition has carried out and represented in Table-2.

UNDER PEER REVIEW

Table.2. Descriptive statistic of 12 traits under drought stress

S. No	Trait	Rasi	ISM	Min	Max	Mean	S.E	CD @ 5%	CD @1%	CV%	Skweness	Kurtosis
1	Days to 50% flowering (Nos)	86.00	110.30	77.00	113.00	96.53	1.48	4.14	5.47	18.89	0.13	2.61
2	Plant Height (cm)	68.70	61.80	34.80	100.40	62.05	4.72	13.18	17.39	10.03	-0.09	2.80
3	Grain Yield per Hectare (kg/ha)	1326.60	550.00	0.00	3500.00	1198.32	170.22	474.79	626.26	19.84	0.47	2.46
4	Harvest Index	0.17	0.04	0.00	0.37	0.13	0.02	0.04	0.05	16.46	0.55	2.73
5	Biomass (g)	220.00	210.00	65.00	325.00	190.28	18.15	50.63	66.79	12.81	-0.40	3.68
6	Plot Yield (g/plot)	261.60	110.00	0.00	700.00	239.66	34.04	94.95	125.25	19.84	0.47	2.46
7	Panicle Length (cm)	18.50	24.60	13.50	24.60	18.42	0.96	2.70	3.56	6.76	-0.13	2.75
8	Panicle Number (Nos)	37.60	20.00	0.00	85.00	46.45	5.84	16.29	21.49	19.37	-0.21	2.81
9	Grain Weighth (g)	35.00	11.00	0.00	120.00	27.23	4.93	13.75	18.14	15.02	0.86	3.24
10	Number of grains per panicle (Nos)	55.30	23.00	0.00	125.40	50.15	6.49	18.11	23.89	16.78	-0.20	2.26
11	Chaffy grains per panicle (Nos)	19.50	93.60	10.00	108.00	46.73	5.30	14.79	19.51	16.33	0.42	2.81
12	Sterility Percentage (%)	26.50	81.07	10.10	100.00	49.49	4.47	12.49	16.48	11.50	0.45	2.29

The days to fifty percent flowering (DFF) under drought stress ranged from 77 ± 0.146 days (RIL-8) to $113 (RIL-82) \pm 0.097$ days, with an overall mean of 96.53 ± 1.48 days, while Rasi and ISM recorded mean values of 86 ± 0.9 days and 110 ± 0.14 days, respectively. Among the population, the late flowering genotypes suffered higher yield reduction than early maturing ones on an average among population. This is mainly because in early flowering genotypes water deficit occurs after the completion of pollination and fertilization and coincides with early grain filling period. Where as in late flowering genotypes drought stress at this period leads to increase in embryo abortion or reduce grain weight (O'Toole and Chang, 1979). O'Toole and Namucow (1983) found that spikelet fertility decreased with decrease in panicle excretion under drought stress, due to entrapment of spikelets in leaf sheath are sterile. Similarly, delay in DFF was reported across several studies in rice under drought stress previously (Lanceras et al., 2004; Venuprasad et al., 2007; 2009, Berneir et al., 2007; Kumar et al., 2007, 2008, 2009; Vikram et al., 2011; Kalmeshwer et al., 2012; Saikumar et al., 2014).

The plant height (PH) under drought stress ranged from 34.8 ± 0.47 cm (RIL-118) to 100.4 ± 0.12 (RIL-54) with an overall mean of 62.05 ± 4.72 , while Rasi and ISM recorded mean values as 68.7 ± 0.029 and 61.8 ± 0.024 , respectively. Similarly, Kumar et al., (2009) reported a reduction in plant height in several field experiments under different stress levels (mild, moderate, and severe stress). The PH reduction was much greater in dry season experiments than in wet season experiments under stress. Drought reduces plant height, leaf area, and crop growth due to impaired mitosis, cell elongation, and expansion (Nonami, 1998). Because of the decrease in turgor pressure, cell growth is the most drought-sensitive (Taiz and Zeiger, 2006). The HI (Harvest Index) under drought stress ranged from 0 (RIL-154) to 0.37 ± 0.0014 (RIL- 33) with an overall mean of 0.13 ± 0.0159 with HI values of 0.17 ± 0.0015 and 0.04 ± 0.00011 respectively, recorded for Rasi and ISM. The Biomass under drought stress ranged from 65 ± 0.97 (RIL-83) to 325 ± 0.97 (RIL- 33) with an overall mean of 190.28 ± 18.15 , with Rasi and ISM recoding values of with 220 ± 0.097 and 210 ± 0.048 , respectively. Grain yield under drought has been reported to be a function of biomass production and harvest index at the vegetative and reproductive stage, respectively (Atlin et al., 2008). The higher severity of reproductive stage stress has caused a significant reduction in HI. Atlin et al., (2008) reported that a large portion of the

variation in yield was explained by variation in HI under reproductive stress conditions. a reduction in yield due to a reduction in biomass, even when there was no significant reduction in HI, thereby producing a further reduction in yield. Genotypes with a greater capacity to produce dry matter in a short growth cycle had the advantage in grain yield. The GYKGPHA (grain yield per Hectare) under drought stress ranged from 0 (RIL-154) to 3500 ± 12.2 (RIL-33), with an overall mean of 1198.32 ± 170.22 , with Rasi and ISM the recorded mean values of 1326.6 ± 2.54 and 550 ± 0.097 respectively. The PLOTYLD (Plot yield) under drought stress ranged from 0 (RIL-154) to $700 \pm .244$ (RIL-33) with an overall mean of 239.66 ± 34.04 , with the parents, Rasi and ISM recording values of $261.6 \pm .097$ and $110 \pm .048$, respectively. The panicle length (PL) under drought stress ranged from 13.5 ± 0.11 (RIL-40) to $24.6 \pm .07$ (66) with an overall mean of 18.42 ± 0.96 . The Rasi and ISM recorded values of $18.5 \pm .03$ and 24.6 ± 0.01 , respectively for PL. The PANNO (No of panicles per 0.5m) under drought stress ranged from 0 (RIL-155) to 85 ± 0.97 (RIL-12) with an overall mean of 46.45 ± 5.84 , while Rasi and ISM recorded values of 37.6 ± 0.04 and $20 \pm .048$ respectively. The GRAINWT (grain weight) under drought stress ranged from 0 (RIL-154) to 120 ± 1.71 (RIL-33) with an overall mean of 27.23 ± 4.93 with Rasi and ISM showing values of 35 ± 0.04 and 11 ± 0.14 , respectively. Significant reduction in grain yield was observed in many drought sensitive genotypes, when drought coincides with reproductive stage in earlier reports (Lanceras et al., 2004; Venuprasad et al., 2007; Berneir et al., 2008; Kumar et al., 2008; Vikram et al., 2011; Kalmeshwer et al., 2012; Krishna et al., 2012; Saikumar et al., 2014). Reduction in GY under stress in relation to control condition was higher in lines with higher DFF due to delay in flowering, which is attributed to slow floral development and reduction in panicle elongation, which are severely influenced by reproductive stage drought stress (Pantuwan et al., 2002; Laffitte et al., 2004).

The GNOPPAN (grain no. per panicle) under drought stress ranged from 0 (RIL-154) to 125.4 ± 1.56 (RIL-152) with an overall mean of 50.15 ± 6.49 . The parents, Rasi and ISM recorded values of 55.3 ± 0.092 and 23 ± 0.097 , respectively. The CNOPPAN (chaffy no. per panicle) under drought stress ranged from 10 ± 0.37 (RIL-112) to 108 ± 0.42 (RIL-155) with an overall mean of 46.73 ± 5.3 , whereas Rasi and ISM recorded values of 19.5 ± 0.019 and 93.6 ± 0.007 , respectively. The STERILITYP (Sterility percentage) under drought stress

ranged from 10.1 ± 0.43 (RIL-112) to $100 \pm .004$ (RIL-154) with an overall mean of 49.49 ± 4.47 . The parents, Rasi and ISM showed values of 26.5 ± 0.01 and 81.07 ± 0.04 respectively. When stress occurs at flowering stage, the most sensitive growth stage, the main yield component affected is the percentage of fertile spikelets and thus the number of grains will be reduced. This trait, highly relevant for any cereal, is critical for rice under water stress (Lafitte, 2002). At reproductive phase, drought results in barrenness, spikelet sterility and reduction in photosynthesis, early senescence and shortens the grain-filling period and decreases remobilization of assimilates to grains leading to reduction in grain number and grain weight collectively (Plaut, 2003; Yang and Zhang, 2006). The failure of panicle exertion alone accounts for approximately 25–30% of spikelet sterility because the unexserted spikelets cannot complete anthesis and shed pollen, even when development is otherwise normal (O'Toole and Namuco, 1983).

A continuous distribution of phenotypic values for various agro morphological traits recorded for drought tolerance supports quantitative inheritance of various traits. In the histogram, the Y-axis (vertical axis) represents the frequency count *i.e.*, number of RILs, while the X-axis (horizontal axis) represents the trait measured (Figure 1).




Figure 1 - Frequency distribution of recombinant inbred lines for 12 traits, based on screening for drought tolerance under stress condition.

In the present study, it was found that the distribution of RILs for plant height, biomass, panicle length, number of panicles and grain number per panicles show negative skewness. In contrast the distribution of days to 50% flowering, grain yield per hectare, harvest index, plot yield, grain weight, chaffy number per panicle, sterility percent showed positive skewness. Maximum skewness was observed for grain weight (0.86) and minimum for plant height (-0.09). Positive skewness towards tolerance is related to complementary gene interaction whereas negative for Duplicate gene interaction. Kurtosis is a statistical measure which shows the degree of the score cluster in peak or tail of a variable frequency distribution. Peak is the tallest part of the distribution curve whereas tails are the ends of the curve. Kurtosis is divided into three –a normal distribution with kurtosis three is termed as mesokurtic, a normal distribution with kurtosis more than three (excess kurtosis > 0) is termed as leptokurtic and a normal distribution with kurtosis less than three (excess kurtosis < 0) is termed as platykurtic (Savitha and Usha, 2015). In the present study, day to 50% flowering (2.61), plant height (2.8), grain yield per hectare (2.46), harvest index (2.73), plot yield (2.46), panicle length (2.75), panicle number (2.81), grain number per panicle (2.26), chaffy number per panicle (2.81), sterility percent (2.29) showed kurtosis value below three, thus platykurtic distribution while the rest traits such as biomass (3.68), grain weight (3.24) showed value above three which means that they are showing leptokurtic distribution. Additionally, a positive kurtosis shows the presence of gene interaction, whereas a negative value denotes the absence of gene interaction. (Pooni et al., 1997).

The study has identified some promising RILs that have a good level of tolerance to drought. A total of 59 lines recorded a very good level of tolerance to drought with reasonable yield levels under drought stress, as compared to the sensitive parent, ISM. Two RILs, RIL-33 and RIL-58 also possessed the highly desirable characters with a medium slender grain type. Both the lines showed better yield related parameters and biomass in

comparison to ISM, indicating a key role for these traits with respect to drought tolerance (Table.3).

Table 3: Comparison between parents and RILs for different trait under drought stress

Entry	DFF	PH	GYKH	HI	BM	PLY	PANNO	GW	GNOPPAN	SP
RIL-33	85	84.1	3500	0.37	325	700	73	120	75.6	29.7
RIL-58	94	80.8	2600	0.27	255	520	42	70	66.6	35.6
ISM	110	61.5	550	0.05	210	110	20	10	22	81.0
Rasi	87	68.4	1300	0.16	220	260	38	35	54.4	26.5

DFF-days to 50%flowering, PH-plant height, GYKH-grain yield per hectare, HI-Harvest index, BM-Biomass, PLY-plot yield, PANNO-number of panicles, GW-grain weight, GNOPPAN-grain number per panicle, SP-sterility percent.

Conclusion:

In conclusion, the two RILs (RIL-33 and RIL-58) have highly desirable characters such as, high yield under drought, early maturing, medium-slender grain type and better plant height along with tolerance to drought condition in comparison to both the parents. We are advancing these two lines along with other RILs possessing high yield under drought stress for evaluation in larger plots. Cultivation of such lines possessing tolerance to drought can enhance the productivity of Samba Mahsuri farmers. The present study also concluded the skewness of the lines towards tolerance suggests a possibility of identifying one or more main QTLs related with tolerance, according to analysis of a few important phenotypic metrics as well as yield data. The population will soon be genotyped using a set of molecular markers for possible identification and characterization of the component loci associated with tolerance.

References

1. Anila, M., Giri, A. and Sundaram, R.M., 2022. Evaluation of recombinant inbred lines for low soil phosphorous tolerance derived from Rasi-a low soil phosphorous tolerant variety. *Electronic Journal of Plant Breeding*, 13(3), pp.800-809.
2. Anila, M., Mahadevaswamy, H.K., Kale, R.R, Bhadana, V.P, Brajendra, P., Hajira, S.K., Pranathi, K., Balachiranjeevi, CH., Praveen, M., Abhilash Kumar, V., Bhaskar, S., Dhanalaxmi, D., Harika, G., Dilipkumar, T., Kousik, MBVN., Yugender, A., Punniakotti, E., Laxmi Prasanna, B., Swapnil, K., Rekha, G., Balachandran, S.M., Neeraja, C.N., Madhav, M.S., Surekha, K., Laha, G.S., Anantha, M.S, Fiyaz, A., Suneetha, K., Gireesh, C., Viraktamath, B.C., Ravindra Babu, V and Sundaram, R.M. 2017. Marker Assisted Introgression of *Pup1*, A Major QTL Associated with Tolerance to Low Soil Phosphorus into Elite Rice Varieties. *ICAR-IIRR Newsletter* –15 (1):26-29.
3. Atlin G, Venuprasad R, Bernier J, Zhao D, Virk P and Kumar A. 2008. Rice germplasm development for drought-prone environments: Progress made in breeding and genetic analysis at the International Rice Research Institute. Drought frontiers in rice: Crop improvement for increased rainfed production. World Scientific Publishing, Singapore, Singapore, and International Rice Research Institute, Los Baños, Philippines. 35-59.
4. Bai, X., Luo, L., Yan, W., Kovi, M. R., Zhan, W., & Xing, Y. (2010). Genetic dissection of rice grain shape using a recombinant inbred line population derived from two contrasting parents and fine mapping a pleiotropic quantitative trait locus qGL7. *BMC genetics*, 11(1), 1-11.
5. Bernier J, Kumar A, Venuprasad R. Spaner D, Atlin GN. 2007. A large-effect QTL for grain yield under reproductive-stage drought stress in upland rice. *Crop Sci.* 47: 507-516.
6. Collard, B. C., Jahufer, M. Z. Z., Brouwer, J. B., & Pang, E. C. K. (2005). An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica*, 142(1), 169-196.
7. Dhawan, G.; Kumar, A.; Dwivedi, P.; Gopala Krishnan, S.; Pal, M.; Vinod, K.K.; Bhowmick, P.K.; Bollinedi, H.; Ellur, R.K.; Ravikiran, K.T.; et al. Introgression of qDTY1. 1 Governing Reproductive Stage Drought Tolerance into an Elite Basmati Rice Variety "Pusa Basmati 1" through Marker Assisted Backcross Breeding. *Agronomy* 2021, 11, 202
8. Fischer KS, Lafitte R, Fukai S, Atlin G, Hardy B. 2003. Breeding Rice for Drought Prone Environments. International Rice Research Institute, Los Banos, Philippines, pp 5- 22.
9. Kalmeshwer PG, Varma CMK, Saikumar S, Kiran B, Shenoy V, Shashidhar HE. 2012. Direct Selection for Grain Yield under Moisture Stress in *Oryza sativa* cv IR58025B × *Oryza meridionalis* Population. *Crop Sci.* 52, 644-653.
10. Kilimo, T. Expanding Rice Markets in the East African Community. Regional Solution to Local Problems; Kilimo Trust Head Quarters: Kampala, Uganda, 2018; p. 78.

11. Krishna GH, Lenie AQ, Vikram P, Swamy BPM, Dixit S, Ahmed H, Jose EH, Teresita HB, Kumar A. 2012. Identification and mapping of a QTL (qDTY1 1) with a consistent effect on grain yield under drought. *Field Crops Res* 131: 88-96.
12. Kumar A, Bernier J, Verulkar S, Lafitte HR, Atlin GN. 2008. Breeding for drought tolerance, direct selection for yield, response to selection and use of drought tolerant donors in upland and lowland-adapted populations. *Field Crops Res* 107: 221-231.
13. Kumar A, Venuprasad R, Atlin GN. 2007. Genetic analysis of rainfed lowland rice drought tolerance under naturally-occurring stress in eastern India, heritability and QTL effects. *Field Crops Res.* 103, 42-52.
14. Kumar A, Verulkar SB, Dixit S, Chauhan B, Bernier J, Venuprasad R, Zhao D, Shrivastava MN. 2009. Yield and yield-attributing traits of rice (*Oryza sativa* L) under lowland drought and suitability of early vigor as a selection criterion. *Field Crops Res.* 114: 99-107.
15. Lafitte HR, Ismail A, Bennett J. 2004. Abiotic stress tolerance in rice for Asia, progress and the future. In, Fischer, T., et al. (Eds.), *New Directions for a Diverse Planet*, Proceedings of the 4th International Crop Science Congress. Brisbane., Australia.
16. Lafitte HR. 2002. Relationship between leaf relative water content during reproductive stage water deficit and grain formation in rice. *Field Crop Res.* 76: 165-174.
17. Lanceras J, Pantuwan G, Jongdee, B, Toojinda T. 2004. Quantitative trait loci associated with drought tolerance at reproductive stage in rice. *Plant Physiol.* 135: 84-399.
18. Li, Z.K. and Xu, J.L., 2007. Breeding for drought and salt tolerant rice (*Oryza sativa* L.): progress and perspectives. *Advances in molecular breeding toward drought and salt tolerant crops*, pp.531-564.
19. Narciso J and Hossain M. 2002. World Rice Statistics, <http://www.irri.org/science/ricestat>.
20. Nonami H. 1998. Plant water relations and control of cell elongation at low water potentials, *J. Plant Res.* 111: 373-382.
21. O'Toole JC, Chang TT. 1979. Drought resistance in cereal. Rice: A case study. In Mussell, H., Staples, R.c. (Eds), *Stress physiology in crop plants*. Wiley. New york, pp. 374- 405.
22. O'Toole JC & Namuco OS. 1983. Role of panicle exertion in water-stress induced sterility. *Crop Sci.* 23: 1093-1097.
23. Pandey S, Bhandari H, Hardy B. 2007, Economic costs of drought and rice farmers' coping mechanisms: a cross-country comparative analysis from Asia. Los Baños (Philippines): International Rice Research Institute, 203.

24. Pandey, V.; Shukla, A. Acclimation and tolerance strategies of rice under drought stress. *Rice Sci.* 2015, 22, 147–161.
25. Pantuwan G, Fukai S, Cooper M, Rajatasereekul S, O'Toole JC. 2002. Yield response of rice (*Oryza sativa* L) genotypes to drought under rainfed lowlands, 2 Selection of drought resistant genotypes. *Field Crops Res.* 73: 169-180.
26. Plaut Z. 2003. Plant exposure to water stress during specific growth stages, *Encyclopedia of Water Science*, Taylor & Francis, pp. 673– 675.
27. Pooni HS, Jinks JL and CornishMA 1977. The causes and consequences of non-normality in predicting the properties of recombinant inbred lines. *Heredity*, 38:329-338.
28. Sahebi, M.; Hanafi, M.M.; Rafii, M.Y.; Mahmud, T.M.M.; Azizi, P.; Osman, M.; Abiri, R.; Taheri, S.; Kalhori, N.; Shabanimofrad, M.; et al. Improvement of drought tolerance in rice (*Oryza sativa* L.): Genetics, genomic tools, and the WRKY gene family. *Biomed. Res. Int.* 2018, 18, 20.
29. Saikumar S, Gouda KP, Saiharini A, Varma CMK, Vineesha O, Padmavathi G, Shenoy VV. 2014. Major QTL for enhancing rice grain yield under lowland reproductive drought stress identified using an *O. sativa/O. glaberrima* introgression line. *Field Crops Res.* 163: 119–131.
30. Sakran, R.M.; Ghazy, M.I.; Rehan, M.; Alsohim, A.S.; Mansour, E. Molecular genetic diversity and combining ability for some Physiological and Agronomic Traits in Rice under Well-Watered and Water-Deficit Conditions, *Plants* 2022, 11(5), 702.
31. Savitha, P., & Usha Kumari, R. (2015). Assessment of genetic variability and correlation studies Among traditional land races and improved cultivars for Segregating generations of rice (*Oryza sativa* L.). *Int. J. Sci and nature*, 6(2), 135-140.
32. Schenker, S. An overview of the role of rice in the UK diet. *Nutr. Bull.* 2012, 37, 309–323.
33. Sundaram, R. M., Vishnupriya, M. R., Biradar, S. K., Laha, G. S., Reddy, G. A., Rani, S. N., Sarma, N. P and Sonti, R. V., 2008, Marker assisted introgression of bacterial blight resistance in Samba mahsuri, an elite *indica* rice variety. *Euphytica*. 160: 411-422.
34. Taiz L, Zeiger E. 2006. *Plant Physiology*. 4th Ed., Sinauer Associates Inc. Publishers, Massachusetts. Yang J.C., Zhang, J.H., 2006. Grain filling of cereals under soil drying. *New Phytologist*. 169: 223–236.
35. V R Babu, Ch Padmavathi, CN Neeraja, D Krishnaveni, Divya Balakrishnan, LV Subba Rao, P Raghuvveer Rao, GSV Prasad, Gururaja Katti, T Ram, D Subrahmanya, RM Kumar, AS Hari Prasad, K Surekha, MS Prasad and U Chaitanya. 2016. 50 years of AICRIP. way forward. Technical Bulletin No.92/2016. ICAR-Indian Institute of Rice Research, Rajendranagar, Hyderabad-500 030, Telangana State, India, 292 pp.
36. Venuprasad R, Dalid CO, Del Valle M, Zhao D, Espiritu M, Sta Cruz MT, Amante M, Kumar A, Atlin GN. 2009. Identification and characterization of large-effect

- quantitative trait loci for grain yield under lowland drought stress in rice using bulksegregant analysis. *Theor. Appl. Genet.* 120: 177-190.
37. Venuprasad R, Lafitte HR, Atlin GN. 2007. Response to direct selection for grain yield under drought stress in rice. *Crop Sci.* 47: 285-293.
38. Vikram P, Swamy BPM, Dixit S, Sta Cruz T, Ahmed HU, Singh AK, Kumar A. 2011. qDTY11, a major QTL for rice grain yield under reproductive-stage drought stress with a consistent effect in multiple elite genetic backgrounds. *BMC Genet.* 12: 89.
39. Waheed, R.; Ignacio, J.C.; Arbelaez, J.D.; Juanillas, V.M.; Asif, M.; Henry, A.; Kretschmar, T.; Arif, M. Drought response QTLs in a Super Basmati x Azucena population by high-density GBS-based SNP linkage mapping. *Plant Breed.* 2021, 140, 758–774
40. Yang JC & Zhang JH. 2006. Grain filling of cereals under soil drying. *New Phytologist.* 169: 223–236.

UNDER PEER REVIEW