

Original Research Article

Dynamics of zinc fractions in soil as affected by zinc fertilization in a maize-maize cropping sequence in Upper Brahmaputra Valley Zone of Assam

ABSTRACT:

Zinc is considered as an important micronutrient for maize which is an promising emerging cash crop for the state of Assam. A field experiment was conducted at the experimental farm of Krishi Vigyan Kendra, Jorhat, Assam Agricultural University during 2018-19 and 2019-20 to study the dynamics of zinc fractions in soil in a maize-maize cropping sequence as affected by zinc fertilization. The experiment was laid down in a randomized block design (RBD) with twelve treatments combinations of basal and foliar application of zinc through zinc sulphate and zinc oxide nano particle (ZnO NPs). The results revealed that zinc fertilization significantly influenced the studied zinc fractions in soil during both the years under investigation. The distribution of different zinc fractions in soils at harvest was found in the following order: water soluble plus exchangeable-Zn < complexed-Zn < crystalline sesquioxide bound-Zn < amorphous sesquioxide bound-Zn < residual-Zn. Among the zinc treatment combinations, soil application of Zn @ 3.75 kg ha⁻¹ in combination with foliar application of 500 ppm ZnO NPs exhibited the highest concentration of water soluble plus exchangeable-Zn content (1.10 ppm) in soil. The highest concentration of complexed-Zn (2.95 ppm), amorphous sesquioxide-bound-Zn (4.11 mg kg⁻¹), crystalline sesquioxide-bound-Zn (3.76 ppm), residual-Zn (125.65 ppm) and total-Zn (137.33 ppm) were recorded in the treatment receiving soil application of Zn @ 7.5 kg ha⁻¹. Among the zinc fractions studied, the concentration and percent contribution of water soluble plus exchangeable-Zn to total-Zn was the lowest while residual-Zn fraction contributed the highest to the total soil zinc pool. Furthermore, path coefficient analysis revealed that the water soluble plus exchangeable-Zn had the highest contribution towards the plant available DTPA extractable-Zn in soil. Moreover, all the fractions of zinc were found to be significantly and positively correlated with each other indicating existence of dynamic equilibrium of zinc in soil.

Keywords: Zinc fractions, cropping sequence, Zinc oxide nano particle, zinc fertilization, foliar application

INTRODUCTION:

Zinc is considered as the fourth most important yield limiting nutrient after nitrogen, phosphorus and potassium. The plant available DTPA extractable zinc in Indian soils is less than 1% of total zinc [1]. Availability of zinc is reported to be associated with the transformation of zinc in soils and plant continuum through various mechanisms like adsorption by clay surfaces, hydrous oxide minerals, organic matter etc which affect Zn uptake by crops. Also, the availability of zinc to plants has been reported to be associated with the distribution of different zinc fractions in soil. Therefore, understanding the distribution of various zinc fractions in soils help to characterize the dynamics of Zn in soils as well as possible contribution of individual zinc fractions towards plant availability.

The distribution of Zn among various chemical forms may vary significantly in response to changing soil properties [2]. For a better perceptive, total Zn in soil can be broadly divided into five mechanistic fractions viz., water soluble plus exchangeable zinc (WSEx-Zn), complexed zinc (Comp-Zn), amorphous sesquioxide bound zinc (ASB-Zn), crystalline sesquioxide bound zinc (CSB-Zn), residual zinc (Res-Zn) and total zinc (Total-Zn) which can be quantified using sequential fractionation schemes [3]. These fractions provides a detailed information on the geological, biological and chemical processes occurring in the soil system and give a detailed account of the available Zn for plants uptake. Oxide bound and residual Zn is known to be more stable while water soluble and exchangeable fractions are more soluble [4]. The dynamic equilibrium between these fractions is influenced by soil properties such as soil texture, pH and soil organic matter status of soil [5].

Maize-maize cropping sequence is the most promising but exhaustive cropping sequence followed in upper Brahmaputra valley zone of Assam. The sequence is very much sensitive to fertilization and often shows deficiency of macro as well as micro nutrients. Furthermore, micronutrient deficiency primarily of zinc is often prevalent in upper Brahmaputra valley zone of Assam. However, systematic information on the dynamics of different zinc fractions in soil in maize-maize cropping sequence and their relationship with soil properties in the acidic soils of Assam is very limited. Therefore, the present study was aimed at obtaining a more detailed and critical information and understanding the dynamics of different fractions of zinc and their relationship with soil properties in a maize-maize cropping sequence in Upper Brahmaputra Valley Zone of Assam.

Materials and Methods

The field experiment was conducted at the experimental farm of Krishi Vigyan Kendra, Jorhat, Assam Agricultural University during 2018-19 and 2019-20 in both *rabi* as well as *kharif* season. The experimental site is situated in the Upper Brahmaputra Valley Zone (UBVZ) of Assam at latitude 26.8308⁰ N and longitude 94.4565⁰ E at an altitude of 112 m above mean sea level. The experiment was laid down in a randomized block design (RBD) with twelve treatments *viz.*, T1-Control, T2- Soil application of Zn @ 2.5 kg ha⁻¹, T3- Soil application of Zn @ 5.0 kg ha⁻¹, T4- Soil application of Zn @ 7.5 kg ha⁻¹, T5- Foliar application of 0.5% Zn in three sprays (knee high, tasseling and silking stage), T6- Foliar application of 500 ppm ZnO NPs in three sprays (knee high, tasseling and silking stage), T7- T₅ + Soil application of Zn @ 1.25 kg ha⁻¹, T8- T₅ + Soil application of Zn @ 2.5 kg ha⁻¹, T9- T₅ + Soil application of Zn @ 3.75 kg ha⁻¹, T10- T₆ + Soil application of Zn @ 1.25 kg ha⁻¹, T11- T₆ + Soil application of Zn @ 2.5 kg ha⁻¹ and T12- T₆ + Soil application of Zn @ 3.75 kg ha⁻¹ with three replications, consisted of zinc applied as soil with or without foliar application through zinc sulphate and zinc oxide nano particle (ZnO NPs). The recommended dose of N, P and K were applied as urea, SSP and MOP, respectively. Soil samples from 0-15 cm depth from each plot were drawn before sowing and after harvest of crop and analysed for zinc fractions. Initial soil properties are given in table 1.

Table 1. Initial physico-chemical properties of the experimental site

Soil Characteristics	Value
A. Textural class	Sandy loam
B. Chemical properties	
i) Soil reaction (pH)	5.29
ii) Organic carbon (%)	0.58
iii) Available N (kg ha ⁻¹)	188.00
iv) Available P ₂ O ₅ (kg ha ⁻¹)	25.89
v) Available K ₂ O (kg ha ⁻¹)	151.34
xii) Water soluble plus exchangeable-Zn (mg kg ⁻¹)	0.33
xiii) Complexed-Zn (mg kg ⁻¹)	1.80
xiv) Amorphous sesquioxide bound-Zn (mg kg ⁻¹)	2.82
xv) Crystalline sesquioxide bound-Zn (mg kg ⁻¹)	2.42
xvi) Residual-Zn (mg kg ⁻¹)	106.63
xvii) Total-Zn (mg kg ⁻¹)	113.86

Chemical fractionation of zinc

Different zinc fractions in soils are sequentially extracted following the procedure of Black, [3] as shown in Table 2.

Table 2. Zinc sequential fractionation procedures.

Fractions	Solution (ml)	Soil (g): solution (ml)	Conditions
(1) Water soluble + exchangeable-Zn (WSE _x -Zn)	1 M NH ₄ OAc at pH 7.2	5:20	Shaken for 1 h, Centrifuge
(2) Complexed-Zn (Comp-Zn)	20 ml 0.5 M Cu (OAc) ₂	5:20	Shaken for 1 h, Centrifuge
(3) Amorphous sesquioxide bound-Zn (ASB-Zn)	0.2 M acidified NH ₄ (OX) ₂ (pH 3.0)	5:20	Shaken for 1 h, Centrifuge
(4) Crystalline sesquioxide bound-Zn (CSB-Zn)	40 ml 0.3 M Sodium Citrate + 5 ml 0.1M NaHCO ₃ + 1 g Na ₂ S ₂ O ₄ Kept on the water bath (70-80°C)	5:20	Stirred and kept on the water bath at a temperature of 70-80°C for 10 min For 15 min with occasional stirring Centrifuge after cooling

Total zinc (Total-Zn)

For analysis of total zinc, 0.1 g of each soil was digested with few drops of H₂SO₄ and 5 ml of HF + 0.5 ml of HClO₄ in a 30 ml capacity platinum crucible [6]. When the residue completely dissolved in 6 N HCl, the content of the crucible was transferred to 50 ml volumetric flask and volume made up. Digests were analyzed for zinc using ICP OES (Inductively coupled plasma-optical emission spectrometry) (Model: Agilent 5110).

Residual zinc (Res-Zn)

Res-Zn is calculated as the difference between total zinc and sum of the zinc fractions *viz.*, WSE_x-Zn, Comp-Zn, ASB-Zn and CSB-Zn.

Statistical analysis

Fisher's method of analysis of variance was resorted to determine the statistical significance among the data recorded from various treatments in randomized block design. The respective F values as well as least significant differences were calculated at 5% probability level to determine the statistical significance among the variances of the different

treatments. Moreover, path analysis was carried out in IBM SPSS to understand the contribution of different Zn fractions to plant available pool.

Results and Discussion

Influence of zinc fertilization on zinc fractions in soil

The zinc fractions studied were water soluble plus exchangeable zinc (WSEx-Zn), complexed zinc (Comp-Zn), amorphous sesquioxide bound zinc (ASB-Zn), crystalline sesquioxide bound zinc (CSB-Zn), residual zinc (Res-Zn) and total zinc (Total-Zn).

Water soluble and exchangeable zinc fractions (WSEX-Zn)

WSEx-Zn content (pooled for two year) of the soil ranged from 0.30 to 1.10 ppm (Table 3). Among the treatments, highest concentration of WSEx-Zn (1.10 ppm) was recorded in the treatment receiving foliar application of 500 ppm ZnO NPs in combination with soil application of Zn @ 3.75 kg ha⁻¹ and the lowest value was recorded (0.30 mg kg⁻¹) in the control which might be due to higher contribution of applied Zn towards this fraction due to foliar application of ZnO NPs as well as soil application of zinc. Moreover, contribution of WSEx-Zn to total zinc (0.86%) was also the highest in the same treatment. Similar results were also reported by Bala *et al.* [7] that foliar application of ZnO NPs significantly influenced Zn content in soil and recorded the highest Zn content in soil with foliar spray treatment with 5.0 g L⁻¹ ZnO NPs. Analogous results were also reported by Verma *et al.* [8] and Ghoneim [9]. The lowest value of WSEx-Zn fractions in soil in the control treatment might be attributed to continuous uptake by the crop with no external application of zinc fertilizer in these plots.

Complexed-Zn (Comp-Zn)

Perusal of data in Table 3 indicated significant increase in the Comp-Zn in soil with increase in the rate of zinc application. Soil application of Zn @ 7.5 kg ha⁻¹ recorded significantly highest value of Comp-Zn (2.95 mg kg⁻¹) while, the lowest content (1.77 mg kg⁻¹) was recorded in the control. However, no significant difference in Comp-Zn in soil was observed among the foliar treatments. The percent contribution of Comp-Zn to Total-Zn (2.16%) was highest in the treatment receiving soil application of Zn @ 7.5 kg ha⁻¹. Results further revealed that increasing rate of soil application of zinc in combination with foliar application either with Zn (T₇ to T₉) or with ZnO NPs (T₁₀ to T₁₂) had recorded an increase in Comp-Zn content in soil (Table 3).

Soil organic matter is an important soil constituent that influence availability of nutrients in soil. It consists of a range of organic compounds such as humic substances, organic acids of low and high molecular weight, carbohydrates, proteins, peptides, amino acids, lipids, waxes, polycyclic aromatic hydrocarbons and lignin fragments [10]. Many of these components of soil organic matter have a strong affinity to bind Zn [11,12]. The most stable organic substances *viz.*, humic and fulvic acids contain a large number of functional groups (OH, COOH, SH, C=O) that have a great affinity for Zn [13]. The amount of zinc in organically bound form showed an increasing trend with increase in the rate of zinc application. Such increase might probably be due to release of applied zinc to available form and subsequent chelation by organic compounds. This increase in Comp-Zn fraction might also be attributed to the mechanism of chemisorption and complexation by organic ligands [14].

Table 3. Effect of zinc fertilization on WSEx-Zn and Comp-Zn (mg kg^{-1}) in soil

Treatments	WSEx-Zn			Comp-Zn		
	Pooled (<i>Khari</i> f 2018-19 and 2019-20)	Pooled (<i>Rabi</i> 2018-19 and 2019-20)	Pooled Sequence	Pooled (<i>Khari</i> f 2018-19 and 2019-20)	Pooled (<i>Rabi</i> 2018-19 and 2019-20)	Pooled Sequence
T₁	0.31 ^e	0.30 ^g	0.30 ^g	1.76 ^d	1.75 ^d	1.77 ^d
T₂	0.49 ^{cd}	0.50 ^{ef}	0.49 ^{ef}	2.34 ^{bc}	2.34 ^{abcd}	2.36 ^c
T₃	0.57 ^{cd}	0.57 ^{de}	0.56 ^{de}	2.52 ^b	2.57 ^{ab}	2.58 ^{ab}
T₄	0.84 ^b	0.81 ^c	0.82 ^c	2.96 ^a	2.90 ^a	2.95 ^a
T₅	0.35 ^e	0.35 ^g	0.35 ^g	1.84 ^d	1.83 ^{cd}	1.86 ^d
T₆	0.55 ^{de}	0.54 ^f	0.55 ^{de}	1.85 ^d	1.86 ^{cd}	1.87 ^d
T₇	0.40 ^e	0.41 ^g	0.41 ^g	2.04 ^{cd}	2.03 ^{bcd}	2.00 ^{cd}
T₈	0.57 ^c	0.56 ^{de}	0.56 ^{de}	2.37 ^b	2.40 ^{abc}	2.40 ^{bc}
T₉	0.62 ^c	0.65 ^d	0.64 ^d	2.52 ^b	2.50 ^{ab}	2.53 ^{ab}
T₁₀	0.78 ^b	0.78 ^{bc}	0.78 ^c	2.05 ^{cd}	2.09 ^{bcd}	2.03 ^{cd}
T₁₁	0.94 ^b	0.93 ^b	0.94 ^b	2.39 ^b	2.41 ^{abc}	2.41 ^{bc}
T₁₂	1.11 ^a	1.11 ^a	1.10 ^a	2.48 ^b	2.47 ^{ab}	2.49 ^b

LSD (p≤0.05)	0.13	0.13	0.12	0.32	0.59	0.38
CV (%)	12.92	12.87	11.73	8.36	15.61	9.98

Mean values in each column followed by similar letters in superscript are not significantly different at 5% probability level

Amorphous sesquioxide bound-Zn (ASB-Zn)

Significantly higher content of ASB-Zn (4.11 ppm) was recorded in the treatment receiving soil application of Zn @ 7.5 kg ha⁻¹ and the lowest (2.69 mg kg⁻¹) in the control (Table 4). Soil application of Zn @ 7.5 kg ha⁻¹ recorded the highest percent contribution of ASB-Zn (3.00%) to total zinc (Table 6). The increase in ASB-Zn fraction with increase in the rate of zinc application could be attributed to addition of zinc fertilizer and subsequent transformation to this fraction due to presence of amorphous sesquioxide in the soil. These findings were in agreement with Wijibandara [15], Veeranagappa *et al.* [16] and Talukdar *et al.* [17]. Higher content of ASB-Zn fraction as compared to CSB-Zn as well as WSEx-Zn and Comp-Zn might be attributed to the greater ability of amorphous sesquioxides to adsorb zinc because of their high specific surface area [18],[19] and [20]. Also, due to the acidic environment which favoured formation of amorphous sesquioxide bound-Zn in soil.

Crystalline sesquioxide bound-Zn (CSB-Zn)

Crystalline sesquioxide bound fraction of zinc (CSB-Zn) in soil is the predominant fraction due to presence of crystalline iron oxide in the soil. Among the treatments, soil application of Zn @ 7.5 kg ha⁻¹ recorded significantly higher content of CSB-Zn (3.76 mg kg⁻¹) and the lowest (2.35 mg kg⁻¹) was recorded in the control in after harvest. The percent contribution of CSB-Zn (2.77%) to Total-Zn was also highest in the treatment receiving soil application of Zn @ 7.5 kg ha⁻¹ (Table 4). The increase in CSB-Zn with increase in the rate of zinc application might be due to chemical affinity or specific adsorption and also due to predominance of crystalline iron oxide content in soil [21]. Zinc is said to be specifically adsorbed because the ion is sorbed by surfaces of synthetic oxides of Fe and Al (goethite, gibbsite) that have a net positive charge, so despite electrostatic repulsion, Zn is still adsorbed in significant amounts [22], [23]. The higher contribution of sesquioxide bound-Zn might be attributed to higher amount of Zn adsorption on the surface of these oxides because of the higher concentrations of these ions in the soil [19].

Table 4. Effect of zinc fertilization on ASB-Zn and CSB-Zn (mg kg⁻¹) in soil

Treatments	ASB-Zn			CSB-Zn		
	Pooled (<i>Kharif</i> 2018-19 and 2019- 20)	Pooled (<i>Rabi</i> 2018-19 and 2019- 20)	Pooled Sequence	Pooled (<i>Kharif</i> 2018-19 and 2019- 20)	Pooled (<i>Rabi</i> 2018-19 and 2019- 20)	Pooled Sequence
T₁	2.75 ^d	2.74 ^d	2.69 ^d	2.40 ^c	2.39 ^e	2.35 ^c
T₂	3.24 ^{bcd}	3.26 ^{bcd}	3.25 ^{bcd}	3.05 ^{abc}	3.03 ^{bcd}	3.09 ^b
T₃	3.73 ^{ab}	3.74 ^{ab}	3.71 ^{ab}	3.43 ^{ab}	3.40 ^{ab}	3.38a ^b
T₄	4.10 ^a	4.13 ^a	4.11 ^a	3.80 ^a	3.82 ^a	3.76 ^a
T₅	2.87 ^d	2.88 ^{cd}	2.82 ^d	2.45 ^c	2.44 ^{de}	2.41 ^c
T₆	2.75 ^d	2.76 ^d	2.73 ^d	2.47 ^c	2.49 ^{cde}	2.48 ^c
T₇	3.14 ^{cd}	3.12 ^{bcd}	3.13 ^{cd}	2.92 ^{bc}	2.89 ^{bcde}	2.85 ^{bc}
T₈	3.45 ^{bc}	3.48 ^{abc}	3.45 ^{bc}	3.11 ^{abc}	3.08 ^{bc}	3.15 ^b
T₉	3.55 ^{abc}	3.56 ^{abc}	3.48 ^{bc}	3.32 ^{ab}	3.41 ^{ab}	3.33 ^{ab}
T₁₀	3.14 ^{cd}	3.14 ^{bcd}	3.13 ^{cd}	2.89 ^{bc}	2.91 ^{bcde}	2.84 ^{bc}
T₁₁	3.46 ^{bc}	3.49 ^{abc}	3.46 ^{bc}	3.14 ^{abc}	3.13 ^b	3.13 ^b
T₁₂	3.53 ^{abc}	3.55 ^{abc}	3.49 ^{bc}	3.37 ^{ab}	3.39 ^{ab}	3.36 ^{ab}
LSD (p≤0.05)	0.58	0.69	0.61	0.75	0.61	0.65
CV (%)	10.26	12.16	10.84	14.64	11.95	12.80

Mean values in each column followed by similar letters in superscript are not significantly different at 5% probability level

Residual-Zn (Res-Zn)

Residual-Zn was the dominant fraction and considered as zinc reservoir in soil. The percent contribution of this fraction to total zinc was the highest among all other fractions studied (Table 5). The effect of zinc fertilization on residual-Zn in soil was found to be significant for both the years under investigation. The highest concentration of residual-Zn (125.65 mg kg⁻¹) was found in the treatment receiving Zn @ 7.5 kg ha⁻¹ as soil application followed by the treatment receiving soil application of Zn @ 5.0 kg ha⁻¹ (120.31 mg kg⁻¹)

while, the lowest concentration residual-Zn (106.25 mg kg⁻¹) was recorded in the control. Residual-Zn fraction consisting of large proportion of total zinc was also reported by Singh [23] and Priyanka *et al.* [24] which might be due to conversion of labile zinc to non-labile form.

Table 5. Effect of zinc fertilization on Res-Zn and Total-Zn (mg kg⁻¹) in soil

Treatments	Res-Zn			Total-Zn		
	Pooled (<i>Kharij</i> 2018-19 and 2019- 20)	Pooled (<i>Rabi</i> 2018-19 and 2019-20)	Pooled Sequence	Pooled (<i>Kharij</i> 2018-19 and 2019- 20)	Pooled (<i>Rabi</i> 2018-19 and 2019- 20)	Pooled Sequence
T₁	106.42 ^g	106.09 ^g	106.25 ^g	113.63 ^h	113.27 ^h	113.45 ^j
T₂	115.74 ^{cd}	115.90 ^{cd}	115.82 ^{de}	124.85 ^{ef}	125.02 ^{de}	124.93 ^{fg}
T₃	120.55 ^b	120.06 ^b	120.31 ^b	130.80 ^b	130.34 ^b	130.57 ^b
T₄	125.39 ^a	126.91 ^a	125.65 ^a	137.09 ^a	137.57 ^a	137.33 ^a
T₅	110.48 ^{fg}	109.68 ^f	109.58 ^f	116.98 ^g	117.17 ^g	117.08 ⁱ
T₆	110.36 ^f	110.61 ^f	110.48 ^f	117.89 ^g	118.15 ^{fg}	118.02 ⁱ
T₇	111.12 ^{ef}	111.87 ^{ef}	111.49 ^f	119.61 ^g	120.33 ^f	119.97 ^h
T₈	116.22 ^{cd}	116.13 ^{cd}	116.17 ^{de}	125.73 ^{de}	125.68 ^{de}	125.71 ^{ef}
T₉	118.85 ^{bc}	118.93 ^{bc}	118.89 ^{bc}	128.86 ^{bc}	129.06 ^{bc}	128.96 ^{bc}
T₁₀	114.04 ^{de}	114.45 ^{de}	114.24 ^e	122.85 ^f	123.38 ^e	123.11 ^g
T₁₁	116.57 ^{cd}	117.51 ^{bc}	117.04 ^{cd}	126.51 ^{cde}	127.48 ^{cd}	126.10 ^{de}
T₁₂	117.34 ^c	118.14 ^{bc}	117.74 ^{cd}	127.82 ^{cd}	128.66 ^{bc}	128.24 ^{cd}
LSD (p≤0.05)	3.89	3.09	2.44	2.75	2.47	1.83
CV (%)	12.22	13.23	12.66	13.24	9.85	10.38

Mean values in each column followed by similar letters in superscript are not significantly different at 5% probability level

Total Zinc (Total-Zn)

The values of total-Zn content enable better depiction of Zn accumulation in the soil. The total-Zn content of soils depends on the parent material from which the soils have been

developed. It is considered to be poor indicator of the zinc supplying capacity of soils for long term management of cropping system. The total-Zn content was found to be the highest among all the fractions and maximum content of total-Zn ($137.33 \text{ mg kg}^{-1}$) was observed in the treatment receiving soil application of Zn @ 7.5 kg ha^{-1} and the lowest concentration ($113.45 \text{ mg kg}^{-1}$) was recorded in the control. Effect of graded application of zinc fertilizer on different zinc fractions (Table 5) indicated that out of Total-Zn, majority of zinc was distributed in the Res-Zn fraction followed by ASB-Zn, CSB-Zn and Comp-Zn while WSEx-Zn occupied a very small portion ($< 1\%$) of the total-Zn pool in soil. Behera *et al.* [25] observed that among different zinc fractions in soil, Res-Zn was the dominant fraction of the total-Zn under maize-wheat cropping system. Jyothi *et al.* [26] reported that the contribution of Res-Zn was the maximum followed by CSB-Zn and ASB-Zn fractions, while, the contribution of WSEx-Zn fraction was only 1-2 per cent.

Data presented in Table 3, 4 and 5 divulged that application of zinc @ 7.5 kg ha^{-1} recorded significantly higher values of different fractions of zinc as compared to the control. Continuous addition of zinc at higher rates in this treatment and conversion of the added zinc resulted in transformation of zinc into these fractions. Vishvakarma [27] also reported an increase in the concentration of WSEx-Zn, organically bound-Zn and complexed-Zn fractions in soil with increase in the zinc levels. The control plots recorded lower values of these fractions probably due to non application of zinc fertilizer externally and continuous uptake by the crop.

Thus, it is evident that zinc fertilization significantly influenced the distribution of different zinc fractions in soil. At harvest, different zinc fractions were found in the order: water soluble plus exchangeable-Zn (WSEx-Zn) $<$ complexed-Zn (Comp-Zn) $<$ crystalline sesquioxide bound-Zn (CSB-Zn) $<$ amorphous sesquioxide bound-Zn (ASB-Zn) $<$ residual-Zn (Res-Zn) fractions. These results were in accordance with the findings of Preetha and Stalin [28], Soltani *et al.* [29] and Spelbar *et al.* [21].

Correlation among the different zinc fractions (Table 7) revealed that all the fractions of zinc were significantly and positively correlated with each other. Results showed significant positive correlation among WSEx-Zn, Comp-Zn, CSB-Zn, ASB-Zn, Res-Zn and Total-Zn fractions indicating existence of dynamic equilibrium of zinc in soil.

Table 6. Percent (%) contribution of different Zn fractions to total-Zn in soil

Treatments	WSEx-Zn	Comp-Zn	ASB-Zn	CSB-Zn	Res-Zn
------------	---------	---------	--------	--------	--------

T₁	0.26	1.75	2.82	2.51	92.66
T₂	0.40	1.87	2.60	2.43	92.70
T₃	0.44	1.93	2.86	2.61	92.14
T₄	0.60	2.16	3.00	2.77	91.49
T₅	0.29	1.57	2.46	2.09	93.59
T₆	0.38	1.57	2.34	2.10	93.61
T₇	0.33	1.70	2.61	2.42	92.93
T₈	0.46	1.89	2.76	2.46	92.42
T₉	0.50	1.95	2.76	2.61	92.19
T₁₀	0.64	1.66	2.53	2.36	92.79
T₁₁	0.74	1.88	2.74	2.47	92.16
T₁₂	0.86	1.92	2.76	2.64	91.81

Table 7. Correlation coefficient (r) among zinc fractions and DTPA-Zn

	WSEx-Zn	Comp-Zn	ASB-Zn	CSB-Zn	Res-Zn	Total-Zn
Complex-Zn	0.594*					
ASB-Zn	0.587*	0.977**				
CSB-Zn	0.634*	0.982**	0.980**			
Residual-Zn	0.627*	0.979**	0.975**	0.973**		
Total-Zn	0.635*	0.982**	0.979**	0.981**	0.997**	
DTPA-Zn	0.942**	0.656*	0.634*	0.672*	0.673*	0.680*

* Correlation is significant at the 0.05 level (2-tailed).

** Correlation is significant at the 0.01 level (2-tailed).

Path analysis

A path analysis was carried out to understand the contributions of different zinc fractions towards the plant available form of Zn in soil i.e., DTPA-Zn (Fig. 1). The direct effects of the studied Zn fractions towards DTPA-Zn were 0.783 (WSEx-Zn), 0.661 (Comp-Zn), 0.003 (ASB-Zn), 0.802 (CSB-Zn), 4.145 (Res-Zn) and 5.244 (Total-Zn). However, the indirect effect of the studied Zn fractions followed an opposite trend, where, the contribution towards DTPA-Zn were 0.052, -0.554, -0.010, -0.901, -4.386 and -4.974 for WSEx-Zn, Comp-Zn, ASB-Zn, CSB-Zn, Res-Zn and Total-Zn, respectively. Likewise, the combined

direct and indirect effect of WSEx-Zn (0.834) was the highest towards DTPA-Zn followed by Total-Zn (0.269) and Comp-Zn (0.106). This indicated that the WSEx-Zn had the highest contribution towards the plant available fraction of Zn in soil.




Fig. 1. A path analysis diagram and coefficient factors affecting DTPA-Zn

Conclusions

The dynamics of the studied zinc fractions in maize maize cropping sequence was significantly influenced by the treatment combination of basal and foliar application of zinc during both the years under investigation. The distribution of different zinc fractions in soils at harvest was found in the following order: water soluble plus exchangeable-Zn < complexed-Zn < crystalline sesquioxide bound-Zn < amorphous sesquioxide bound-Zn < residual-Zn. WSEx-Zn fraction least contributed to total-Zn however, had the highest

contribution towards the plant available DTPA extractable-Zn in soil as divulged by path analysis. Foliar application with zinc nano particle significantly increased WSEx-Zn in soil. Furthermore, all the fractions of zinc were found to be significantly and positively correlated with each other indicating existence of dynamic equilibrium of zinc in soil.

References:

- [1] P. Takkar and M.S. Mann, Evaluation of analytical methods for estimating available zinc and response of maize to applied zinc in major soil series of Ludhiana, Punjab. *Agrochimica*, 19 (1975) 420-429.
- [2] T. Adhikari and R. Rattan, Distribution of zinc fractions in some major soils of India and the impact on nutrition of rice. *Commun. Soil Sci. Plant Anal.*, 38 (2007) 2779–2798.
- [3] A.S.P. Murthy, Zinc fractions in wetland rice soils and their availability to rice. *Soil Science*, 133 (1982) 150-154.
- [4] B. Rahmani, M.M. Tehrani, A. Khanmirzaei and K. Shahbazi, Cadmium fractions and its uptake by the wheat plant in some calcareous soils of Iran. *Int. J. Agric. Res. Review.*, 2 (2012) 461-466.
- [5] S. Ramzan, M.A. Bhat, N.A. Kirmani and R. Rasool, Fractionation of zinc and their association with soil properties in soils of Kashmir Himalayas. *Int. J. Agric. Soil Sci.*, 2 (2014) 132-142.
- [7] R. Bala, A. Kalia and S.G. Dhaliwal, Evaluation of efficacy of ZnO nanoparticles as remedial zinc nano fertilizer for rice. *J. Soil Sci. Plant Nutri.*, 1 (2019) 1-12.
- [8] Y. Verma, S.K. Singh, H.S. Jatav, V.D. Rajput and T. Minkina, Interaction of zinc oxide nanoparticles with soil: Insights into the chemical and biological properties. *Environmental and Geochemical Health*, (2021) [https://doi.org/10.1007/s10653-021-00929-8\(0123456789\)](https://doi.org/10.1007/s10653-021-00929-8(0123456789)).
- [6] C.A. Black, *Soil plant relationship*, 2nd edition. New York., USA, 1965, pp. 515-516.
- [9] A.M. Ghoneim, Effect of different methods of Zn application on rice growth, yield and nutrients dynamics in plant and soil. *J Agri Ecol Res Int.*, 6 (2016) 1-9.
- [10] F.J. Stevenson and M.S. Ardakani, Organic matter reactions involving micronutrients in soils. In: J.J. Mortvedt, P.M. Giordano and W.L. Lindsay (Eds.) *Micronutrients in Agriculture*, Soil Science Society of America, Madison, USA, 1972, pp. 79-114.
- [11] M. Schnitzer and S.I.M. Skinner, Organo-metallic interactions in soils: 5. Stability constants of Cu⁺⁺, Fe⁺⁺ and Zn⁺⁺ fulvic acid complexes. *Soil Science.*, 102 (1966) 361-365.
- [12] M. Schnitzer, In: *Soil Organic Matter*. Schnitzer, M. and Khan, S. U. (Eds.). Elsevier: Amsterdam 1978.

- [13]F.J. Stevenson, Organic matter-micronutrient reactions in soil. In: J.J. Mortvedt, F.R. Cox, L.M. Shuman and R.M. Welch (Eds.) *Micronutrients in Agriculture*. Soil Science Society of America, Madison, USA, 1991, pp. 145-186.
- [14]L. Kiekens, Zinc. In: *Heavy metals in soils*. B.J. Alloway, (Ed.). Blackie: Glasgow, UK, 1990, pp. 261-279.
- [15]D.M.D.I. Wijebandara, Studies on distribution and distribution of soil zinc and response of rice to nutrient in traditional and system of rice intensification (SRI) methods of cultivation. Ph.D. Thesis, University of Agricultural Science, Dharwad, 2007.
- [16]P. Veeranagappa, H.C. Prakasha, K.R. Ashoka, M.N. Venkatesha and M.B. Kumar, Effect of zinc enriched compost on soil chemical properties and nutrient availability. *An Asian J. Soil Sci.*, 6 (2011) 189-194.
- [17]K. Talukder, I. Ahmed, M. Islam, M. Islam, M. Asaduzzaman and M. Hossain, Incubation studies on exchangeable zinc for varying levels of added zinc under aerobic and anaerobic conditions in grey terrace soils, non calcareous floodplain soils and calcareous floodplain soils. *J. Sci. Foundation*, 9 (2013), 9-15.
- [18]H.K. Kumar, Chemistry and fertility zincs selected paddy soils of Tungabhadra command. M.Sc. Thesis, University of Agricultural Science, Dharwad (2007).
- [19]D.M.D.I. Wijebandara, G.S. Dasong and P.L. Patil, Zinc fractions and their relationship with soil properties in paddy growing soils of northern dry hill zones of Karnataka. *Journal of the Indian Society of Soil Science*, 59 (2011), 141-147.
- [20]G.G. Kandali, N.G. Barua, A. Basumatary, B.K. Medhi, and K. Kurmi, Zinc fractions and their availability to wetland rice in acidic soils of Assam. *J. Indian Soc. Soil Sci.*, 64 (2016).
- [21]E. Spalbar, A.K. Mondal, and A.P. Rai, Distribution of zinc fractions and its association with soil properties in some rice-wheat growing soils of Jammu Region, India. *Int. J. Current Microbiol. App. Sci.*, 6 (2017): 711-721.
- [22]E.A. Forbes, A.M. Posner and J.P. Quirk, The specific adsorption of divalent Cd, Co, Cu, Pb and Zn on goethite. *J. Soil Sci.*, 27 (1976) 154-166.
- [23]M.D.A. Bolland, A.M. Posner and J.P. Quirk, Zinc adsorption by goethite in the absence and presence of phosphate. *Aust. J. Soil Res.*, 15 (1977) 279-286.
- [23]K.K. Singh, C.S. Praharaj, A.K. Choudhary, N. Kumar and M.S. Venkatesh, Zinc response in pulses. *Indian J. Fert.*, 7 (2011) 118-126.
- [24]Priyanka, S.K. Sharma and R.H. Meena, Fractionation and distribution of zinc under integrated nutrient management system on maize-wheat cropping system in Typic Haplustepts. *J. Pharmaco. Phytochem.*, 6 (2017) 2301-2305.
- [25]K.S. Behera, D. Singh, B.S. Dwivedi, S. Singh, K. Kumar and D.S. Rana, Distribution of fractions of zinc and their contribution towards availability and plant uptake of zinc under long term maize and wheat cropping on Inceptisol. *Aust. J. Soil Res.*, 46 (2008) 83-89.

- [26]T.Y. Jyothi, Y.V. Shetty, M.D. Kumar and H.V. Rudramurthy, Distribution of zinc fractions in traditional arecanut garden soils of Karnataka. *Environ Ecol*, 28 (2010) 1229-1234.
- [27]D. Vishvakarma, Effect of zinc application with or without organic manure on transformation of in maize grown on vertisols. M.Sc. Thesis, Jawarharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, 2012.
- [28]P.S. Preetha and P. Stalin, Different forms of soil zinc - their relationship with selected soil properties and contribution towards plant availability and uptake in maize growing soils of Erode district, Tamil Nadu. *Indian J. Sci. Technol.*, 7 (2014) 1018-1025.
- [29]S.M. Sotlani, M.M. Hanafi, S.A. Wahid and S.M.S. Khardiah, Zinc fractions of tropical paddy soils and their relationship with selected soil properties. *Chemical Spec. Bioavail.*, 27 (2015): 53-61.

UNDER PEER REVIEW