

## Original Research Article

### Evaluation of IPM modules against major pest of okra (*Abelmoschus esculentus*-L.) in Saran District (Bihar), India

#### Abstract :

Experiments were carried out at ten locations of Saran district, Bihar during *kharif* (Autumn) season 2017 with okra crop cv 'Supper green'. The assessment revealed that module (M<sub>4</sub>) proved the most effective treatment against okra pest and ~~diseases~~ *i.e.* shoot and fruit borer, *Earias vittella* F., yellow vein ~~mosaemosaic~~ virus, *Bemisia tabaci* (Genn.) and red spider mites, *Tetranychus urticae* -Koch in which lowest incidence was recorded as compared to other IPM modules, farmer practices and control plot during 2017. Module (M<sub>3</sub>) was the next effective treatment against the major pest of okra crop. Significantly, maximum fruit yield was observed in module (M<sub>4</sub>) 210.10 q/ha. -This was significantly superior over all other treatments. The impact of various IPM modules showed that net return was also higher in module (M<sub>4</sub>) which was Rs.76430.00/ha. It is conducted that judicious use of neonicotinoids pesticides in modules III and IV.

**Keywords**:- IPM modules, Okra, shoot and fruit borer, yellow vein mosaic virus, red spider mites, neonicotinoids

#### INTRODUCTION:

Okra (*Abelmoschus esculentus*), also known as lady's finger or bhendi, belongs to family Malvaceae and is an important vegetable crop grown throughout the year in India. Besides India, it is grown in many tropical and subtropical parts of the world. India grows okra on about 4.52 lakh hectares with an annual production of 48.03 lakh tons and productivity of 10.61 t/ha. -The productivity of our country is low compared to other countries due to yield losses caused by insect pests, diseases and nematodes. The crop is ~~attaek~~ *attacks* by more than 72 insect pests; and the infestation of the crop starting from the seedling to the harvest stage [1] (~~Kedar et al., 2013~~). Amongst various insect-pests causing damage, ~~;~~ shoot and fruit borer, *Earias vittella* F. (Lepidoptera: Noctuidae), yellow vein mosaic virus (YVMV) transmitted by whitefly, *Bemisia tabaci* (Gen), and the red spider mites, *Tetranychus urticae* Koch are considered as the most ~~limiting-effective~~ factors [2, 3] (~~Rawat and Saha, 1973; Alagar and Srivasubramanian, 2006~~). The excessive and indiscriminate use of pesticides to control these pests has resulted in undesirable ecological changes [4] (~~Mahapatra and Gupta, 1998~~). In view of the above, evaluation and adoption of IPM module in holistic manner incorporating judicious use of newly introduced modern pesticides seem to be best alternative. Hence, the present

Formatted: Font: Italic

Formatted: Double strikethrough, Highlight

Formatted: Double strikethrough

Formatted: Double strikethrough, Highlight

investigation to assessment of the effectiveness of different IPM modules in Saran district of Bihar.

The author forgets to mention the aim of this study. Please revise.

#### **Materials and Methods:**

Field experiments were carried out at ten locations ~~of~~ near Krishi Vigyan Kendra, Manjhi Block of Disstrict Saran, Bihar, under Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar in 2017. The trials were laid out randomized block design with ten replications. Okra “Supper green” was sown in first week of July with spacing 45 x 30 cm. All the agronomical practices recommended to raise the good crops were followed. Observations were made on ten ~~leaves were~~ randomly selected leaves (yellow vein mosaic virus transmitted by whitefly, *Bemisia tabaci* (Gen) and red spider mite) at fortnightly intervals after 50 days of transplanting. Observation were also recorded by counting healthy fruit and damaged of shoot and fruits for shoot and fruit borer insect at each picking and their weight were noted. The weight of healthy and infested fruits was taken separately and ~~per cent~~percentage of infestation was ~~worked out~~calculated. ~~Per cent~~Percentage of infestation and its reduction due to various treatments were transformed to Arcsine before subjected to analysis of variance. The yield data in various IPM modules were also recorded. Finally the crop was harvested in last week of October.

**Comment [m1]:** It is well-known that okra are planted as seeds, but not as seedlings, please revise this sentence!

**Comment [m2]:** Why? The real values is enough. Please revise

The author did not mention the total areas for his experiments (4 modules, comprising 17 treatments) with 10 replicates. The descriptions of the experimental design needs to rewritten with more details.

IPM module I : Untreated

-IPM module II : Farmers practice

i) Spraying of imidacloprid 0.5 ml/L water once at 30 days after sowing.

ii) Application of quinalphos @ 2 ml/L water 2 to 3 time at 7-10 days interval started 65-70 days after sowing.

iii) Spraying of fenvalerate @ 1 ml/L water twice at 7 days interval commenced 85-90 days after sowing.

-IPM module III

i) Deep summer ploughing

ii) Seed treatment with imidacloprid 70% WS @ 10g /kg-Kg of seeds

- iii) Use of neem cake @ ~~250kg~~250Kg/ha before **transplanting**
- iv) Spraying of spinosad @ 0.4 ml/~~L~~L water 40 and 50 days after **transplanting**.
- v) Application of imidacloprid 0.5ml/~~L~~L water 70 days after **transplanting**.
- vi) Spraying of abamectin @ 1 ml/~~L~~L water between 60 and 70 days after **transplanting**.

#### IPM module IV

- i) Deep summer ploughing
- ii) Seed treatment with imidacloprid 70% WS @ 10g /~~kg~~Kg of seeds
- iii) Use of neem cake @ ~~250kg~~250Kg/ha before **transplanting**.
- iv) Hand picking and destruction of infested leaves, shoots and fruits of shoot and fruit borer and mite infested leaf at initial stage.
- v) Spraying of indoxacarb @ 0.4 ml/~~L~~L water 45 and 60 days after **transplanting**.
- vi) Application of thiomethoxam 1g/~~L~~L water 75 days after **transplanting**.
- vii) Spraying of hexythiazox (acaricides) @ 0.5 ml/~~L~~L water between 65 and 85 days after **transplanting**.

The author did not mention the total volume of water applied per Hectare/Feddan which carrying the treatment insecticide concentration

#### Results and Discussion:

The result revealed that impact of IPM modules as compared to farmers' practices and untreated plot are presented in Table 1 and 2 during 2017. The incidence of shoot and fruit borer varies from 5.85 ~~per-centpercent~~ to 12.24 ~~per-centpercent~~ during 2017 in vegetative stage. The module (M<sub>4</sub>) showed lowest shoot damage *i. e.* 5.85 ~~per-centpercent~~ followed by module (M<sub>3</sub>) 7.02 ~~per-centpercent~~ shoot damage was recorded with compared to farmer's practices module (M<sub>3</sub>) 9.36 ~~per-centpercent~~ and untreated plot module (M<sub>1</sub>) 12.24 ~~per-centpercent~~. Module (M<sub>4</sub>) was observed significantly superior to module (M<sub>2</sub>) and module (M<sub>1</sub>). Similarly, fruit damage was higher in IPM modules (M<sub>1</sub>) 18.12 ~~per-centpercent~~ and modules (M<sub>2</sub>) 14.86 ~~per-centpercent~~ with compared to modules (M<sub>3</sub>) 9.87 ~~per-centpercent~~ and modules (M<sub>4</sub>) 7.25 ~~per-centpercent~~, respectively. The study revealed that module (M<sub>4</sub>) was significantly superior to module (M<sub>2</sub>) and module (M<sub>1</sub>).

Almost same trend was recorded in ~~per-centpercent~~ incidence of yellow vein mosaic virus and red spider mites. The incidence of yellow vein mosaic virus was higher as compared to shoot

and fruit borer damage which varies from 14.04 ~~per cent~~ module (M<sub>4</sub>) to 24.42 ~~per cent~~ in module (M<sub>1</sub>), but red spider mites was lowest in module (M<sub>4</sub>) which was only 1.89 ~~per cent~~ and highest in module (M<sub>1</sub>) i. e. 9.05 ~~per cent~~. The data recorded on incidence of shoot and fruit borer, yellow vein mosaic virus and red spider mites during field trials in various IPM modules recorded that module (M<sub>4</sub>) significantly reduced all the pest and demonstrated its superiority.

The yield data, gross return cost of cultivation, net return and benefit cost ratio due to adoption of different IPM modules were also worked out. The fruit yield various from 132.50 q/ha to 210.10 q/ha in during 2017. The maximum mean fruit yield was found in module (M<sub>4</sub>) i.e. 210.10 q/ha, which was significantly superior over all IPM modules. The mean gross return various from 99375 Rs/ha to 157575 Rs/ha in various IPM modules. However mean cost of cultivation was higher in module (M<sub>4</sub>) which was 81145 Rs/ha followed by module (M<sub>3</sub>) 79825 Rs/ha. Though cost of cultivation was higher in module (M<sub>4</sub>) but net return was maximum is 76430 Rs./ha in comparison with module (M<sub>1</sub>) i.e. 39090 Rs/ha and mean cost benefit ratio was also found maximum in module (M<sub>4</sub>) i.e. 0.94 followed by module (M<sub>3</sub>) 0.90, module (M<sub>2</sub>) 0.85 and module (M<sub>1</sub>) 0.65, respectively.

In general, the incidence of shoot and fruit borer, yellow vein mosaic virus and red spider mites was minimum and fruit yield was significantly higher in module (M<sub>4</sub>). It might be because of the influence of hand picking and destruction of infested leaves, shoots and fruits, seed treatment with imidacloprid, application of indoxacarb, thiomethoxam, hexythiazox, deep summer ploughing and use of neem cake @ 250 kg/ha before sowing. The results showed that application of neonicotinoids or chloronicotinyl insecticides in module (M<sub>1</sub>) along with other IPM strategies gave effective management from shoot and fruit borer, yellow vein mosaic virus and red spider mites and received higher yield as compared to other insecticides like quinalphos and fenvalerate.

The study conducted by Kodandaram *et al.*, [5] ~~(2010)~~ suggests that because of the relatively low risk to non-target organism and environment, high target specificity and their versatility in application methods, neonicotinoids are more useful Preetha and Nadarajam [6], ~~(2007)~~, Singh *et al.* [7] ~~(2008)~~ at Jaipur, Rajasthan; and Singh *et al.* [8], ~~(2010)~~ revealed that imidacloprid and indoxacarb treated okra crop showed maximum reduction in incidence of shoot and fruit borer and yellow vein mosaic virus and received higher yield as compared to other

insecticides like quinalphos and fenvalerate. Satpathy *et al.* [9], (2019) suggest that a good control of okra pest for seed treatment and seedling root dip is 0.02-0.03 per cent percent solution of Imidacloprid. Campos and Omoto [10] (2002) reported that widely used in controlling many phytophagous mites in vegetable crops.

### References

[example] Ravisankar N, Chandrasekaran B, Raja R, Din M, Chaudhuri SG. Influence of integrated weed - management practices on productivity and profitability of wet - seeded rice (*Oryza sativa*). Indian Journal of Agronomy. 2008; 53(1): 57-61.

All references needs to edit as the above example, as journal instructions

- 1- Kedar SC, Kumaranag KM, Nitin T. Integrated pest management in okra. 2013. [www.krishisewa.com](http://www.krishisewa.com).
- 2- Rawat RR, Saha HR. Estimation of losses in growth and yield of okra due to *Empoasca devastans* Dist. And *Earias* sp. *Indian Journal of Entomology*, 1973; **35** (2): 14-20.
- 3- Alagar, M. And Sivasubramanian. 2006. Field efficacy of botanicals insecticides and their combination against major pests of okra, *Abelmoschus esculentus* (L.) Moench. *Indian Journal of Entomology*, 68 (8): 369-374.
- 4- Mahapatro, G.K. and Gupta, G. P. 1998. Pesticide induced resurgence. *Pestology*, 22 (12): 14-20.
- 5- Kodandaram, M. H., Rai, A. B. and Halder Jaydeep. 2010. Noval insecticides for management of insects pest in vegetable crops: A Review. *Vegetable Science*, 37 (20):109-123.
- 6- Preetha, G. and Nadarajan, L. 2007. Validation of IPM modules against sucking pests of okra in Karaikal. *Indian Journal of Entomology*, **69** (3): 201-214.
- 7- Singh, Swaroop, Choudhary, D.P., Sharma, H.C., Mahla, R.S., Mathur, Y.S. and Ahuja, D.B. 2008. Effect of insecticidal molecules against jassid and shoot and fruit borer in okra. *Indian Journal of Entomology*, **70** (3): 197-199.
- 8- Singh, S.S., Rai, M.K. and Singh, V.B. 2010. Bio-efficacy of bio rational insecticides and Bt. Based bio-insecticides against shoot and fruit borer in okra. *Vegetable Science*, 37 (2):171-174.

- 9- Satpathy S., Rai A.B., Shivalingaswamy T.M., Kodandram M.H. and Kumar Akhilesh. 2010. Effect of novel biorational molecules - polymer combination as seed treatments on jassid population and seed vigour in okra. National Conference on Production of quality seeds planting material, health management in horticultural crops”, held on March 11-14, 2010 at New Delhi, pp. 84.
- 10- Campos, F. J. and Omoto, C. 2002. Resistance to hexythiazox in *Brevipalpus phoenicis* (Acari: Tenuipalpidae) from Brazilian citrus. *Experimental and Applied Acarology*, 26 : 243-251.

### References

- ~~Alagar, M. And Sivasubramanian. 2006. Field efficacy of botanicals insecticides and their combination against major pests of okra, *Abelmoschus esculentus* (L.) Moench. *Indian Journal of Entomology*, 68 (8): 369-374.~~
- ~~Campos, F. J. and Omoto, C. 2002. Resistance to hexythiazox in *Brevipalpus phoenicis* (Acari: Tenuipalpidae) from Brazilian citrus. *Experimental and Applied Acarology*, 26 : 243-251.~~
- ~~Kodandaram, M. H., Rai, A. B. and Halder Jaydeep. 2010. Novel insecticides for management of insects pest in vegetable crops: A Review. *Vegetable Science*, 37 (20):109-123.~~
- ~~Mahapatro, G.K. and Gupta, G. P. 1998. Pesticide induced resurgence. *Pestology*, 22 (12): 14-20.~~
- ~~Preetha, G. and Nadarajan, L. 2007. Validation of IPM modules against sucking pests of okra in Karaikal. *Indian Journal of Entomology*, 69 (3): 201-214.~~
- ~~Rawat, R. R. And Saha, H.R. 1973. Estimation of losses in growth and yield of okra due to *Empoasca devastans* Dist. And *Earias* sp. *Indian Journal of Entomology*, 35 (2): 14-20.~~
- ~~Santosh Kumar C. Kedar, Kumaranag K.M\* and Nitin Thodsare. 2013. Integrated pest management in okra. [www.krishisewa.com](http://www.krishisewa.com).~~
- ~~Satpathy S., Rai A.B., Shivalingaswamy T.M., Kodandram M.H. and Kumar Akhilesh. 2010. Effect of novel biorational molecules - polymer combination as seed treatments on jassid population and seed vigour in okra. National Conference on Production of quality seeds planting material, health management in horticultural crops”, held on March 11-14, 2010 at New Delhi, pp. 84.~~
- ~~Singh, S.S., Rai, M.K. and Singh, V.B. 2010. Bio efficacy of bio rational insecticides and Bt. Based bio insecticides against shoot and fruit borer in okra. *Vegetable Science*, 37 (2):171-174.~~
- ~~Singh, Swaroop, Choudhary, D.P., Sharma, H.C., Mahla, R.S., Mathur, Y.S. and Ahuja, D.B. 2008. Effect of insecticidal molecules against jassid and shoot and fruit borer in okra. *Indian Journal of Entomology*, 70 (3): 197-199.~~

Formatted: Double strikethrough

Tables must be comprises the 17<sup>th</sup> treatments within the 4 modules with clear statistical comparison

**Table 1: Incidence of major pests in different IPM modules conducted in okra crop during kharif 2017**

Treatments	Shoot & fruit borer (% infestation) on shoot	Shoot & fruit borer (% infestation) on fruits	Yellow vein Mosaic Virus (% infestation)	Red spider mites (% infestation)	Yield (q/ha)
IPM modules I (Untreated)	12.24 (20.48)	18.12 (25.19)	24.42 (29.61)	9.05 (17.51)	132.5
IPM modules II (Farmers' practices)	9.36 (17.81)	14.86 (22.67)	19.23 (26.01)	6.35 (14.60)	172.7
IPM modules III	7.02 (15.36)	9.87 (18.31)	16.25 (23.71)	3.35 (10.55)	202.4
IPM modules IV	5.85 (14.00)	7.25 (15.62)	14.04 (22.01)	1.89 (7.90)	210.1
SEM±	0.26	0.24	0.35	0.15	1.78
CD-5%	0.90	0.84	1.19	0.53	6.23

Figures in parenthesis are the Arc sine  $\sqrt{\text{percentage}}$

**Table 2: Economic viability of different IPM modules in okra crop during kharif 2017**

Treatments	Gross return (₹/ha)	Cost of cultivation (₹/ha)	Net return (₹/ha)	Benefit Cost ratio
IPM modules I (Untreated)	99375	60285	39090	0.65
IPM modules II (Farmers' practices)	129525	70175	59350	0.85
IPM modules III	151800	79825	71975	0.90
IPM modules IV	157575	81145	76430	0.94

Net : The sale price of fruit was considered as Rs. 750 per quintal