

Original Research Article

Character association studies for yield and quality traits in rice (*Oryza sativa* L.)

ABSTRACT

Background: The production of rice needs to grow to fulfill the requirement of the rapidly growing population. **Objective and Methods:** The field experiment was conducted to study character association among yield and its attributes among 58 rice genotypes at Indira Gandhi Krishi Vishwavidyalaya, College of Agriculture, Raipur (C.G.) in India during *Kharif* 2020 by following a randomized complete block design with two replications. **Results:** we found that Traits like panicle weight per plant, fertile spikelets per panicle, harvest index, and biological yield per plant are having a high direct effect and significant correlation with grain yield per plant, and traits like milled grain length-width ratio, hulling %, and alkali spreading value are having a high direct effect and significant correlation with head rice recovery. **Conclusions:** Direct selection based on these traits for improvement of rice grain yield and grain quality can be effective.

KEYWORDS: Rice, Correlation, path coefficient, and yield.

INTRODUCTION

Rice (*Oryza sativa* L.) is the most extensively grown cereal worldwide. since it is the principal source of calories It supplies most of the requirements of starch, protein, and micronutrient for over half of the world's population. Insufficient availability of grain yield is also typically associated with poverty in many regions of the globe. Knowledge of the inter-relationship between characters and grain yield is essential for the improvement of crop production. A correlation coefficient is a statistical measure that is used to find out the degree and direction of the relationship between two or more variables working at the same time. It also helps to know how the improvement in one character will bring simultaneous change in other characters. While path analysis splits

the correlation coefficient into direct and indirect effects which specify the relative contribution of each character (Shanthala *et al.*, 2004). A study about the contribution of the direct and indirect effects of each character towards yield or quality traits could be an added advantage in aiding the selection process (Babu *et al.*, 2012). In the present paper, the correlation and path coefficients have been evaluated to estimate the contribution of characters on grain yield in rice.

MATERIALS AND METHODS

Experimental Site

The field experiment was conducted at Indira Gandhi Krishi Vishwavidyalaya, College of Agriculture, Raipur (C.G.) in India, during *the Kharif season* 2020. Geographically, this Indian state is located between 17°14' N and 24°45' N latitude and longitude of 79°16' E. The experiment was laid out by following a randomized complete block design with two replications. All the standard agronomic practices were adopted during the crop season.

Plant Materials

A total of 58 accessions including 06 checks *i.e.*, Dubraj Sel-1, CG Zinc Rice 1, CG Zinc Rice 2, Zinco Rice MS, Rajeshwari, and Protezin were used in the experiment. All the genotypes were accessed from the Department of Genetics and Plant Breeding, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.). Randomly five plants were selected from each replication for further studies.

METHOD:

Correlation coefficient analysis

The correlation coefficient was determined to illustrate the mutual relation between seventeen agro-morphological traits like days to 50% flowering, leaf length (cm), leaf width (cm), leaf area, plant height (cm), panicle length (cm), total tillers per plant, effective tillers per plant, fertile spikelets per panicle, sterile spikelets per panicle, total spikelets per panicle, spikelet fertility percentage, biological yield per plant (g), panicle weight per plant, harvest index, hundred seed weight (g) and grain yield per plant (g) in all the genotypes. The correlation coefficients at phenotypic and genotypic standards were evaluated from the analysis of variance and covariance. The linear relationship between two variables was estimated by using Karl Pearson's coefficient of

correlation which is based on the variance and covariance of the variables. The significance of the correlation coefficient was tested by comparing the observed value of the correlation coefficient with tabulated values for n-2 degree of freedom. If the observed value is more than the table value, the correlation coefficient is significant.

Path coefficient analysis

Path coefficient analysis promotes the partition of the correlation into the measures of direct and indirect effects. The genetic composition of yield must be dealt with the genetic improvement of all other characters, affecting it directly or indirectly, the phenotypic and genotypic correlation coefficients were significantly partitioned into direct and indirect effects with the help of path coefficient analysis, as suggested by Wright, 1921 and elaborated by Dewey and Lu, 1959. The path coefficient was assessed by clarifying and observing a series of simultaneous equations, specifying the basic relationship between correlation and path coefficients.

$$r_{iy} = P_{iy} + r_{i1}P_{1y} + r_{i2}P_{2y} + \dots + r_{i(i-1)}P_{iy}$$

$$i = 1, 2, 3, \dots, n$$

Where i is the number of independent characters r_{1y} to r_{iy} stands for the coefficient of correlation between independent variables (1...i) with dependent character y; r_{i1} to $r_{i(i-1)}$ stands for the coefficient of correlation among all feasible consolidations of caused factors (independent variables); P_{1y} to P_{iy} stands for the direct effects of independent variables (1...i) on the dependent variable y.

List 1: Results of path coefficient analysis are interpreted as per the scale suggested by Lenka and Mishra (1973)

S. No.	Value of direct/indirect effect	Rate/Scale
1	0.00 to 0.09	Negligible
2	0.10 to 0.19	Low
3	0.20 to 0.29	Moderate
4	0.30 to 0.99	High
5	>1.00	Very High

STATISTICAL ANALYSIS:

Statistical analysis for association study was evaluated through the standard software (Windostat version 9.2 from Indostat services, and MS-Excel-2010, etc.)

where Replicated data of each trait were used for the calculation of genotypic and phenotypic correlation coefficients and path correlation effect.

RESULTS AND DISCUSSION:

Association analysis is an important approach in a breeding program. It gives an idea about the relationship among the various characters and determines the component characters, on which selection can rely for genetic improvement in the grain yield. It helps to know how the improvement in one character will bring simultaneous change in other characters. Phenotypic and genotypic correlations for various yield and quality attributing traits were estimated.

Correlation study based on agro-morphological traits

In the present study grain yield per plant showed a positive and significant correlation at the phenotypic level with panicle weight per plant (0.817), harvest index (0.802), biological yield per plant (0.331), fertile spikelet per panicle (0.290), effective tillers per plant (0.261), total spikelet per panicle (0.20), total tillers per plant (0.194) and spikelet fertility % (0.182). where hundred seed weight (0.135) and days to 50% flowering (0.116) showed a positive correlation with rice grain yield per plant. Whereas it showed a significant negative correlation with leaf length (-0.094), leaf width (-0.052), leaf area (-0.111), plant height (-0.027), panicle length (cm) (-0.043), and sterile spikelet per panicle (-0.068), respectively. Singh *et al.*, (2018) and Bagudam *et al.*, (2018) also revealed a positive and highly significant correlation of yield with harvest index, days to 50% flowering, productive tillers per plant, tillers per plant, and biological yield per plant to emerge as the most important associates of grain yield in rice.

Correlation study based on grain quality traits

The phenotypic and genotypic correlations for grain quality attributing traits were estimated. The grain quality traits viz., milling% (0.336), hulling % (0.287), iron content (0.295), alkali spreading value (0.267), elongation ratio (0.262), milled grain L/B (0.245), zinc (0.238) and decorticated grain L/B (0.218) were having positive and significant correlation at phenotypic level with head rice recovery. Whereas head rice recovery showed a significant negative correlation with grain length (-0.260), decorticated grain length (-0.242), milled grain length (-0.232), and cooked grain length

(-0.195). Thus, correlation analysis enables breeders to comprehend the shared character traits that might serve as the basis for selection for genetic improvement. In addition to this, understanding the type and extent of genetic diversity is crucial for enacting genetic improvement and for employing the appropriate selection techniques.

Path coefficient analysis based on agro morphological traits

Path coefficient analysis splits the correlation coefficient into the measure of direct and indirect effects of a set of independent variables on the dependent variable. If the correlation between yield and character is due to the direct effects of character, it reflects the true relationship between them. In the present study, a genotypic path coefficient was utilized for the study where grain yield per plant was considered as the dependent variable for agro-morphological traits and the head rice recovery was the dependent variable for grain quality traits, and the rest all the traits were considered as independent while correlation coefficient between grain yield and its attributing characters were partitioned into direct and effects in path coefficient analysis. Further effects were scaled based on the scale of Lenka and Mishra (1973) for path analysis.

Path analysis revealed that the very high positive direct contribution on grain yield per plant was expressed by fertile spikelets per panicle (1.041), while harvest index (0.765), biological yield per plant (0.527) and sterile spikelets per panicle (0.438) showed high positive direct effect whereas panicle weight per plant (0.288) showed a moderate positive direct effect on yield. Singh *et al.*, (2018) and Sivasankar *et al.*, (2018) also reported that the biological yield per plant, harvest index, panicle bearing tillers per plant, panicle length, spikelet per panicle, biological yield, harvest index is the benefitted and efficient for selection for the enhancement of rice productivity.

Linking correlation with path analysis gives a better picture for selection compared to correlation alone, by employing this concept in this study it was observed that the trait viz; panicle weight per plant, fertile spikelets per panicle, harvest index, biological yield per plant showed high direct effect and positive significant correlation on grain yield per plant. However, grain quality traits like milled grain L/B, hulling %, and alkali spreading value are having a high direct effect and significant positive correlation with head rice recovery. This shows their true and potent relationship and direct selection in those traits (Singh and Narayana, 2009) will directly lead to high

yield and good head rice recovery. Similar results were reported by Hossain *et al.*, (2018); Parimala *et al.*, (2020); and Samudin *et al.*, (2021).

CONCLUSION

In conclusion, character association analysis revealed that direct selection can be possible for the traits which are having a very high direct effect and strong positive correlation with grain yield per plant and head rice recovery for improvement in yield and grain quality in rice respectively.

CONFLICT OF INTEREST STATEMENT

The author declares that there's no conflict-of-interest present in this study, its respondents, and its researcher. No Funding was provided for this article. However, the declaration of no conflict of interest still stands since the publication of research is a core requirement in the author's doctoral degree.

REFERENCES:

- Babu, V.R., Shreya, K., Dangi, K.S., Usharani, G. and Nagesh, P. 2012. Genetic variability studies for qualitative and quantitative traits in popular rice (*Oryza sativa* L.) hybrids of India. *Int. J. Scient. Res. Publ.*, 2(6): 1-5.
- Bagudam, R., Eswari, K.B., Badri, J. and Rao, R.P. 2018. Correlation and Path Analysis for Yield and its Component Traits in NPT Core Set of Rice (*Oryza sativa* L.). *Int.J.Curr.Microbiol.App.Sci.*, 7(9): 97-108.
- Dewey, J.R. and Lu, K.H. 1959. Correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron J.*, 51: 515-518.
- Hossain, S., Salim, M., Azam, M. G. and Noman, S. 2018. Variability, correlation and path analysis in drought tolerant rice (*Oryza sativa* L.). *Journal of Bioscience and Agriculture Research*, (02):1521-1530.
- Lenka, D. and Mishra, B. 1973. Path coefficient analysis of yield in rice varieties. *Indian J. Agric. Sci.*, 43: 376-379
- Parimala, K., Surender, Raju, C.H., HariPrasad, A.S., Kumar, S.S. and Reddy, S.N. 2020. Studies on genetic parameters, correlation, and path analysis in rice (*Oryza sativa* L.). *J. pharmacogn. phytochem.*, 9(1): 414-417.

Samudin, S., Maemunah, M., Ete, A., Mustakim, U., Yusran and Effendy. 2021. Determination of selection criteria to increase local upland rice yields. PCBMB, 22(11-12):165-176.

Singh, P. and Narayanan, S.S. 2009. Biometrical Techniques in Plant Breeding. 4th Edition Kalyani Publishers, New Delhi; 56-65.

Shanthala, J., Latha, J. and Shailaja, H. 2004. Path coefficient analysis for grain yield with yield components in hybrid rice. Environ. Ecol., 22(4): 734-736.

Sivasankar, R., Suresh, B.G., Ashish, S. and Sudheer, T.R. 2018. Correlation and Path Coefficient Analysis in Elite Germplasm of Rice (*Oryza sativa* L.). Int. J. Curr. Microbiol. App. Sci. 7(07): 3454-3459.

Agashe Nehatai Wamanrao, Vinod Kumar and Dronkumar Meshram. 2020. Correlation and Path Coefficient Analysis of Grain Yield and its Growth Components in Soybean (*Glycine max.* L.). Int.J.Curr.Microbiol.App.Sci. 9(03): 2445-2451.

UNDER PEER REVIEW

Table no.1 Phenotypic (p) and genotypic (g) correlation coefficients of seventeen agro-morphological traits in rice.

Traits		DFF	LL	LW	LA	PH	PL	TTPP	ETPP	FSP	SSPP	TSPP	SF	BYPP	PWPP	HI	HSW	
Days to fifty % flowering	g	0.529**	0.108	0.414	0.392	0.201	-0.38	-0.325	0.307	0.155	0.338	-0.04	0.27	0.239	-0.07	-0.24	-0.009	
	p	0.478**	0.102	0.375**	0.383**	0.159	-0.29**	-0.238*	0.288**	0.144	0.319**	-0.038	0.274**	0.228*	-0.063	-0.232*	0.116	
Leaf length	g		0.305**	0.844	0.428	0.427	-0.43	-0.397	0.217	-0.01	0.194	0.044	0.22	0.002	-0.27	-0.13	0.27	
	p		0.321**	0.852**	0.403**	0.344**	-0.335**	-0.305**	0.211*	-0.011	0.181	0.068	0.207*	0.01	-0.217*	-0.122	0.14	
Leaf width	g			0.764**	0.203	0.19	-0.23	-0.26	-0.31	-0.09	-0.305	-0.03	0.09	-0.02	-0.15	0.254	0.331	
	p			0.763**	0.190*	0.163	-0.221*	-0.228*	-0.281**	-0.084	-0.285**	-0.012	0.098	-0.021	-0.135	0.239**	0.154	
Leaf area	g				0.4**	0.374	-0.42	-0.412	-0.05	-0.07	-0.063	0.018	0.18	-0.04	-0.28	0.039	0.358	
	p				0.373**	0.306**	-0.347**	-0.329**	-0.027	-0.067	-0.054	0.044	0.177	-0.026	-0.230*	0.038	0.181	
Plant height	g					0.583**	-0.46	-0.507	0.217	-0.07	0.16	0.134	0.47	0.089	-0.37	0.075	0.36	
	p					0.486**	-0.366**	-0.394**	0.209*	-0.064	0.156	0.121	0.443**	0.09	-0.324**	0.073	0.195*	
Panicle length	g						-0.35**	-0.379	0.215	0.081	0.228	0.008	0.53	0.091	-0.32	-0.02	0.22	
	p						-0.279**	-0.281**	0.164	0.058	0.17	0.013	0.447**	0.08	-0.313**	-0.026	0.154	
Total tiller per plant	g							0.88**	0.008	-0.15	-0.084	0.183	0	0.202	0.197	-0.05	-0.158	
	p							0.897**	-0.005	-0.13	-0.073	0.134	0.03	0.221*	0.188*	-0.056	-0.025	
Effective tiller per plant	g								0.056	-0.12	-0.029	0.165	0.07	0.285	0.196	-0.11	-0.204	
	p								0.025	-0.101	-0.035	0.116	0.102	0.283**	0.204*	-0.104	-0.086	
Fertile spikelet per panicle	g									0.064	0.865	0.386	0.2	0.323	0.185	-0.59	0.091	
	p									0.053	0.860**	0.386**	0.186*	0.308**	0.172	-0.569**	0.045	
Sterile spikelet per panicle	g										0.552**	-0.84	0.01	0.076	-0.08	-0.21	-0.166	
	p										0.547**	-0.830**	0.013	0.068	-0.072	-0.204*	-0.042	
Total spikelets per panicle	g											-0.11	0.18	0.308	0.099	-0.59	-0.009	
	p											-0.119	0.174	0.291**	0.092	-0.569**	0.022	
Spikelet fertility	g												0.02	0.058	0.179	-0.1	0.22	
	p												0.017	0.06	0.163	-0.098	0.063	
Biological yield per plant	g													0.413**	-0.31	0.032	0.108	
	p													0.401**	-0.280**	0.02	0.011	
Panicle weight per plant	g														0.589**	0.226	-0.038	
	p														0.574**	0.202*	-0.058	
Harvest index	g															0.137	-0.114	
	p															0.119	-0.13	
Hundred seed weight	g																	-0.049
	p																	-0.019
Grain yield per plant	g	0.1199	-0.13	-0.065	-0.149	-0.04	0.003	0.162	0.221	0.32	-0.079	0.221	0.2	0.336	0.856	0.784	0.159	
	p	0.116	-0.094	-0.052	-0.111	-0.027	-0.043	0.194	0.261	0.29	-0.068	0.2	0.182	0.331	0.817	0.802	0.135	

*, ** significant at 5% and 1% probability levels

Table no. 2 Phenotypic (p) and genotypic (g) correlation coefficients for twenty-grain quality traits of rice.

Traits		M%	GL	GW	DGL	DGW	DGL/B	MGL	MGW	MGL/B	CGL	CGW	CL/CB	ER	ASV	GC	AC%	Iron	Zinc
Hulling %	g	0.86**	0.59**	-0.65**	0.64**	-0.56**	-0.62**	0.63**	-0.65**	-0.63**	0.78**	-0.54**	0.84**	-0.61**	-0.68**	0.04 ^{NS}	0.28**	-0.52**	-0.67**
	p	0.71**	0.378**	-0.403**	0.366**	-0.357**	-0.392**	0.368**	-0.412**	-0.392**	0.442**	-0.302**	0.439**	-0.379**	-0.388**	0.049 ^{NS}	0.166 ^{NS}	-0.321**	-0.394**
Milling %	g		0.53**	-0.73**	0.61**	-0.62**	-0.71**	0.58**	-0.75**	-0.72**	0.78**	-0.49**	0.76**	-0.68**	-0.79**	0.21*	0.45**	-0.64**	-0.74**
	p		0.378**	-0.512**	0.403**	-0.420**	-0.511**	0.393**	-0.469**	-0.521**	0.486**	-0.308**	0.446**	-0.488**	-0.508**	0.160 ^{NS}	0.281**	-0.440**	-0.445**
Grain length	g			-0.702**	0.955**	-0.636**	-0.627**	0.959**	-0.713**	-0.704**	0.917**	-0.403**	0.830**	-0.823**	-0.822**	0.312**	0.425**	-0.833**	-0.842**
	p			-0.699**	0.947**	-0.613**	-0.619**	0.942**	-0.677**	-0.696**	0.882**	-0.344**	0.753**	-0.815**	-0.787**	0.305**	0.404**	-0.693**	-0.763**
Grain width	g				-0.741**	0.929**	0.846**	-0.74**	0.932**	0.895**	-0.818**	0.613**	-0.774**	0.885**	0.916**	-0.388**	-0.506**	0.741**	0.906**
	p				-0.736**	0.895**	0.835**	-0.73**	0.888**	0.884**	-0.784**	0.512**	-0.698**	0.879**	0.878**	-0.376**	-0.484**	0.619**	0.821**
Decorticated grain length	g					-0.694**	-0.609**	0.988**	-0.752**	-0.712**	0.943**	0.406**	0.849**	-0.849**	-0.839**	0.324**	0.462**	-0.795**	-0.857**
	p					-0.662**	-0.594**	0.975**	-0.713**	-0.699**	0.908**	-0.345**	0.772**	-0.840**	-0.799**	0.312**	0.436**	-0.661**	-0.775**
Decorticated grain width	g						0.690**	-0.68**	0.973**	0.779**	-0.771**	0.637**	-0.755**	0.841**	0.805**	-0.357**	-0.493**	0.595**	0.796**
	p						0.627**	-0.63**	0.938**	0.728**	-0.692**	0.501**	-0.635**	0.805**	0.744**	-0.324**	-0.458**	0.479**	0.692**
Deco. grain length/width	g							-0.61**	0.727**	0.983**	-0.722**	0.509**	-0.624**	0.854**	0.915**	-0.371**	-0.540**	0.860**	0.891**
	p							-0.60**	0.663**	0.971**	-0.689**	0.421**	-0.560**	0.841**	0.874**	-0.369**	-0.515**	0.713**	0.804**
Milled grain length	g								-0.750**	-0.705**	0.943**	-0.392**	0.837**	-0.850**	-0.834**	0.338**	0.415**	-0.786**	-0.866**
	p								-0.690**	-0.681**	0.895**	-0.351**	0.766**	-0.843**	-0.792**	0.323**	0.394**	-0.642**	-0.782**
Milled grain width	g									0.800**	-0.843**	0.616**	-0.815**	0.882**	0.837**	-0.370**	-0.452**	0.643**	0.818**
	p									0.722**	-0.756**	0.490**	-0.689**	0.833**	0.773**	-0.318**	-0.429**	0.542**	0.730**
Milled grain length/width	g										-0.791**	0.540**	-0.700**	0.915**	0.969**	-0.386**	-0.613**	0.901**	0.941**
	p										-0.762**	0.455**	-0.625**	0.897**	0.914**	-0.380**	-0.563**	0.735**	0.835**
Cooked grain length	g											-0.412**	0.886**	-0.878**	-0.904**	0.342**	0.514**	-0.867**	-0.888**
	p											-0.351**	0.848**	-0.816**	-0.826**	0.310**	0.453**	-0.687**	-0.795**
Cooked grain width	g												-0.733**	0.598**	0.552**	-0.188**	-0.247**	0.511**	0.555**
	p												-0.711**	0.509**	0.424**	-0.11 ^{NS}	-0.15 ^{NS}	0.373**	0.426**
Cooked grain length/width	g													-0.782**	-0.808**	0.272**	0.393**	-0.802**	-0.808**
	p													-0.685**	-0.690**	0.212**	0.327**	-0.611**	-0.685**
Elongation ratio	g														0.942**	-0.393**	-0.542**	0.862**	0.941**
	p														0.902**	-0.384**	-0.520**	0.717**	0.846**
Alkali spread. value	g															-0.408**	-0.569**	0.865**	0.942**
	p															-0.401**	-0.584**	0.774**	0.914**
Gel consistency	g																0.142 ^{NS}	-0.437**	-0.381**
	p																0.153 ^{NS}	-0.342**	-0.336**
Amylose content%	g																	-0.549**	-0.491**
	p																	-0.492**	-0.497**
Fe content	g																		0.831**
	p																		0.764**
Zn content	g																		
	p																		
Head rice recovery	g	0.287**	0.336**	-0.260**	0.146 ^{NS}	-0.242**	0.103 ^{NS}	0.218*	-0.232*	0.134 ^{NS}	0.245**	-0.195*	-0.037 ^{NS}	-0.115 ^{NS}	0.262**	0.267**	-0.096 ^{NS}	-0.181 ^{NS}	0.295**
	p	0.26**	-0.17 ^{NS}	-0.696**	0.369**	-0.612**	0.350**	0.606**	-0.634**	0.379**	0.685**	-0.454**	0.087 ^{NS}	-0.284**	0.724**	0.557**	-0.272**	-0.342**	0.658**

*, ** significant at 5% and 1% probability levels.

Table no.3 Path coefficient analysis for seventeen agro-morphological traits for grain yield in rice.

Characters	Days to fifty % flowering	Leaf length	Leaf width	Leaf area	Plant height	Panicle length	Total tiller per plant	Effective tiller per plant	Fertile spikelet per panicle	Sterile spikelet per panicle	Total spikelets per panicle	Spikelet fertility	Biological yield per plant	Panicle weight per plant	Harvest index	Hundred seed weight	Grain yield per plant	Partial R ²
Days to fifty % flowering	-0.012	-0.069	-0.003	0.038	-0.047	-0.006	0.019	0.025	0.320	0.068	-0.381	0.004	0.144	0.069	-0.054	-0.009	0.120	-0.001
Leaf length	-0.006	-0.131	-0.009	0.076	-0.052	-0.013	0.021	0.031	0.225	-0.004	-0.219	-0.004	0.117	0.001	-0.209	-0.005	-0.014	0.018
Leaf width	-0.001	-0.040	-0.030	0.069	-0.025	-0.006	0.012	0.020	-0.322	-0.040	0.344	0.003	0.047	-0.007	-0.112	0.010	-0.065	0.002
Leaf area	-0.005	-0.110	-0.023	0.091	-0.048	-0.011	0.021	0.032	-0.048	-0.032	0.071	-0.002	0.095	-0.010	-0.212	0.002	-0.149	-0.014
Plant height	-0.005	-0.056	-0.006	0.036	-0.121	-0.018	0.023	0.039	0.226	-0.029	-0.180	-0.013	0.248	0.026	-0.282	0.003	-0.056	0.004
Panicle length	-0.002	-0.056	-0.006	0.034	-0.070	-0.030	0.018	0.030	0.224	0.036	-0.257	-0.001	0.277	0.026	-0.246	-0.001	0.002	0.000
Total tiller per plant	0.005	0.056	0.007	-0.038	0.055	0.011	-0.050	-0.069	0.009	-0.068	0.095	-0.018	-0.006	0.058	0.151	-0.002	0.162	-0.008
Effective tiller per plant	0.004	0.052	0.008	-0.037	0.061	0.011	-0.044	-0.078	0.058	-0.053	0.033	-0.016	0.037	0.082	0.150	-0.004	0.221	-0.017
Fertile spikelet per panicle	-0.004	-0.028	0.009	-0.004	-0.026	-0.007	0.000	-0.004	1.041	0.028	-0.974	-0.037	0.105	0.093	0.141	-0.023	0.320	0.333
Sterile spikelet per panicle	-0.002	0.001	0.003	-0.007	0.008	-0.002	0.008	0.009	0.067	0.438	-0.622	0.080	0.005	0.022	-0.063	-0.008	-0.079	-0.035
Total spikelets per panicle	-0.004	-0.025	0.009	-0.006	-0.019	-0.007	0.004	0.002	0.900	0.242	-1.126	0.011	0.098	0.089	0.075	-0.023	0.221	-0.249
Spikelet fertility	0.001	-0.006	0.001	0.002	-0.016	0.000	-0.009	-0.013	0.401	-0.367	0.126	-0.096	0.011	0.017	0.137	-0.004	0.201	-0.019
Biological yield per plant	-0.003	-0.029	-0.003	0.016	-0.057	-0.016	0.001	-0.005	0.207	0.004	-0.208	-0.002	0.527	0.119	-0.235	0.001	0.336	0.177
Panicle weight per plant	-0.003	0.000	0.001	-0.003	-0.011	-0.003	-0.010	-0.022	0.336	0.033	-0.347	-0.006	0.218	0.288	0.451	0.009	0.856	0.247
Harvest index	0.001	0.036	0.004	-0.025	0.045	0.010	-0.010	-0.015	0.192	-0.036	-0.111	-0.017	-0.162	0.170	0.765	0.005	0.784	0.600
Hundred seed weight	0.003	0.018	-0.008	0.004	-0.009	0.001	0.003	0.008	-0.610	-0.094	0.660	0.010	0.017	0.065	0.105	0.038	0.159	0.006

(R²=0.9804, Residual effect = 0.1401) diagonal Bold values represent the direct effect

(DFF = days to 50% flowering, LL= leaf length, LW= leaf width, LA= leaf area, PH= plant height, PL= panicle length, TTPP= total tillers per plant, ETPP= effective tillers per plant, FSPP= fertile spikelets per plant, SSPP= sterile spikelets per panicle, TSPP= total spikelets per panicle, SF= spikelet fertility %, BYPP= biological yield per plant, PWPP= panicle weight per plant, HI= harvest index, HSW= head rice recovery, GYPP= grain yield per plant).

Table no.4 Path coefficient analysis for twenty-grain quality traits in rice.

Characters	Hulling %	Milling %	Grain Length	Grain Width	Grain length	Decoricated Grain width	Decoricated Grain L/B	Decoricated Length	Milled Grain Width	Milled L/B	Milled Grain Length	Cooked Grain Width	Cooked L/B	Ratio	Elongation Spreading value	Alkali	Gel Consistency	Amylose Con%	Fe (ppm)	Zn (ppm)
H%	1.437	-0.627	1.188	0.818	0.229	2.403	0.191	-2.779	-4.8	-3.6	3.09	0.69	-2.7	0.42	-0.496	0.013	-0.044	0.096	0.286	
M%	1.248	-0.723	1.068	0.925	0.219	2.683	0.219	-2.545	-5.52	-4.2	3.09	0.62	-2.438	0.47	-0.577	0.066	-0.069	0.119	0.314	
GL	0.851	-0.385	2.007	0.881	0.342	2.728	0.193	-4.196	-5.2	-4.1	3.61	0.51	-2.661	0.56	-0.599	0.095	-0.065	0.153	0.356	
GW	-0.937	0.533	-1.409	-1.255	-0.265	-3.981	-0.26	3.25	6.797	5.16	-3.23	-0.77	2.482	-0.6	0.667	-0.117	0.078	-0.136	-0.383	
DGL	0.922	-0.443	1.916	0.929	0.358	2.975	0.187	-4.326	-5.48	-4.1	3.72	0.51	-2.721	0.58	-0.611	0.098	-0.071	0.146	0.362	
DGW	-0.806	0.452	-1.277	-1.166	-0.248	-4.287	-0.212	2.982	7.099	4.49	-3.04	-0.8	2.421	-0.57	0.586	-0.108	0.076	-0.109	-0.336	
DGL/B	-0.894	0.515	-1.259	-1.061	-0.218	-2.959	-0.307	2.702	5.305	5.67	-2.85	-0.64	2.002	-0.58	0.666	-0.112	0.083	-0.158	-0.376	
MGL	0.913	-0.42	1.924	0.932	0.354	2.92	0.19	-4.377	-5.47	-4.1	3.72	0.49	-2.684	0.58	-0.607	0.102	-0.064	0.144	0.366	
MGW	-0.946	0.547	-1.431	-1.17	-0.269	-4.172	-0.223	3.283	7.295	4.61	-3.33	-0.78	2.614	-0.6	0.61	-0.112	0.069	-0.118	-0.345	
MGL/B	-0.908	0.526	-1.413	-1.123	-0.255	-3.34	-0.302	3.084	5.839	5.77	-3.12	-0.68	2.244	-0.62	0.706	-0.117	0.094	-0.165	-0.398	
CGL	1.13	-0.57	1.84	1.03	0.34	3.3	0.22	-4.13	-6.15	-4.6	3.94	0.52	-2.84	0.6	-0.66	0.1	-0.08	0.16	0.38	
CGW	-0.79	0.36	-0.81	-0.77	-0.15	-2.73	-0.16	1.71	4.5	3.11	-1.62	-1.26	2.35	-0.41	0.4	-0.06	0.04	-0.09	-0.23	
CL/CB	1.21	-0.55	1.67	0.97	0.3	3.24	0.19	-3.67	-5.95	-4	3.49	0.92	-3.21	0.53	-0.59	0.08	-0.06	0.15	0.34	
ER	-0.88	0.5	-1.65	-1.11	-0.3	-3.61	-0.26	3.72	6.43	5.28	-3.46	-0.75	2.51	-0.68	0.69	-0.12	0.08	-0.16	-0.4	
ASV	-0.98	0.57	-1.65	-1.15	-0.3	-3.45	-0.28	3.65	6.11	5.59	-3.56	-0.7	2.59	-0.64	0.73	-0.12	0.09	-0.16	-0.4	
GC	0.06	-0.16	0.63	0.49	0.12	1.53	0.11	-1.48	-2.7	-2.2	1.35	0.24	-0.87	0.27	-0.3	0.3	-0.02	0.08	0.16	
AC	0.41	-0.33	0.85	0.63	0.17	2.11	0.17	-1.82	-3.29	-3.5	2.03	0.31	-1.26	0.37	-0.41	0.04	-0.15	0.1	0.21	
Fe	-0.75	0.47	-1.67	-0.93	-0.28	-2.55	-0.26	3.44	4.69	5.19	-3.42	-0.64	2.57	-0.59	0.63	-0.13	0.08	-0.18	-0.35	
Zn	-0.97	0.54	-1.69	-1.14	-0.31	-3.41	-0.27	3.79	5.97	5.42	-3.5	-0.7	2.59	-0.64	0.69	-0.12	0.08	-0.15	-0.42	

UNDER PEER REVIEW