

DEVELOPMENT AND EVALUATION OF HYBRIDS RESISTANT TO LATE BLIGHT AND LEAF CURL VIRUS DISEASES IN TOMATO

Abstract

The experimental material included 21 F₁ hybrids (developed by half diallel fashion), 7 parents and standard check were all planted in randomized complete block design with three replications. Among the parents Pusa Rohini, Arka Vikash and *S. pimpinellifolium* whereas, among the crosses, Pusa Rohini × Arka Vikash, H-86 × Arka Vikash and Arka Vikash × *S. pimpinellifolium* were showed -highly resistant for per cent disease incidence and coefficient of infection of late blight and ToLCV. Maximum heterosis over better parent and standard parent in desirable direction were found in Pusa Rohini × Arka Vikash (-48.61, -52.84), (-72.12, -76.42) and Arka Vikash × Arka Abha (-42.02, -50.35), (-61.35, -75.17) for per cent disease incidence and coefficient of infection of late blight. Whereas heterosis for PDI and CI of ToLCV was found in cross combinations Pusa Rohini × *S. pimpinellifolium* (-46.35, -65.08), (-77.50, -88.35) and Arka Vikash × *S. pimpinellifolium* (-60.85, -64.32) (-86.95, -88.27) over better parent and standard parent. The cross identified as best specific combiners Arka Vikash × *S. pimpinellifolium* for per cent disease incidence and coefficient of infection of late blight and ToLCV. Among the parents for late blight the per cent disease incidence and coefficient of infection were significantly shown by Arka Vikash and Arka Abha. Against ToLCV (PDI and CI) resistance, parents Pusa Rohini, *S. pimpinellifolium* and Arka Vikash were exhibited significantly negative gca effect.

Key words: Heterosis, Combining Ability, Late Blight, ToLCV

Introduction

Tomato (*Solanum lycopersicum* L. Mill) is the second most important vegetable crop in the world and is grown practically throughout India. India is the second top tomato growing country after China contributed about 11 percent of the world tomato production (Anonymous, 2011). Worldwide growth and spread of tomato as a vegetable crop is limited by the fact that it is affected by a number of diseases causing substantial yield loss and also affecting the quality of fruits. Besides fungal, bacterial and mycoplasmal infection, it is also affected by large number of viral diseases. In tropics and subtropics, tomatoes are affected

Comment [AAS1]: Spell out all abbreviation as appear first in MS.

Write the location, duration and institute name where research was conducted. Define the objective of experiment in abstract clearly.

Comment [AAS2]: Spell out as appear first in MS.

Comment [AAS3]: See for grammatical error; split in 2 sentence if possible without losing information.

Comment [AAS4]: Is it general combining ability? (GCA)? Spell out if any other meaning?

Comment [AAS5]: Provide reference to the statements made in introduction section to make it scientifically correct; Provide current status of information available on disease resistance in tomato for different diseases in general and late blight and ToLCV in particular; Provide the information on significance of selected diseases in more elaborative form rather than citing single reference showing losses caused; Define the research gap which was addressed in present investigation followed by specific objectives of study;

with many diseases, which include late blight caused by *Phytophthora infestans* and tomato leaf curl virus disease, a viral disease. They cause huge losses and deterioration to fruit quality, quantity as well as yield (Nelson-SC, 2008). These phyto-pathogens have huge capability to generate new forms of infestation and infection, which can cause much destruction of the crop, leading to crop failure. *Phytophthora infestans* (Mont.) de Bary is not a true fungus, but rather is regarded as a fungus-like organism. This pathogen is currently classified as an Oomycete, which are members of the kingdom Chromista (Stramenopiles or Straminopiles): (Nelson-S, 2008). Among the diseases, the occurrence of ToLCV and late blight in tomato is a major constraint in cultivation of tomato during summer and rainy season in India. ToLCV is a monopartite, Gemini virus known to be transmitted by the vector white fly, *Bemisia tabaci* Genn. ToLCV is known to infect the crop at all the stages starting from nursery to fruit formation. Saikia and Muniyappa (1989) reported cent per cent infection and fruit yield losses up to 90 per cent. Host plant resistance is an important disease control strategy and environmentally safe, with low running costs. Therefore, screening tomato cultivars possessing inbuilt resistance is an appropriate approach for disease management.

Materials and Methods

The experiment was conducted during 2015-17 and the study comprised of genotypes collected from IIVR, Varanasi, ICAR-NBPGR, New Delhi and ICAR-IIHR, Bengaluru. After screening for disease, quality and yield parameters, seven tomato lines along with one check and 21 F₁ hybrids produced from crossing parental lines in a 7 × 7 half diallel fashion. The parental genotypes and 21 F₁ hybrids were grown in randomized block design with three replications and other agronomic practices were followed as per package of practices given by Fageria *et al.*, (2003). To assess the resistant of given strain symptom severity grade designated with numerical value of 0-4 scale were given on the basis of visual observation to quantify the disease severity and calculation were made according to the method described by Kalloo and Banerjee (1987). -This calculation used for parents and F₁s screening under natural condition. The present disease incidence and coefficient infection were calculated by the formula-

$$\text{PDI} = \frac{\text{No. of diseased plants}}{\text{No. of total plants}} \times 100$$

Comment [AAS6]: Is this is specific objective of study?

Comment [AAS7]: Incomplete and need to rewrite. Mention duration (Is it for 2 years?) in year as well as seasons for growing crops; Also some information on climatic condition of selected area.

Provide name of institute and georeferenced location of study area where research was conducted;

Write at least little (may in tabular form) about input addition, package of practices followed (Is it on field study of in potted plant or sterilized seed bed or some other specification).

Is any measure taken to insure management of disease other than selected disease (LB and To LCV).

Provide information on methodologies/ software followed for analysis of data

Comment [AAS8]: Lengthy sentence; split in to 2 sentences.

Coefficient Infection (CI) = Percent Disease Incidence (PDI) \times Response value (RV)

Result and Discussion

The per cent disease incidence and coefficient of infection of late blight in parents ranged from 36.39 (Arka Vikash) to 69.27 (BSS-488) and 21.85 (Arka Vikash) to 73.19 (Arka Ahuti). Among F_1 s populations it ranged from 32.66% (Pusa Rohini \times Arka Vikash) to 73.96% (Pusa Rohini \times *S.pimpinellifolium*) and 16.33 (Pusa Rohini \times Arka Vikash) to 73.96 (Pusa Rohini \times *S.pimpinellifolium*), respectively. The results were in accordance with the findings of Narayan *et al.* (2018). The resistant parents and crosses viz., Arka Vikash and *S. pimpinellifolium* and Pusa Rohini \times Arka Vikash, H-86 \times Arka Vikash, CLNB \times Arka Vikash and Arka Vikash \times *S.pimpinellifolium* can be utilized in future breeding programme. Per cent disease incidence and coefficient of infection of ToLCV among parents ranged from 22.58% (*S.pimpinellifolium*) to 60.09% (H-86) and 5.64 (*S.pimpinellifolium*) to 45.07 (H-86). Whereas among crosses it was varied between 19.25% (Pusa Rohini \times *S.pimpinellifolium*) to 70.00% (CLNB \times Arka Ahuti) and 4.81 (Pusa Rohini \times *S.pimpinellifolium*) to 70.00 (CLNB \times Arka Ahuti). The above findings are in agreement with the findings of Chellimi *et al.* (1994) and Bhattarai and Panthee (1998). The resistant parents and crosses viz., Pusa Rohini, Arka Ahuti and *S. pimpinellifolium* and Pusa Rohini \times Arka Ahuti, Pusa Rohini \times Arka Vikash, Pusa Rohini \times *S.pimpinellifolium*, H-86 \times Arka Vikash and Arka Vikash \times *S.pimpinellifolium*.....

The cross combination exhibiting negative and significant heterosis in case of (diseases) i.e., PDI and CI of late blight is an indication of low disease incidence it was observed in Pusa Rohini \times Arka Vikash (-48.61, -52.84), (-72.12, -76.42) and Arka Vikash \times Arka Abha (-42.02, -50.35), (-61.35, -75.17) over better parent and standard parent. Sixteen crosses exhibited non-significant (desirable) heterosis over the better parent and standard parent in the needful direction for PDI and CI of late blight. Associated characters were also reported by Arora *et al.* (2022). Sixteen crosses showed desirable heterosis over mid parent and seventeen crosses over standard parent. Whereas maximum desirable heterosis for PDI and CI of ToLCV was found in cross combinations Pusa Rohini \times *S.pimpinellifolium* (-46.35, -65.08), (-77.50, -88.35) and Arka Vikash \times *S.pimpinellifolium* (-60.85, -64.32) (-86.95, -88.27) over better parent and standard parent. Fourteen crosses out of twenty-one exhibited desirable negative heterosis over better parent and standard parent for PDI and CI

Comment [AAS9]: Discuss the findings in more elaborative way such as, effect of environment and environment \times genotype interaction, need and significance of outliers (both desirable or undesirable) for important characters; Avoid writing "results are in co-ordination with" and explain the relation of past studies with the data presented in present investigation; If the findings of earlier workers are already exist, than for what purpose the author has repeated the work? Is the yield was calculated (As ultimately the economic damage is expressed in terms of yield increase or decrease? If yes, than provide the data of yield discuss in terms of impact of diseases incidence on yield for all combination. Cite table number at appropriate places in result and discussion section. Avoid citing too old references if new recent references were available. Is there any relation of selected diseases with other diseases or pest or growing condition; If yes describe it as part of discussion (If no write the reasons for the same to strengthen discussion section). The result presented describe the results of screen of genotypes and crosses for two diseases incidence/impact and there is no any development activity involved in this MS- See the title.

Comment [AAS10]: Is it complete sentence? Please check.

of ToLCV Similar reports were also reported by Narayan *et al.* 2018, Sowjanya and Sridevi 2020, Arora *et al.* 2022 and Sundharaiya *et al.* 2018.

Among parents for PDI and CI of late blight the varieties, Arka Vikash and Arka Abha (-13.80, -1.41), (-18.05, -4.62) were found good general combiners they exhibited negative and significant gca effect. Against ToLCV (PDI and CI) resistance, among parents Pusa Rohini, *S. pimpinellifolium* and Arka Vikash (-6.51, -5.47), (-7.67, -5.47), (-4.85, -5.09) were found good general combiner. Hence, these three parents Arka Vikash, Pusa Rohini and *S.pimpinellifolium* may be used extensively in breeding programme aimed at the development of high yielding with quality tomato hybrids along with resistance to late blight and ToLCV diseases. Similar findings were also reported by Kulkarni 1999 and Arora *et al.* 2022. Out of 21 crosses, 6 for PDI and 5 for CI of late blight exhibited significant SCA effects in the desirable direction. The maximum SCA effects in the desirable direction was exhibited by H-86 × Arka Ahuti and Arka Vikash × *S.pimpinellifolium* for PDI (-14.23, -9.86) and CI (-22.34, -13.89) of late blight. Elsayed *et al.* (2016) were also reported significant SCA effect in desirable direction for late blight. The negative and significant SCA effect was expressed by six crosses for PDI and CI of ToLCV. Whereas the maximum significant SCA effect was exhibited by CLNB × Arka Abha and Arka Vikash × *S.pimpinellifolium* for PDI (-24.55, -12.80) and CI of ToLCV (-27.68, -13.21) and this is also concordant with Singh *et al.* 2011, Singh *et al.* 2014 and Arora *et al.* 2022.

Conclusion/ Conclusions: Missing?

References

Elsayed, A.Y., Elsaid, E. M. and Elsherbiny, E. A. (2016). The performance of late blight gene ph-3 in tomato under the effect of local populations from *phytophthora infestans*. *J. Plant Production, Mansoura Univ.*, 7 (3): 361- 371.

Anonymous (2011). National Horticulture Board, Department of Agriculture and cooperation, Government of India. Retrieved from www.nhb.gov.in.

Arora, H., Jindal, S. K., Sharma, A., Gill, R. and Chawla, R. (2022). Development and evaluation of hybrids resistant to late blight and leaf curl virus diseases in tomato. *Genetika*. 54(2): 801-816.

Bhattarai, S. P. and Panthee, D. R. 1998. Identification of the parents for production of bacterial wilt resistant tomato lines. *Himachal J. Agric. Res.*, 23: 40-44.

Comment [AAS11]: ?

Formatted: Font: Bold

Formatted: Indent: First line: 0"

- Chellemi, D. O., Dankers, H. A., Olson, S. M., Hodge, W. C. and Scoot, J. W. (1994). Evaluating bacterial wilt resistant genotypes using a regional approach. *J. American Soc. Hort. Sci.*, 119 (2): 325-329.
- Fageria, M. S., Chaudhury, B. R. and Dhaka, R. S. 2003. Vegetable crops production technology. 25–40p.
- Kaloo, G. and Banerjee, M. K. (1989). Transfer of tomato leaf curl virus resistance from *Lycopersicon hirsutum f. glabrum* to *L. esculentum*. *Plant Breed.* 105, 156–159.
- Kulkarni, G. P. (1999). Heterosis, combining ability and reaction to tomato leaf curl virus in tomato. M.Sc. (Agri.) Thesis, Univ. Agric. Sci., Dharwad, pp. 1-107.
- Narayan, R. P. J., Mallesh, S. B., Patil, M. G. and Dhotre, M. (2018). Heterosis and combining ability for tomato leaf curl virus (ToLCV) and Bacterial wilt disease in tomato (*Solanum lycopersicum* L.) *Electronic Journal of Plant Breeding*, 9(1): 73-81
- Nelson Scot C. (2008). Late Blight of Tomato (*Phytophthora infestans*). *Plant Disease*. PD-45.
- Saikia, A. K. and Muniyappa, V. (1989). Epidemiology and control of tomato leaf curl virus in Southern India. *Trop. Agric. (Trinidad)*, 66: 350-364
- Singh, R. K., Rai, N., Singh, M., Singh, A. K., Kumar, P., Singh, R. and Singh, S. N. (2011) Molecular diversity in tomato genotypes and their exploitation of heterosis against tomato leaf curl virus (ToLCV) and yield traits. Abstract in National Symposium on Vegetable Biodiversity, pp. 47–48. Department of Horticulture, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, India.
- Singh, R. K., Rai, N., Singh, M., Singh, S. N. and Srivastava, K. (2014). Genetic analysis to identify good combiners for ToLCV resistance and yield components in tomato using interspecific hybridization. *J. Genet.* 93(3): 623–629.
- Sowjanya, B. A. and Sridevi, O. (2020). Combining ability and heterosis studies in tomato (*Solanum lycopersicum* L.) under ToLCV disease stress condition. *International Journal of Chemical Studies*. 8(3): 2134-2141.

Sundharaiya, K., Karuthamani, M. and Sathish, G. (2018). Estimate of heterosis and per se performance of tomato f_1 hybrids for leaf curl virus resistance. *Int.J.Curr.Microbiol.App.Sci* 6: 189-196.

UNDER PEER REVIEW

Table: 1 Mean, sca effects for late blight and ToLCV incidence, coefficient of infection in tomato

Crosses	Per cent late blight incidence		Late blight coefficient of infection		Per cent ToLCV incidence		ToLCV coefficient of infection	
	Mean	SCA effects	Mean	SCA effects	ToLCV (%)	SCA effects	Mean	SCA effects
Pusa Rohini×H-86	50.89	-6.09*	38.17	-12.36*	35.74	-2.29	17.87	-3.74
Pusa Rohini×CLNB	55.36	-3.90	41.52	-9.73*	59.33	11.53**	44.50	9.57**
Pusa Rohini×Arka Ahuti	55.74	-5.82*	46.82	-7.36	31.13	-7.14**	15.57	-8.30*
Pusa Rohini×Arka Vikash	32.67	-10.41**	16.34	-14.06**	27.71	-3.45	11.83	-3.95
Pusa Rohini×Arka Abha	61.12	5.66*	56.52	12.69*	36.65	-0.12	21.92	0.70
PusaRohini× <i>S.pimpinellifolium</i>	73.97	10.98**	73.97	16.42**	19.25	-11.29**	4.82	-10.67**
H-86×CLNB	53.94	-3.52	45.48	-4.90	56.82	0.49	42.61	-0.72
H-86×Arka Ahuti	45.63	-14.13**	30.97	-22.34**	36.00	-10.79**	21.48	-10.80**
H-86×Arka Vikash	41.78	0.50	31.34	1.81	24.10	-15.58**	12.05	-12.13**
H-86×Arka Abha	46.93	-6.74*	35.20	-7.76	42.95	-2.34	25.63	-4.00
H-86× <i>S.pimpinellifolium</i>	71.52	10.32**	71.52	14.84**	42.50	3.43	25.19	1.29
CLNB×Arka Ahuti	61.21	-0.83	51.43	-2.60	70.00	13.43**	70.00	24.41**
CLNB×Arka Vikash	38.18	-5.37	21.40	-8.84	51.17	1.72	38.38	0.88
CLNB×Arka Abha	52.94	-3.01	39.70	-3.98	30.51	-24.55**	15.26	-27.68**
CLNB× <i>S.pimpinellifolium</i>	68.06	4.59	68.06	10.67*	59.42	10.58**	54.81	17.60**
Arka Ahuti×Arka Vikash	58.34	12.48**	48.87	15.69**	37.69	-2.23	18.84	-7.60*
Arka Ahuti×Arka Abha	55.03	-3.22	41.27	-5.34	44.17	-1.36	29.84	-2.04
ArkaAhuti× <i>S.pimpinellifolium</i>	59.56	-6.21*	49.80	-10.52	51.37	12.06**	38.52	12.37**
Arka Vikash×Arka Abha	34.40	-5.37	17.20	-5.63	37.28	-1.14	22.20	-1.59
ArkaVikash× <i>S.pimpinellifolium</i>	37.43	-9.86**	22.65	-13.89**	19.40	-12.80**	4.85	-13.21**
ArkaAbha× <i>S.pimpinellifolium</i>	58.01	-1.67	43.51	-6.46	53.83	16.02**	40.37	16.87**
C.V.	9.03		18.50		10.54		19.49	
S.E.	2.89		4.95		2.61		3.26	
C.D. 5%	8.19		14.02		7.40		9.24	

Table: 2 Mean, gca effects for late blight and ToLCV incidence, coefficient of infection in tomato

Genotypes	Per cent late blight incidence		Late blight coefficient of infection		Per cent ToLCV incidence		ToLCV coefficient of infection	
	Mean	SCA effects	Mean	SCA effects	Mean	SCA effects	Mean	SCA effects
Pusa Rohini	63.57	1.91*	58.61	2.95	35.89	-6.51**	21.41	-7.67**
H-86	65.02	0.11	65.02	2.08	60.09	2.02*	45.07	0.74
CLNB	65.75	2.38*	60.79	2.80	59.50	11.79**	44.63	14.06**
Arka Ahuti	73.19	4.68**	73.19	5.73**	45.05	2.26**	30.51	3.00**
Arka Vikash	36.39	-13.80**	21.85	-18.05**	49.55	-4.85**	37.16	-5.09**
Arka Abha	59.33	-1.41	44.50	-4.62**	50.79	0.76	38.09	0.34
<i>S.pimpinellifolium</i>	63.13	6.12**	58.15	9.10**	22.59	-5.47**	5.65	-5.38**
BSS-488	69.27	1.91*	69.27	2.95	55.13	-6.51**	41.35	-7.67**
C.V.	9.03		18.50		10.54		19.49	
S.E.	2.89		4.95		2.61		3.26	
C.D. 5%	8.19		14.02		7.40		9.24	

Table:3 Heterosis (%) over better parent and standard parent for late blight and ToLCV incidence, coefficient of infection in tomato

Crosses	Per cent late blight incidence		Late blight coefficient of infection		Per cent ToLCV incidence		ToLCV coefficient of infection	
	BPH (%)	SPH (%)	BPH (%)	SPH (%)	BPH (%)	SPH (%)	BPH (%)	SPH (%)
Pusa Rohini×H-86	-21.73**	-26.54**	-41.29**	-44.90**	-40.53**	-35.18**	-60.35**	-56.78**
Pusa Rohini×CLNB	-15.80*	-20.08**	-31.70**	-40.06**	-0.29	7.61	-0.28	7.62
Pusa Rohini×Arka Ahuti	-23.85**	-19.54**	-36.03**	-32.41**	-30.89**	-43.54**	-48.97**	-62.35**
Pusa Rohini×Arka Vikash	-48.61**	-52.84**	-72.12**	-76.42**	-44.08**	-49.75**	-68.17**	-71.39**
Pusa Rohini×Arka Abha	-3.85	-11.76	-3.56	-18.41	-27.84**	-33.53**	-42.46**	-46.99**
PusaRohini× <i>S.pimpinellifolium</i>	16.35*	6.78	26.21*	6.78	-46.35**	-65.08**	-77.50**	-88.35**
H-86×CLNB	-17.97**	-22.14**	-30.05**	-34.34**	-5.45	3.05	-5.46	3.06
H-86×Arka Ahuti	-37.65**	-34.13**	-57.69**	-55.30**	-40.09**	-34.70**	-52.35**	-48.06**
H-86×Arka Vikash	-35.75**	-39.69**	-51.80**	-54.76**	-59.90**	-56.29**	-73.26**	-70.85**
H-86×Arka Abha	-27.82**	-32.25**	-45.87**	-49.19**	-28.53**	-22.10**	-43.14**	-38.02**
H-86× <i>S.pimpinellifolium</i>	9.99	3.24	9.99	3.24	-29.27**	-22.91**	-44.11**	-39.08**
CLNB×Arka Ahuti	-16.37**	-11.64	-29.74**	-25.76*	17.65**	26.96**	56.86**	69.29**
CLNB×Arka Vikash	-41.93**	-44.88**	-64.79**	-69.10**	-13.99*	-7.18	-14.00	-7.18
CLNB×Arka Abha	-19.49**	-23.58**	-34.69**	-42.69**	-48.72**	-44.66**	-65.81**	-63.10**
CLNB× <i>S.pimpinellifolium</i>	3.52	-1.75	11.96	-1.75	-0.13	7.78	22.82*	32.55**
Arka Ahuti×Arka Vikash	-20.30**	-15.79*	-33.23**	-29.45**	-23.94**	-31.64**	-49.30**	-54.43**
Arka Ahuti×Arka Abha	-24.82**	-20.57**	-43.61**	-40.42**	-13.02	-19.88**	-21.67	-27.84*
ArkaAhuti× <i>S.pimpinellifolium</i>	-18.62**	-14.02*	-31.96**	-28.11**	14.03	-6.83	26.25	-6.84
Arka Vikash×Arka Abha	-42.02**	-50.35**	-61.35**	-75.17**	-26.59**	-32.38**	-41.72**	-46.31**
ArkaVikash× <i>S.pimpinellifolium</i>	-40.71**	-45.97**	-61.05**	-67.30**	-60.85**	-64.82**	-86.95**	-88.27**
ArkaAbha× <i>S.pimpinellifolium</i>	-8.11	-16.26**	-25.18	-37.19**	5.99	-2.37	5.99	-2.36