

EVALUATION OF TOTAL VIABLE BACTERIA COUNTS AND *IN SITU* CHARACTERISTICS IN DRINKING WATER SOURCES IN SAGBAMA TOWN, BAYELSA STATE, NIGERIA

ABSTRACT

Before the 21st century, most of Nigeria's rural and coastal communities obtained their drinking water from surface water. However, due to development, many coastal towns are sourcing their drinking water from groundwater. Nevertheless, surface water remains a crucial source of drinking water in communities where groundwater is inaccessible. This study evaluated the density of total heterotrophic bacteria and *in situ* characteristics of drinking water sources in Sagbama town in the Sagbama Local Government Area of Bayelsa State, Nigeria. Triplicate water samples were obtained from five stations for each of the water types (ground and surface water). The water samples were analyzed following standard procedures. The results of ground and surface water ranged from 2.33 to 2.86 (overall mean of 2.52) Log CFU/mL and 2.70 to 3.87 (overall mean of 3.24) Log CFU/mL for total heterotrophic bacteria counts, 7.05 to 7.68 (overall mean 7.40) and 7.06 to 7.31 (overall mean 7.22) for pH, 146.67 to 232.00 (overall mean 169.13) mg/L and 40.23 to 45.60 (overall mean 41.50) mg/L for total dissolved solids, 207.57 to 327.67 (overall mean 239.31) μ S/cm and 56.73 to 64.47 (overall mean 58.55) μ S/cm for conductivity, and 0.11 ppt and 0.03 ppt (in all the stations) for salinity. The temperatures of both kinds of water were in the order of 26°C range. Except for the pH, salinity, and conductivity of groundwater, an analysis of variance reveals no significant differences ($p > 0.05$) across sites for any water type. The overall t-test revealed that all parameters, except pH and temperature, were statistically distinct ($p < 0.05$) between the various water types. According to Pearson's correlation conductivity correlates positively with total dissolved solids for both kinds of water. In contrast, salinity correlates positively with conductivity and total dissolved solids for surface water at $p < 0.01$. Except for total heterotrophic bacteria counts on surfaces, which exceeded World Health Organization standards, other parameters fell within Standard Organization of Nigeria and World Health Organization guidelines. Therefore, drinking surface water from the research location without first purifying it increases the chance of avoidable ill health consequences.

Keywords: Human activities, Microorganisms, Pollution, Water-borne diseases, Water Quality

1. Introduction

Water is a vital resource essential to the existence of all living things [1-3]. Solid water (ice), gaseous water (vapor), and liquid water are the three distinct forms of water used to categorize various water resources. Water may also be found underground, on the surface, or as precipitation from the sky, i.e., rainfall. According to their origins, the three types of surface

water resources are fresh water, salt water (also known as estuarine water), and brackish water (estuarine water, salt, and freshwater interphase).

Additionally, water offers a home for several plant and animal species [4]. As a consequence, the life cycles of species responsible for transmitting illnesses to people and other animals, such as tadpoles, mosquitoes, and parasites, are limited to water [5-10]. Human disease transmissions are due to vectors such as *Aedes aegypti* (yellow and dengue fever), *Culex quinquefasciatus* (lymphatic filariasis), and *Anopheles gambiae*, *A. funestus*, *A. arabiensis*, and *A. melas* (malaria) [11-13].

Surface water is the most commonly used resource for transportation-related activities, although it has the lowest overall availability. The most prominent surface water uses are in recreational activities such as swimming, bathing, washing, cooking, and drinking. Additionally, groundwater is an essential resource for home use. Izah et al. [14] and Izah and Srivastav [4] stated that rain, ground, and surface waters are sources of potable water in Nigeria. In the coastal areas of the Niger Delta, human drinking water sources are often contaminated [15, 16]. Human actions and, to a lesser degree, natural occurrences are often responsible for the pollution of water supplies. Inappropriate residential and industrial waste disposal (municipal, solid, and effluents) contributes to surface water contamination. These include dredging, oil and gas operations, agricultural operations, and food production [15-22]. Additionally, improper management and conservation techniques contribute to groundwater pollution.

The leaching of over-filled soakaways causes groundwater pollution, resulting from poor home development in many coastal locations and insufficient sanitation services. Due to the high-water tables in certain coastal regions, notably in Bayelsa State [2], it is easy for effluents and runoff to penetrate the coastal aquifer. In addition, water pollution also increases nutrient levels, resulting in eutrophication, acidification, and hydrological changes. Consequently, it might harm the productive capacity of such waterways in terms of fish population composition and dispersion [23-25]. Consequently, changes in the entire home and industrial water demand are possible. The presence of microorganisms, heavy metals, and the water's basic physicochemical features are the three most essential factors for determining the overall quality of water [24 – 33]. In addition, in several regions of Bayelsa State, surface water [27,30] and groundwater [34] reports on drinking water sources.

Microorganisms are widespread. Moreover, it is well-known that they cause several infectious illnesses. Water contaminated by microorganisms, such as harmful bacteria, parasites, and viruses, increase the chances of human illness. As a result, poor-quality water can potentially serve as a transmission medium for harmful microorganisms such as bacteria and viruses. Bayelsa State's drinking water sources (surface and groundwater) include several microbial species. Among these microorganisms are *Pseudomonas*, *Enterobacter*, *Bacillus*, *Citrobacter*, *Erminia*, *Klebsiella*, *Shigella*, *Salmonella*, *Proteus*, *Serratia*, *Streptococcus*, *Micrococcus*, *Yersia*, *Corynebacterium*, *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus niger* [24, 29]. In addition, some of these microorganisms occur in soil and air habitats [25], as well as in ready-to-eat meals and food beverages in different regions of Bayelsa State [36-38]. Despite these studies, the literature has minimal information on the drinking water quality in Sagbama town, Sagbama Local Government Area of Bayelsa State. Therefore, the focus of this study is to assess the microbial quality and in-situ characteristics of potable water in Sagbama town. The findings

of this work will be of benefit to the users of the various water types under study and the general public.

2. Materials and Methods

2.1 Study Area

The study was conducted in Sagbama town, which stands as the headquarters of the Sagbama local government area in Bayelsa State. The primary occupation of the indigenous people of the area is farming, including fishing. Few other people are civil servants. Like other major communities in the area, surface water passes through the town. The surface water is a major receiver of municipal solid waste. Also, the water is also used for domestic purposes before now although some people still use the water. Due to industrialization, many people in the area are now using groundwater as their drinking water supply.

2.2 Sample collection

A total of thirty water samples were collected from the study area, which were labeled A, B, C, D, and E. Therefore, fifteen samples were collected for a particular water type (ground and surface water). For each type of water, triplicate samples were collected. The water samples were collected in sterile containers.

2.3 Enumeration of total heterotrophic bacteria counts

Nutrient Agar was used as a medium to enumerate total heterotrophic bacteria. Previously reported by Pepper and Gerba [39] and Benson [40], the pour plate method was adopted. The nutrient agar used was prepared following the manufacturer's directions. A 1.0 ml water sample was aseptically plated and incubated inverted at 37°C for 24 to 48 hours. For each water sample, the outcome was a colony count expressed as colony-forming units.

2.4 Determination of the in-situ characteristics of the water

The manufacturers' guide determined all the in-situ parameters using a multipurpose meter that analyses pH, total dissolution, conductivity, salinity, and temperature.

2.5 Statistical analysis

The statistical analysis was done by SPSS software version 20, which expressed the data as mean and standard deviation. One-way analysis of variance at $P = 0.05$ and Waller-Duncan's statistics discern the source of the observed differences for each water type across the various locations. A t-test compares the overall water quality for each parameter. Finally, the Spearman rho correlation matrix identifies the relationship between the in-situ parameters and total heterotrophic bacteria counts.

3. Results and Discussion

The levels of total heterotrophic bacteria count and in-situ characteristics of ground and surface waters from Sagbama town, Bayelsa State, Nigeria, are shown in Tables 1 and 2, respectively. The total heterotrophic bacteria counts ranged from 2.33–2.86 Log CFU/mL and 2.70–3.87 Log CFU/mL for ground and surface water, respectively. There was no statistical deviation ($p > 0.05$)

in the different locations for each water type. On average, the mean values of total heterotrophic bacteria counts were 2.52 Log CFU/mL and 3.24 87 Log CFU/mL for ground and surface water, respectively, being statistically different ($p = 0.001$) (Table 3). The values showed that the total heterotrophic bacteria counts were higher in the surface water than in the groundwater since several human activities occur in the surface water. The lack of significant variation in the overall heterotrophic bacteria counts across the various water types and locations implies that similar activities contaminate each water type in the region [24].

This study contradicts prior research on drinking water quality in Bayelsa State. 1.78 to 9.30×10^6 CFU/ml from river nun at Amassoma axis [23]; 5.38 to 6.74 Log CFU/ml from Epie stream[24]; 4.53 to 5.82 logs CFU/ml in Taylor Creek [41]; 0.74 to 8.43×10^6 CFU/ml from Ikoli creek [17] and $2.97 - 6.03 \times 10^4$ CFU/ml from Imiringi [34]. Variations in these values reflect human activities that influence the density of total heterotrophic bacteria counts in drinking water sources. The levels above the World Health Organization/Food and Agriculture Organization permissible limit of 1.0×10^2 CFU/ml for surface water [29] were within the range for surface water. Consequently, the overall density of heterotrophic bacteria in surface water exceeded the permissible level and potential health risks exist.

The pH range for ground and surface water was 7.05 to 7.68 and 7.06 to 7.31, respectively. For surface water, there was no statistical variance across sites; however, there was a statistical deviation for groundwater. The post hoc test revealed no divergence between stations A-C and C-E. Nonetheless, there was a difference between A and D. The average pH values for groundwater and surface water were 7.40 and 7.22, respectively, with no statistically significant difference ($p = 0.057$). (Table 3). The modest variation in pH levels for groundwater may be attributable to variances in compounding variables, such as the depth of the aquifer, direction of flow of the water, materials, and water treatment procedures. Essentially, the values are comparable to those obtained in some regions of Bayelsa State.

Some of the studies include Nun River at the Gbarantoru and Tombia towns (6.27–6.45) [42]; 5.80–7.01 from Epie Creek [25]; Nun River at Amassoma axis (6.53–7.35) [2,33]. The discrepancy is attributable to the quantity of acidic material leached into the water due to several human activities [25,42]. Nevertheless, based on the water quality standard established by the Standard Organization of Nigeria (SON) and the World Health Organization (WHO), which is between 6.5 and 8.5, the results discovered in this research indicate that the water is safe to consume.

Table 1: Level of Total heterotrophic bacteria counts and in-situ characteristics of groundwater from Sagbama town, Bayelsa State, Nigeria

Locations	THB, Log CFU/mL	pH	Salinity, ppt	Conductivity, μ S/cm	TDS, mg/L	Temperature, $^{\circ}$ C
A	2.33 \pm 0.11 a	7.68 \pm 0.23c	0.11 \pm 0.01 a	207.57 \pm 11.66 a	146.67 \pm 7.51a	26.60 \pm 0.10a
B	2.86 \pm 0.06 a	7.54 \pm 0.13bc	0.11 \pm 0.01 a	229.00 \pm 20.66 a	162.00 \pm 15.10 a	26.47 \pm 0.06a
C	2.40 \pm 0.11 a	7.50 \pm 0.39ab c	0.11 \pm 0.08 a	327.67 \pm 65.58 a	232.00 \pm 46.81 a	26.47 \pm 0.23a
D	2.64 \pm 0.60 a	7.05 \pm 0.08a	0.11 \pm 0.01 b	220.33 \pm 12.90 b	154.33 \pm 12.66 a	26.50 \pm 0.20a
E	2.35 \pm 0.04 a	7.20 \pm 0.13ab	0.11 \pm 0.01 a	212.00 \pm 17.69 a	150.67 \pm 12.58 a	26.43 \pm 0.06a

Data is expressed as mean \pm Standard deviation; Different alphabets along the column indicate significance difference ($P < 0.05$) according to Waller Duncan statistics

Table 2: Level of Total heterotrophic bacteria counts and in-situ characteristics of surface water from Sagbama town, Bayelsa State, Nigeria

Locations	THB, Log CFU/mL	pH	Salinity, ppt	Conductivity, μ S/cm	TDS, mg/L	Temperature, $^{\circ}$ C
A	3.87 \pm 0.49a	7.30 \pm 0.17a	0.03 \pm 0.01a	57.63 \pm 3.28a	40.75 \pm 2.22a	26.53 \pm 0.12a
B	3.04 \pm 0.52a	7.28 \pm 0.12a	0.03 \pm 0.00a	56.73 \pm 2.94a	40.23 \pm 2.12a	26.50 \pm 0.10a
C	2.70 \pm 0.16a	7.06 \pm 0.06a	0.03 \pm 0.00a	56.80 \pm 1.31a	40.40 \pm 0.98a	26.50 \pm 0.10a
D	3.05 \pm 0.63a	7.14 \pm 0.06a	0.03 \pm 0.00a	57.13 \pm 2.58a	40.50 \pm 1.64a	26.57 \pm 0.06a
E	3.53 \pm 1.18a	7.31 \pm 0.23a	0.03 \pm 0.01a	64.47 \pm 13.28a	45.60 \pm 9.70a	26.57 \pm 0.15a

Data is expressed as mean \pm Standard deviation; Different alphabets along the column indicate significance difference ($P < 0.05$) according to Waller Duncan statistics

Table 3: Comparative analysis of drinking water sources (surface and ground water) in Sagbama town, Bayelsa State, Nigeria

Parameters	N	Ground water	Surface water	t-value	P-value	Implications

		Mean± Std. Deviation	Mean± Std. Deviation			
THB, Log CFU/mL	15	2.52±0.32	3.24±0.72	-3.558	0.001	Significant deviation
pH	15	7.40±0.31	7.22±0.16	1.982	0.057	No significant deviation
Salinity, ppt	15	0.11±0.03	0.03±0.00	9.422	0.000	Significant deviation
Conductivity, μS/cm	15	239.31±53.96	58.55±6.22	12.889	0.000	Significant deviation
TDS, mg/L	15	169.13±38.54	41.50±4.46	12.742	0.000	Significant deviation
Temperature, °C	15	26.49±0.14	26.53±0.10	-0.913	0.369	No significant deviation

The salinity of groundwater was 0.11 ppt, while that of surface water was 0.03 ppt. There was no statistical difference between the sites for both kinds of water. However, a significant difference ($p < 0.001$) existed across the water types (Table 3). There was no noticeable variation in water quality across the various water types, indicating that their freshness levels are equivalent [25]. However, the statistical variance found for the various water types indicates that their properties vary. The results found in this study are comparable to those obtained in earlier research from the Nun River in Gbaratoru and Tombia (0.03 mg/l) [42] and Ikoli Creek (0.01-0.03 mg/l) [32]. Furthermore, the levels were within the WHO guidelines for drinking water (100–200ppt).

The range of total dissolved solids for groundwater was 146.67 to 232.00 mg/L, whereas the range for surface water was 40.23 to 45.60 mg/L. There was no statistical difference between the sites for both kinds of water. The mean values of total dissolved solids for groundwater and surface water were 169.13 mg/L and 41.50 mg/L, respectively, which were statistically distinct ($p < 0.001$) (Table 3). The discrepancy between the two kinds of water revealed changes in salt content. Again, water has various chemical qualities and human activity fluctuations can alter its quality and geology. In general, the values reported are comparable to those previously documented in Epie Creek [25] and are greater than those in the Nun River [2,33] and Kolo Creek [31]. However, the total dissolved solids observed in this study are far below the SON and WHO limits of 500 mg/L.

The range of conductivity for ground and surface water was 212.00–327.67 μS/cm and 56.73–64.47 μS/cm, respectively. For surface water, there was no statistical variance across sites. However, there was a statistical deviation for groundwater. Station D is the cause of the observed statistical variation based on the post hoc test. Overall, the mean conductivity values for ground and surface water were 239.31S/cm and 58.55S/cm, respectively, which were statistically distinct ($p < 0.001$) (Table 3). The statistical dissimilarity identified in the groundwater for location D is due to depth and treatment materials.

The values reported in this research are comparable to those obtained previously from Epie Creek [25] and Kolo Creek [31]. The conductivity of water may provide information about the overall concentration of ionic solutes in water [25,33]. According to WHO and SON's

recommendations for safe drinking water, the water's conductivity falls within the acceptable range of 1000 S/cm.

Ground and surface water temperatures are within a range of 26 degrees Celsius. There was no statistical difference between the sites for both kinds of water. In addition, there was no significant difference ($p = 0.369$) in the mean values of the two water types (Table 3). The absence of statistical variance indicates that the ambient air environment during the water sample was comparable. Also, it indicates that human activities did not statically modify the water over the research period. The average values reported are within the range of values that promote the survival of many mesophilic species within the temperature range for potable water. These values are marginally lower than those found in the Nun River [2], Kolo Creek [31], and Ikoli Creek [32] but slightly higher than those found in the Nun River by Ogamba et al. [33]. This discrepancy depends on the sample period's meteorological circumstances, such as the season and time of day.

Tables 4 and 5 illustrate the Pearson correlation between ground and surface water characteristics in Sagbama, Bayelsa State, Nigeria. At $p = 0.01$, there was a statistically significant positive correlation between conductivity, total dissolved groundwater (Table 4), and surface water (Table 5). Moreover, surface water salinity correlates with conductivity and total dissolved solids at a significance level of $p = 0.01$. Similar factors may have a positive effect on parameters that are positively correlated. The ions in the water may have affected the connections between conductivity and total dissolved solids and between conductivity, total dissolved solids, and salinity observed in the study. According to Rusydi [43], conductivity and total dissolved solids are utilized to determine salt levels as water quality indicators. In addition, the author said that total dissolved solids are crucial because they assist explain groundwater quality, namely inorganic salts, organic detritus, and saltwater intrusion. Conductivity and total dissolved solids are essential for determining the water's salinity. Therefore, there is no need for a link between salinity, conductivity, and total dissolved solids. Natural and artificial or manufactured variables, such as the salinity of the water, may be the primary determinants of the relationships.

Table 4: Pearson correlation of the ground water characteristics in Sagbama town, Bayelsa State, Nigeria

		THB	pH	Salinity	Conductivity	TDS	Temperature
THB	Pearson Correlation	1.000					
	Sig. (2-tailed)						
	N	15.000					
pH	Pearson Correlation	-0.063	1.000				
	Sig. (2-tailed)	0.823					
	N	15.000	15.000				
Salinity	Pearson Correlation	-0.018	0.441	1.000			
	Sig. (2-tailed)	0.950	0.100				
	N	15.000	15.000	15.000			
Conductivity	Pearson Correlation	0.000	0.178	-0.055	1.000		
	Sig. (2-tailed)	0.999	0.527	0.845			
	N	15.000	15.000	15.000	15.000		
TDS	Pearson Correlation	0.012	0.184	-0.058	0.999**	1.000	
	Sig. (2-tailed)	0.966	0.513	0.838	0.000		
	N	15.000	15.000	15.000	15.000	15.000	
Temperature	Pearson Correlation	0.233	0.272	-0.051	0.205	0.202	1.000
	Sig. (2-tailed)	0.403	0.326	0.857	0.463	0.470	
	N	15.000	15.000	15.000	15.000	15.000	15.000

** . Correlation is significant at the 0.01 level (2-tailed).

Table 5: Pearson correlation of the surface water characteristics in Sagbama town, Bayelsa State, Nigeria

		THB	pH	Salinity	Conductivity	TDS	Temperature
THB	Pearson Correlation	1.000					
	Sig. (2-tailed)						
	N	15.000					
pH	Pearson Correlation	0.300	1.000				
	Sig. (2-tailed)	0.278					
	N	15.000	15.000				
Salinity	Pearson Correlation	0.061	0.284	1.000			
	Sig. (2-tailed)	0.830	0.305				
	N	15.000	15.000	15.000			

Conductivity	Pearson Correlation	-0.073	0.415	0.787**	1.000		
	Sig. (2-tailed)	0.797	0.124	0.001			
	N	15.000	15.000	15.000	15.000		
TDS	Pearson Correlation	-0.072	0.418	0.779**	0.999**	1.000	
	Sig. (2-tailed)	0.799	0.121	0.001	0.000		
	N	15.000	15.000	15.000	15.000	15.000	
Temperature	Pearson Correlation	-0.041	-0.211	-0.139	-0.435	-0.452	1.000
	Sig. (2-tailed)	0.883	0.450	0.622	0.105	0.090	
	N	15.000	15.000	15.000	15.000	15.000	15.000

** . Correlation is significant at the 0.01 level (2-tailed).

Conclusion

This study evaluated the total viable bacterial density and in-situ characteristics of drinking water sources (surface and groundwater) in Sagbama town, in the Sagbama Local Government area of Bayelsa State. The results of the evaluation revealed that the density of bacteria was on the order of 10^2 – 10^3 CFU/ml, with a higher density found in the surface water, and that physico-chemical parameters showed an allowable level for drinking water. The characteristics of the water under study showed that they are within the Standard Organization of Nigeria and World Health Organization guidelines, except for the total heterotrophic bacteria in the surface water. The consumption of the surface water without treatment could expose users to health hazards. Therefore, there is a need to purify the water before use to prevent any potential health risks.

REFERENCES

1. Izah SC, Angaye TCN (2016) Heavy metal concentration in fishes from surface water in Nigeria: Potential sources of pollutants and mitigation measures. *Sky Journal of Biochemistry Research* 5(4): 31-47.
2. Agedah EC, Ineyougha ER, Izah SC, Orutugu LA (2015). Enumeration of total heterotrophic bacteria and some physico-chemical characteristics of surface water used for drinking sources in Wilberforce Island, Nigeria. *Journal of Environmental Treatment Techniques*, 3(1): 28 – 34.
3. Izah SC, Iyiola AO, Richard G (2023). Impacts of Pollution on the Hydrogeochemical and Microbial Community of Aquatic Ecosystems in Bayelsa State, Southern Nigeria. In: Madhav S, Singh VB, Kumar M, Singh S (Editors). *Hydrogeochemistry of Aquatic Ecosystems*. Published by John Wiley & Sons Ltd. Pp. 283 – 305. <https://doi.org/10.1002/9781119870562.ch13>.
4. Izah SC, Srivastav AL (2015). Level of arsenic in potable water sources in Nigeria and their potential health impacts: A review. *Journal of Environmental Treatment Techniques* 3(1):15 – 24.

5. Izah SC, Angaye TCN (2016). Ecology of Human Schistosomiasis intermediate host and Plant Molluscicides used for control: A review. *Sky Journal of Biochemistry Research*, 5(6): 075- 082
6. Youkparigha FO, Izah SC (2019). Larvicidal efficacy of aqueous extracts of *Zingiber officinale* Roscoe (ginger) against malaria vector, *Anopheles gambiae* (Diptera: Culicidae) *International Journal of Environmental and Agricultural Sciences*, 3:020.
7. Seiyaboh EI, Odubo TC, Izah SC (2020) Larvicidal Activity of *Tetrapleura tetraptera* (Schum and Thonn) Taubert (Mimosaceae) extracts against *Anopheles gambiae*. *International Journal of Advanced Research in Microbiology and Immunology*, 2(1): 20-25.
8. Seiyaboh EI, Seiyaboh Z, Izah SC (2020). Environmental Control of Mosquitoes: A Case Study of the Effect of *Mangifera Indica* Root-Bark Extracts (Family Anacardiaceae) on the Larvae of *Anopheles gambiae*. *Annals of Ecology and Environmental Science*, 4(1), 33-38.
9. Izah SC, Chandel SS, Epidi JO, Venkatachalam T, Devaliya R (2019). Biocontrol of *Anopheles gambiae* larvae using fresh ripe and unripe fruit extracts of *Capsicum frutescens* var. *baccatum*. *International Journal of Green Pharmacy*, 13 (4) | 338 – 342.
10. Izah SC (2019). Activities of Crude, Acetone and Ethanolic Extracts of *Capsicum frutescens* var. *minima* Fruit Against Larvae of *Anopheles gambiae*. *Journal of Environmental Treatment Techniques*, 7(2):196-200.
11. Ndiok EO, Ohimain EI, Izah SC (2016). Incidence of Malaria in Type 2 Diabetic patients and the effect on the liver: a case study of Bayelsa state. *Journal of Mosquito Research*, 6(15): 1-8.
12. Bassey SE, Izah SC (2017). Some determinant factors of Malaria Prevalence in Nigeria. *Journal of Mosquito Research*, 7(7): 48-58.
13. Bassey SE, Izah SC (2017). Nigerian plants with insecticidal potentials against various stages of mosquito development. *ASIO Journal of Medical and Health Sciences Research*, 2(1): 07-18
14. Izah SC, Chakrabarty N, Srivastav AL (2016) A Review on Heavy Metal Concentration in Potable Water Sources in Nigeria: Human Health Effects and Mitigating Measures. *Exposure and Health* 8:285–304.
15. Izah SC, Aigberua AO, Srivastav AL (2022). Factors influencing the alteration of microbial and heavy metal characteristics of river systems in the Niger Delta region of Nigeria. In: *Ecological Significance of river ecosystem: Challenges and management*. Madhav S, Kanhaiya S, Srivastav AL, Singh VB and Singh P (Editors). Published in United Kingdom by Elsevier Pp. 51-78. <https://doi.org/10.1016/B978-0-323-85045-2.00005-4>.
16. Izah SC, Ngun CT, Richard G (2022). Microbial quality of groundwater in the Niger Delta region of Nigeria: Health implications and effective treatment technologies. In: *Srivastav AL, Madhav S, Bhardwaj AK, Valsami-Jones E (Editors). Urban Water Crisis and Management: Strategies for Sustainable Development. Current Directions in Water Scarcity Research*. Pp. 149-172. Elsevier. <https://doi.org/10.1016/B978-0-323-91838-1.00010-5>
17. Seiyaboh EI, Izah SC (2017). Bacteriological assessment of a tidal creek receiving slaughterhouse wastes in Bayelsa state, Nigeria. *Journal of Advances in Biology and Biotechnology*, 14(1): 1 -7.

18. Seiyaboh EI, Izah SC (2017). A Review of Impacts of Gas Flaring on Vegetation and Water Resources in the Niger Delta Region of Nigeria. *International Journal of Economy, Energy and Environment*, 2(4): 48-55.
19. Seiyaboh EI, Izah SC (2017). Review of Impact of Anthropogenic Activities in Surface Water Resources in the Niger Delta region of Nigeria: A case of Bayelsa state. *International Journal of Ecotoxicology and Ecobiology*. 2(2): 61 – 73.
20. Izah, SC, Richard G, Sawyer WE (2021). Distribution of Fungi density and diversity in a Surface water of Epie Creek in Yenagoa Metropolis, Nigeria. *Archives of Epidemiology and Public Health*. doi: 10.15761/AEPH.1000121 Volume 3: 1-5.
21. Aigberua AO, Izah SC, Richard G (2021). Hazard Analysis of Trace Metals in Muscle of *Sarotherodon melanotheron* and *Chrysichthys nigrodigitatus* from Okulu River, Rivers State, Nigeria. *Journal of Environmental Health and Sustainable Development*, 6(3): 1340-1356.
22. Seiyaboh EI, Izah SC (2019). Impacts of Soil Pollution on Air Quality under Nigerian Setting. *Journal of Soil and Water Science*, 3(1):45-53
23. Seiyaboh EI, Izah SC, Bokolo JE (2017). Bacteriological quality of water from river nun at Amassoma Axes, Niger Delta, Nigeria. *ASIO Journal of Microbiology, Food Science & Biotechnological Innovations*, 3(1), 22 – 26.
24. Ben-Eledo VN, Kigigha LT, Izah SC, Eledo BO (2017). Bacteriological Quality Assessment of Epie Creek, Niger Delta Region of Nigeria. *International Journal of Ecotoxicology and Ecobiology*, 2(3): 102-108
25. Ben-Eledo VN, Kigigha LT, Izah SC, Eledo BO (2017). Water quality assessment of Epie creek in Yenagoa metropolis, Bayelsa state, Nigeria. *Archives of Current Research International*, 8(2): 1 – 24.
26. Seiyaboh EI, Izah SC, Oweibi S (2017). Assessment of Water quality from Sagbama Creek, Niger Delta, Nigeria. *Biotechnological Research*, 3(1):20-24
27. Seiyaboh EI, Inyang IR, Izah SC (2016). Seasonal Variation of Physico-Chemical Quality of Sediment from Ikoli Creek, Niger Delta. *International Journal of Innovative Environmental Studies Research*, 4(4): 29-34.
28. Seiyaboh EI, Inyang IR, Izah SC (2016b). Spatial Variation in Physico-chemical Characteristics of Sediment from Epie Creek, Bayelsa State, Nigeria. *Greener Journal of Environment Management and Public Safety*, 5(5): 100 - 105
29. Izah SC and Ineyougha ER (2015). A review of the microbial quality of potable water sources in Nigeria. *Journal of Advances in Biological and Basic Research*. 1(1): 12 - 19.
30. Ogamba EN, Charles EE, Izah SC (2021). Distributions, pollution evaluation and health risk of selected heavy metal in surface water of Taylor creek, Bayelsa State, Nigeria. *Toxicology and Environmental Health Sciences* 13(2):109 – 121. DOI: 10.1007/s13530-020-00076-0.
31. Ogamba EN, Seiyaboh EI, Izah SC, Ogbugo I, Demedongha FK (2015). Water quality, phytochemistry and proximate constituents of *Eichhornia crassipes* from Kolo creek, Niger Delta, Nigeria. *International Journal of Applied Research and Technology*, 4(9): 77 – 84.
32. Ogamba EN, Izah SC, Toikumo BP (2015). Water quality and levels of lead and mercury in *Eichhornia crassipes* from a tidal creek receiving abattoir effluent, in the Niger Delta, Nigeria. *Continental Journal of Environmental Science*, 9(1): 13 – 25.

33. Ogamba EN, Izah SC, Oribu T (2015). Water quality and proximate analysis of *Eichhornia crassipes* from River Nun, Amassoma Axis, Nigeria. *Research Journal of Phytomedicine*. 1(1): 43 – 48.
34. Seiyaboh EI, Youkparigha FO, Izah SC, Daniels ID (2020). Bacteriological Quality of Groundwater in Imiringi Town, Bayelsa State, Nigeria. *Journal of Biotechnology and Biomedical Science*, 2(2): 34 - 40
35. Izah SC, Richard G, Aseibai ER (2021). Public Health implications of Fungi-aerosol contamination around a major dumpsite in Bayelsa State, Nigeria. *Journal of Environmental Treatment Techniques*, 9(2): 458 – 462.
36. Seiyaboh EI, Izah SC (2020). Assessment of Microbial Characteristics of Processed Palm Weevil "*Rhynchophorus phoenicis*" Larvae sold in some Market Areas in Bayelsa State, Nigeria. *Journal of Advanced Research in Medical Science & Technology*; 7(1): 24-29.
37. Izah SC, Etebu EN, Aigberua AO, Odubo TC, Iniamagha I (2022). A meta-analysis of microbial contaminants in selected ready-to-eat foods in Bayelsa State, Nigeria: Public Health implications and risk-reduction strategies. *Hygiene and Environmental Health Advances*. 4:100017. <https://doi.org/10.1016/j.heha.2022.100017>.
38. Kigigha LT, Igoya UOS, Izah SC (2016). Microbiological Quality Assessment Of Unpeeled Groundnut Sold in Yenagoa Metropolis, Nigeria. *International Journal of Innovative Biochemistry & Microbiology Research*, 4(4):11- 22.
39. Pepper IL, Gerba CP (2005). *Environmental microbiology. A laboratory manual*. Second edition. Elsevier academic press.
40. Benson HJ (2002). *Microbiological Applications: Laboratory Manual in General Microbiology*. complete version, 5th edition. McGraw-Hill, New York.
41. Seiyaboh EI, Youkparigha FO, Izah SC, Mientei K (2020). Assessment of bacteriological characteristics of surface water of Taylor creek in Bayelsa state, Nigeria. *Noble International Journal of Scientific Research*, 4(4), 25-30
42. Aghoghovwia OA, Umoru OD, Izah SC (2018). Physicochemical characteristics of nun river at Gbarantoru and Tombia Axis in Bayelsa State, Nigeria. *Bioscience Methods*, 9(1): 1-11
43. Rusydi AF (2018). Correlation between conductivity and total dissolved solid in various type of water: a review. *IOP Conf. Series: Earth and Environmental Science* 118: 012019 doi :10.1088/1755-1315/118/1/012019.