

Efficacy of Arbuscular mycorrhizal inoculations on tree species germination and growth

Abstract

Researchers are very concerned about addressing or managing the growing problem of global warming or climate change. Global warming is caused by greenhouse gases, of which carbon dioxide (CO₂) alone contributes 60%. By absorbing carbon dioxide (CO₂) during photosynthesis and naturally storing it as biomass, trees serve as a sink for CO₂ because trees have the ability to store carbon and can lessen the effects of climate change and global warming, planting trees or restoring forests may be a solution to this growing problem. This review paper can be helpful since it discusses the interaction between arbuscular mycorrhizal fungi (AMF) and the roots of trees, which **benefits** the plant for better germination and growth. **dormancy** of seeds or a delay in growth can be identified as a disadvantage in a plantation program. With the hope that this information would advance the field of research, we have concentrated on mycorrhizal association and its impact on different species of trees in this work. It can be useful for tree species that have lower biomass or slower growth.

Keywords: Global warming, photosynthesis, arbuscular mycorrhizal fungi (AMF), Dormancy and Biomass.

Introduction

Seed dormancy, a physiological phenomenon in plants that can be brought on by internal or external stimuli, prevents seeds from sprouting even under ideal circumstances. Hard seed coats, underdeveloped embryos, primitive embryos, and inhibitory compounds can all contribute to seed dormancy (Mousavi et al., 2019). Some species' seeds cannot fully germinate because the embryo is restricted by the structures in its environment. Embryos removed from these seeds are not dormant; this phenomenon is known as seed coat augmented dormancy or physical dormancy. In seeds of other species with underdeveloped embryos, a second kind of dormancy known as morphophysiological dormancy is also present, but it also has a physiological component. As a result, these seeds need a treatment to break their dormancy, such as a specific mix of warm and/or cold stratification (Baskin & Baskin, 2004). Large-scale cultivation is hampered in forestry and home garden plantation initiatives by delayed nursery development and deprived seed germination (Alamgir and Hossain 2005b; Azad et al., 2006a & 2006b).

Bonner (1984) stated that germination is defined as “the resumption of active growth

in an embryo which results in its emergence from the seed and development of those structures essential for plant development”. It is the culminating event of seed maturation, the establishment of the seedling (Kramer and Kozlowski, 1979). Pre-sowing treatments of seed are crucial for constructing a nursery of a specific species in order to forecast the largest number of high-quality seedlings with the least amount of money, time, and labor (Das, 2015). To overcome this dormancy in forest seeds, various pre-sowing seed treatments are used.

One of the soil microorganisms that makes up the fundamental components of a good soil-plant connection is the arbuscular mycorrhizal fungus (AMF). By reducing stresses through symbiosis, **the** mycorrhizal fungus is an essential part of the soil microbial community that promotes plant growth and survival (Sylvia & Williams, 1992). The advantages that these fungi provide their host include improved phosphorus uptake (Goussous and Mohammad, 2009), increased nitrogen absorption (Rotor and Delima, 2010), the production of plant growth hormones (Herrera-Medina et al., 2007), root defense against soil-borne diseases (Bakhtiar et al., 2010), and improved plant growth and productivity (Duponnois et al., 2005).

The ability of the AM fungus to colonize roots and act as a biofertilizer and bioprotectant to protect plants against parasitic nematodes and fungi as well as to boost plant development and yield has been demonstrated (Berruti et al., 2016). (Wei et al., 2016). AMF is a form of symbiotic soil fungi that colonizes the roots of about 80% of vascular plants, according to Vierheilig (2004). It is one of the essential soil microbes that contribute to a balanced soil-plant system (Budi et al., 2012). They improve plant fitness and soil quality (Barea et al., 2002). Since it has been demonstrated that the AMF colonizes its roots in prodigious quantities, the neem tree species is extremely dependent on mycorrhizal fungus (Habte et al., 1993). In order to lessen the negative effects of slow growth and low germination **rates** in tree species, this review will analyze the benefits, drawbacks, and role of seed treatment with arbuscular mycorrhizal fungi. It will also consider potential AMF that may be applied through seed treatment or inoculation for better growth.

Effect of Mycorrhiza on growth and germination of different tree species

Although arbuscular mycorrhiza fungi **are** naturally present in the roots of higher tree species, this literature includes **the** effect of some AM fungi on **the** growth and germination of tree species (Table 1). Khaiper *et al.* (2021) conducted an experiment on *Melia azedarach*, in which she treated seeds with different pre-sowing treatments and then sow **these** seeds into soils inoculated with *Glomus mosseae* and reported that seeds showed maximum seed

germination percentage in seeds treated with cow dung slurry for 6 days within the soil inoculated with *Glomus mosseae* as compared to control. Shukla *et al.* (2012) investigated the effect of AM fungi (*Glomus cerebriforme* and *Acaulospora scrobiculata*) on two multipurpose tree species *i.e.*, *Eucalyptus tereticornis* and *Albizia procera* and found that shoot length, dry weight and nutrient uptake increased significantly after inoculations with AM fungi. The best results were obtained with *G. cerebriforme* in both the tree species than other inoculants.

Muthukumar *et al.* (2013) concluded that the seedlings of *Casuarina equisetifolia* inoculated with *Glomus aggregatum* and *Acaulospora scrobiculata* had significantly higher plant growth and nutrient parameters measured. Nevertheless, the response was higher for seedlings inoculated with *G. aggregatum* compared to those inoculated with *A. scrobiculata*.

Basumatary *et al.* (2014) reported that an increase in length, diameter, circumference and biomass yield along with nutrient availability in the rhizosphere *i.e.* Nitrogen, Phosphorus, Potassium and carbon in Rubber tree (*Hevea brasiliensis*) seedlings inoculation with *Acaulospora* sp. and *Glomus* sp. over control.

Singh *et al.* (2014) revealed that the maximum biomass accumulation, phosphorous and nitrogen content in *Acacia nilotica* seedlings were achieved under soil treated with *Glomus mosseae* along with phosphate and nitrogenous fertilizers.

Reena and Bagyaraj (1990) conducted an experiment on effect of 13 different VAM fungi on the seedlings of *Acacia nilotica* and *Calliandra calothyrsus* and reported that inoculated plants had greater plant height, leaf number, stem girth, biomass and phosphorus and zinc content. They also had more mycorrhizal root colonization, spores and external hyphae in soil. *A. nilotica* seedlings responded best at inoculation with *G. mosseae* followed by *G. caledonicum* whereas, *C. calothyrsus* responded best with *G. velum* and *G. merredum*.

Herrmann *et al.* (2016) revealed that the management of Arbuscular mycorrhizal fungi (AMF) would contribute to maintaining or restoring soil fertility, leading to a better tree growth and optimized latex yield of Rubber trees (*Hevea brasiliensis*).

Chen *et al.* (2017) reported that Liquirice (*Glycyrrhiza uralensis*) plant inoculated with *G. mosseae* improved the features of the root system and increased in photosynthetic efficiency. The uptake of P and K in plant was also increased when inoculated with *G. mosseae*.

Filho *et al.* (2017) found that the application of *Rhizofagus clarum* and *Glomus etunicatum* in soil grown with star fruit (*Averrhoa carrambola* L), provided increments of

49% in height, 99% in dry matter production and 86, 129 and 108% in the content of N, K and calcium respectively, in relation to the control.

Jamaluddin and Shukla (2012) conducted a laboratory experiment with three treatments viz., AM fungi (*Glomus mosseae*, *Acaulospora* sp. and *Gigaspora* sp.) + root exudates, AM fungi alone and root exudates alone. Out of which the treatment containing AM fungi + root exudates showed maximum colonization, phosphatase enzyme activity, phosphorus uptake and leaf protein as compared with control and other treatments under studied.

Hamidi *et al.* (2017) studied the effect of endomycorrhizae *i.e.* *Glomus etunicatum* on the seedlings of cork oak plants (*Quercus suber*) and observed that plants inoculated with mycorrhizae showed the stimulatory effect on aerial part, aerial part weight, average root length and average fresh weight and the leaves number as compared to control.

Jha *et al.* (2012) studied the effect of AM fungi namely *Glomus fasciculatum*, *Glomus aggregatum*, *Glomus cerebriforme*, *Glomus diaphanum* and *Glomus etunicatum* on *Dalbergia sissoo*, *Acacia procera* and *Acacia nilotica* seedlings and revealed that the application of AM fungi improves the uptake of phosphorus, minor elements and water and enhances plant growth and resistance to root diseases as compared to non-mycorrhizal plants.

An experiment conducted by Young (1990) showed the effect of single and mixed inoculations with phosphorus solubilizing bacteria and VAM fungi on the growth of *Leucaena leucocephala*, *Acacia confuse*, *A. mangium* and *Liquidamber formosana*. Growth of *L. leucocephala* tree increased by 22-99% with VAM fungal inoculation. The growth of *Acacia confuse* (14-63%), *A. mangium* (7-88%) and *Liquidamber formosana* (24-280%) promoted by mixed inoculants.

Sumana and Bagyaraj (1999) conducted a experiment and used eight different VAM fungi (*Acaulospora laevis*, *Gigaspora margarita*, *Glomus caleldonicum*, *Glomus fasciculatum*, *Glomus intraradices*, *Glomus leptotichum*, *Glomus macrocarpum* and *Glomus mosseae*) for selecting best symbiont for *Dalbergia latifolia* and reported that inoculated plants had greater plant height, stem girth, dry weight and phosphorus content when compared to non-inoculated plants. *D. latifolia* responded best to inoculation with *Glomus leptotichum* as well as *Glomus fasciculatum* as compared to the other AM fungi.

Youpensuk *et al.* (2005) conducted an experiment on Seedlings of *Macaranga denticulate* inoculated with spores of fungi (*Glomus* species, *Glomus fasciculatum*, *Acaulospora* species and mixed species of AM fungi) in pot and concluded that Nutrient contents (N, P and K) of plant inoculated with *Acaulospora* species or mixed species of AM fungi were higher than plants inoculated with *Glomus* spp. and *G. fasciculatum*.

Table 1: Effect of mycorrhizal treatment on germination and growth of different tree species

S. No.	Tree spp.	Mycorrhizal spp.	Description	reference
1.	<i>Melia azedarach</i>	<i>Glomus mosseae</i>	The germination percentage (75.87) and other such as mean daily germination, germination value and speed of germination were recorded highest in treatment with cow dung slurry for 6 days + <i>Glomus mosseae</i> .	Khaiper <i>et al.</i> (2021)
2.	<i>Eucalyptus tereticornis</i> and <i>Albizia procera</i>	<i>Glomus cerebriforme</i> and <i>Acaulospora scrobiculata</i>	Shoot length, dry weight and nutrient uptake increased significantly after inoculations with AM fungi. Best results were obtained with <i>G. cerebriforme</i> in both the tree species then other inoculants	Shukla <i>et al.</i> (2012)
3.	<i>Azadirachta indica</i>	<i>G. intraradices</i> and <i>G. gegosporum</i>	Neem (<i>Azadirachta indica</i>) seedlings and inoculate with <i>G. intraradices</i> and <i>G. gegosporum</i> have increase mycorrhizal colonization, greater plant height, leaf area and number of root, biomass, phosphorus, nitrogen, collar diameter and potassium content and seedling quality.	Muthukumar <i>et al.</i> (2001)
4.	<i>Casuarina equisetifolia</i>	<i>Glomus aggregatum</i> and <i>Acaulospora scrobiculata</i>	Higher plant growth and nutrient parameters measured. Nevertheless, the response was higher for seedlings inoculated with <i>G. aggregatum</i> compared to those inoculated with <i>A. scrobiculata</i> .	Muthukumar <i>et al.</i> (2013)

5.	<i>Hevea brasiliensis</i>	<i>Acaulospora</i> sp. and <i>Glomus</i> sp.	Increase in length, diameter, circumference and biomass yield along with nutrient's availability in rhizosphere <i>i.e.</i> Nitrogen, Phosphorus, Potassium and carbon in seedlings inoculated with <i>Acaulospora</i> sp. and <i>Glomus</i> sp. over control.	Basumatary <i>et al.</i> (2014)
6.	<i>Acacia nilotica</i>	<i>Glomus mosseae</i>	Maximum biomass accumulation, phosphorous and nitrogen content in seedlings were achieved under soil treated with <i>Glomus mosseae</i> along with phosphate and nitrogenous fertilizers.	Singh <i>et al.</i> (2014)
7.	<i>Hevea brasiliensis</i>	Arbuscular mycorrhizal fungi	Arbuscular mycorrhizal fungi (AMF) would contribute to maintaining or restoring soil fertility, leading to a better tree growth and optimized latex yield of Rubber trees (<i>Hevea brasiliensis</i>).	Herrmann <i>et al.</i> (2016)
8.	<i>Glycyrrhiza uralensis</i>	<i>G. mosseae</i>	Plant inoculated with <i>G. mosseae</i> improved the features of the root system and increased in photosynthetic efficiency. The uptake of P and K in plant was also increased when inoculated with <i>G. mosseae</i> .	Chen <i>et al.</i> (2017)
9.	<i>Averrhoa carrambola</i> L	<i>Rhizofagus clarum</i> and <i>Glomus etunicatum</i>	Increments of 49% in height, 99% in dry matter production and 86, 129 and 108% in the content of N, K and calcium respectively, in relation to the control.	Filho <i>et al.</i> (2017)
10.	<i>Quercus</i>	<i>Glomus</i>	Plants inoculated with mycorrhizae	Hamidi <i>et al.</i>

	<i>suber</i>	<i>etunicatum</i>	showed the stimulatory effect on aerial part, aerial part weight, average root length and average fresh weight and the leaves number as compared to control.	(2017)
--	--------------	-------------------	--	--------

Conclusion: Trees act as a sink for carbon dioxide (CO₂) by absorbing it during photosynthesis and naturally storing it as biomass, which helps to mitigate the consequences of climate change and global warming. The relationship that develops between arbuscular mycorrhizal fungus (AMF) and tree roots, enabling the plant to germinate and grow more successfully. A disadvantage in a plantation program can be the dormancy of seeds or a delay in growth. We have focused on mycorrhizal association and its effects on several tree species in this work with the aim that this information would progress the field of research for tree species with lower biomass or slower growth.

References:

1. Alamgir M, Hossain MK. Effect of pre-sowing treatments on germination and initial seedling development of *Albizia saman* in the nursery. *Journal of Forestry Research*. 2005;16:200-204.
2. Azad MS, Islam MW, Matin MA, Bari MA. Effect of pre-sowing treatment on seed germination of *Albizia lebbeck* (L.) Benth. *South Asian Journal of Agriculture*. 2006;12:32-34.
3. Azad MS, Matin MA, Islam MW. Effect of pre-sowing treatment on seed germination of Lohakath (*Xylia kerrii* Craib & Hutch). *Khulna University Studies*. 2006; 22:33-36.
4. Bakhtiar Y, Yahya S, Sumaryono W, Sinaga MS, Budi SW, Tajuddin T. Isolation and identification of Mycorrhizosphere bacteria and their antagonistic effects towards *Ganoderma boninense* in vitro. *Microbiology Indonesia*. 2010;4:9-9.
5. Barea JM, Azcón R and Azcon-Aguilar C. Mycorrhizosphere interactions to improve plant fitness and soil quality. *Antonie van Leeuwenhoek*. 2002;81(1-4): 343-351.
6. Baskin CC, Baskin JM, Guerrant EO, Havens K, Maunder M. Determining dormancy-breaking and germination requirements from the fewest seeds. *Ex situ plant conservation: supporting species survival in the wild*. 2004;162-179.
7. Basumatary N, Parkash V, Tamuli AK, Saikia AJ, Teron R. Arbuscular mycorrhizal inoculation affects growth and rhizospheric nutrient availability in *Hevea brasiliensis*

- (Willd. ex A. Juss.) Mull. Arg. clones. International Journal of Current Biotechnology. 2014;2:14-21.
8. Berruti A, Lumini E, Balestrini R, Bianciotto V. Arbuscular mycorrhizal fungi as natural biofertilizers: let's benefit from past successes. *Frontiers in microbiology*. 2016;6:1559.
 9. Bonner FT. Glossary of seed germination terms for tree seed workers. US Department of Agriculture, Forest Service, Southern Forest Experiment Station; 1984.
 10. Budi SW, Bakhtiar Y, May NL. Bacteria associated with arbuscular mycorrhizal spores *Gigaspora margarita* and their potential for stimulating root mycorrhizal colonization and neem (*Melia azedarach* Linn) seedling growth. *Microbiology Indonesia*. 2012;6:180-188.
 11. Chen M, Yang G, Sheng Y, Li P, Qiu H, Zhou X, Huang L, Chao Z. *Glomus mosseae* inoculation improves the root system architecture, photosynthetic efficiency and flavonoids accumulation of liquorice under nutrient stress. *Frontiers in plant science*. 2017;8:1-10.
 12. Das N. The effect of different pre-sowing treatments on the germination of *Aquilaria agallocha* and *Shorea robusta* seeds in the nursery. *Indian Forester*. 2015;141:285-292.
 13. Duponnois R, Colombet A, Hien V, Thioulouse J. The mycorrhizal fungus *Glomus intraradices* and rock phosphate amendment influence plant growth and microbial activity in the rhizosphere of *Acacia holosericea*. *Soil biology and Biochemistry*. 2005;37:1460-1468.
 14. Filho, J. A. V., Mendonça, F. M. S., Martins, M. A., dos, S. P. C. and Cordeiro, C. A. J. . Arbuscular mycorrhizal fungi and phosphate fertilization on star fruit tree seedlings. *Revista Brasileira de Ciências Agrárias* 2017;12:14-19.
 15. Goussous SJ, Mohammad MJ. Comparative effect of two arbuscular mycorrhizae and N and P fertilizers on growth and nutrient uptake of onions. *International Journal of Agriculture and Biology*. 2009;11:463-467.
 16. Habte M, Muruleedhara BN, Ikawa H. Response of neem (*Azadirachta indica*) to soil P concentration and mycorrhizal colonization. *Arid Land Research and Management*. 1993;7:327-333.
 17. Hamidi O, Talbi Z, Chliyeh M, Touhami AO, Selmaoui K, Benkirane R, Douira A. Effect of Endomycorrhizal Inoculation on the Young Cork Oak Plants (*Quercus suber*) Growth. *Annual Research & Review in Biology*. 2017;13:1-11.

18. Herrera-Medina MJ, Steinkellner S, Vierheilig H, Ocampo Bote JA, Garcia Garrido J. Abscisic acid determines arbuscule development and functionality in the tomato arbuscular mycorrhiza. *New Phytologist*. 2007;175:554-564.
19. Herrmann L, Bräu L, Robin A, Robain H, Wiriyakitnatekul W, Lesueur D. High colonization by native arbuscular mycorrhizal fungi (AMF) of rubber trees in small-holder plantations on low fertility soils in North East Thailand. *Archives of Agronomy and Soil Science*. 2016;62:1041-1048.
20. Jamaluddin M, and Shukla R. Effect of root exudates on colonisation of am fungi in *Jatropha curcas*. *Indian Forester*. 2012;138:113-115.
21. Jha A, Kumar A, Shukla A, Chakravarty N. Response of four multipurpose tree species to arbuscular mycorrhizal inoculations. *Indian Phytopathology*. 2012;65:297.
22. Khaiper, M., Dhanda, S. K., Ahlawat, K. S., Chugh, R. S., Jangra, M. and Verma, P. Effect of pre-sowing treatments on drupe germination of *Melia azedarach* at nursery stage. *Agricultural Mechanization in Asia, Africa and Latin America*. 2021; 52:4569-4577.
23. Kramer P, and Kozłowski T. Physiology of woody plants. *New York Academic Press*, 1979; 811.
24. Mousavi L, Wan Ishak WR, Mousavi M. Evaluation of physicochemical methods for dormancy breakage and germination of *Datura stramonium* seeds. *Journal of Chemical Health Risks*. 2019;9:217-224.
25. Muthukumar T, Udaiyan K, Rajeshkannan V. Response of neem (*Azadirachta indica* A. Juss) to indigenous arbuscular mycorrhizal fungi, phosphate-solubilizing and asymbiotic nitrogen-fixing bacteria under tropical nursery conditions. *Biology and Fertility of Soils*. 2001;34:417-426.
26. Muthukumar T, Uma E, Priyadharsini P. Arbuscular Mycorrhizal Fungal Strains and Soil Type Influence Growth, Nodulation, and Nutrient Uptake of *Casuarina equisetifolia*. *Microbiological Research In Agroecosystem Management* 2013;35-52.
27. Reena J, Bagyaraj DJ. Response of *Acacia nilotica* and *Calliandra calothyrsus* to different VA mycorrhizal fungi. *Arid Land Research and Management*. 1990;4:261-268.
28. Rotor AV, Delima PC. Mycorrhizal association, N fertilization and biocide application on the efficacy of bio-N on Corn (*Zea mays* L.) growth and productivity. *International Journal of Scientometric Research* 2010;2:267-290.

29. Shukla A, Kumar A, Jha A, Rao DV. Phosphorus threshold for arbuscular mycorrhizal colonization of crops and tree seedlings. *Biology and Fertility of Soils*. 2012;48:109-116.
30. Singh KP, Yadav R, Kumari P, Bhadauria S. Biomass production, phosphorus and nitrogen content in *Acacia nilotica* under am fungi and nutrient treated soil. *Indian Forester*. 2014;140:489-493.
31. Sumana DA, Bagyaraj DJ. Selection of efficient VA-mycorrhizal fungi for inoculating Neem. In *Proceedings of the National conference of Mycorrhiza*. Singh S (ed) Barkatulla University, Bhopal 1999;10.
32. Sylvia DM, Williams SE. Vesicular-arbuscular mycorrhizae and environmental stress. *Mycorrhizae in sustainable agriculture*. 1992;54:101-124.
33. Vierheilig, H. Regulatory mechanisms during the plant-arbuscular mycorrhizal fungus interaction. *Canadian Journal of Botany*. 2004; 82:1166-1176.
34. Young CC. Effects of phosphorus-solubilizing bacteria and vesicular-arbuscular mycorrhizal fungi on the growth of tree species in subtropical-tropical soils. *Soil Science and Plant Nutrition*. 1990;36:225-231.
35. Youpensuk S, Rerkasem B, Dell B, Lumyong S. Effects of arbuscular mycorrhizal fungi on a fallow enriching tree (*Macaranga denticulata*). *Fungal Diversity*. 2005;18:189-199.