

Original Research Article

BIOLOGICAL MANAGEMENT OF FUSARIUM LENTIL WILT THROUGH EXPLORING THE POTENTIAL OF PSEUDOMONAD ISOLATES

Abstract

Lentil (*Lens culinaris*), also known as "masoor," is a crucial pulse crop in the *Leguminosae* family known for their high protein and nutrient content. Wilt incited by *Fusarium oxysporum* f.sp. *lentis* can result in significant yield losses for lentil farmers. Chemical management through fungicides can reduce crop losses but has negative effects on the environment and human health. To overcome this, use of plant growth-promoting rhizobacteria (PGPR) as a bio-inoculant is gaining attention as a sustainable approach to manage plant diseases. PGPR are soil bacteria that live in the rhizosphere of plants and promote growth. The use of PGPR-based agents has increased in agriculture in recent years. In present investigation, isolation and evaluation of novel pseudomonad isolates were performed from various soil sources for controlling lentil wilt caused by *Fusarium oxysporum* f.sp. *lentis* and assess their ability to inhibit the pathogen *in vitro*.

Keywords: Lentil (*Lens culinaris*), *Fusarium oxysporum* f.sp. *lentis*, Lentil wilt Plant growth-promoting rhizobacteria (PGPR)

Introduction

The legume crop is unique in world because of its high protein content (15-35%), fiber, minerals, carbohydrates, and nutrients (Nadia *et al.*, 2019). In the family, Leguminosae (*Fabaceae*), lentils (*Lens culinaris* M.) are a deployed species (2n =14) self-pollinating crop (Dubey and Pandey, 2020). Wilt of lentil is incited by *Fusarium oxysporum* Schlecht. emend. Snyder & Hansen f. sp. *lentis* Vasudeva and Srinivasan (Fol) (Belabid and Fortas 2002). It brings about significant economic losses for lentil farmers in India and other nations depending on the severity and stage (pre-podding to pre-harvest) of wilt, yield loss could reach 0-100% (Dubey, and Singh, 2018; Tiwari *et al.*, 2018). Pathogen can also survive within soil as chlamydospores, which can persist for several years (Bayaa *et al.*, 1997). It has been shown that chemical-based management (fungicides) has the potential to reduce crop losses caused by plant-pathogenic organisms. However, due to the unfortunate and negligent application of synthetic pesticides, there can be

Comment [H1]: Not mentioned in reference

phytotoxicity and fungicidal residues that are harmful to the environment (natural enemies, flora and fauna) as well as human health (Aioub *et al.*, 2022; Eldeeb *et al.*, 2022). Use of plant growth-promoting rhizobacteria (PGPR) as a bio-inoculant is gaining attention as a sustainable approach to manage plant diseases and overcome the negative effects of fungicides. This is supported by recent research (Backer *et al.*, 2018; Ashok *et al.*, 2015; Turan *et al.*, 2021). PGPR are soil bacteria that live in rhizosphere of plants and promote growth by various mechanisms. They can grow on, in, or around plant tissues (Godbole *et al.*, 2021). For the past decade, use of PGPR-based agents (*Pseudomonas* spp.) as seed bio-inoculant, soil amendment, or soil drenching in crop production systems has increased in agriculture (Abbouni *et al.*, 2018). Due to PGPRs potential to manage soil-borne pathogens by colonizing plant roots (Hisamuddin *et al.*, 2012) and detoxifying the environment, they were considered a suitable solution for biological control (Kumar *et al.*, 2018; Abbouni *et al.*, 2018; Adhikari *et al.*, 2021). The study aimed to isolate and evaluate PGPR from various soil sources for controlling lentil wilt. It was done under *in vitro* conditions and PGPRs were screened for their ability to inhibit pathogens and improve the environment.

Comment [H2]: Not mentioned in reference

Comment [H3]: Not mentioned in reference

Comment [H4]: Not mentioned in reference

Materials and Methods

Isolation and purification of pathogen

F. oxysporum was isolated and purified from wilt infected lentil plants. Samples were collected, surface sterilized, and transferred onto PDA medium (Mondal *et al.*, 2021). Purification was done by growing hyphal tips and the pathogen was identified based on morphological characteristics. It confirmed Koch's postulate. The pathogen was maintained in refrigerated subcultures.

Rhizospheric soil samples collection and isolation of PGPR.

Rhizosphere soil was collected from different crops, and a bacterial suspension was obtained by shaking 1g of soil in sterilized water (Godbole *et al.*, 2021). Bacterial isolation was done using KMB medium with benomyl and by dilution plate technique. Colonies of *Pseudomonas* spp. were purified and kept at 4°C. The isolates were evaluated for plant growth promotion and antagonistic potential (Tsegaye *et al.*, 2019).

Evaluation of potential PGPR as antagonists against *F. oxysporum* for lentil wilt management

The potential of Pseudomonads spp. as antagonists against *Fusarium oxysporum* f.sp. *lentis* was evaluated in vitro by streaking bacterial isolates around pathogen in KMB medium and incubating at 28°C. Percent inhibition of mycelial growth was calculated using formula $I = (C - T)/C \times 100$ (Kalantari *et al.*, 2018).

Comment [H5]: Fol

Results and Discussion

Screening of native Pseudomonad isolates against *F. oxysporum* pathogen

Dual culture test revealed that all PGPRs (*Pseudomonad* isolates) have potential in controlling mycelium growth of *F. oxysporum* except isolate PGPR4 (Fig.1 & Table 1). In this present study, out of 20 Pseudomonad isolates, PGP 18 exhibited highest percent mycelial growth inhibition (67.41%) followed by PGP 6, 16 and 17 (61.85, 61.48 and 59.63) respectively. Mycelial growth inhibition was 57.78%, 55.56% and 51.11% in isolates PGP 11, 15 and 20 respectively. Lowest percent of mycelial growth inhibition was exhibited by PGP 4 (0.00 %) followed by PGP 5, 1 and 2 (25.93, 29.26% and 30.37%) respectively as compared to control. Results of present study indicate that various PGPR pseudomonad isolates showed varying levels of inhibition against *F. oxysporum*, likely due to their varying antifungal abilities. The results of our study are partially or fully aligning with the findings those of Harsha *et al.* (2023), who evaluated 20 inherent *Pseudomonas* bacteria for their ability to combat *F. oxysporum* and found that isolate CRS-PF1 was particularly effective, exhibiting 51.84% inhibition of mycelium growth. The pseudomonad isolate PGPR-WS demonstrated strong inhibition against *F. oxysporum* in chickpeas, resulting in 75% reduction in growth of mycelia as compared to the control (Kumar *et al.*, 2018). Similarly, twenty-four strains of *Bacillus* (B4, B7 and B12) showed excellent antagonistic activity against investigated pathogenic fungi (Abbouni, *et al.*, 2018). Experimental findings were also in accordance with previous results (Sachdev and Singh, 2018) who identified and characterized *Pseudomonas aeruginosa* with positive antagonistic potentials toward *Fusarium lycopersici* and demonstrated radial inhibitions 67.85%. Isolation and antagonistic activity of rhizospheric bacteria against various disease-causing pathogens were also reported previously (Sahu *et al.*, 2017).

Table.1: Screening of native PGPRs (Pseudomonad isolates) against *F. oxysporum* f.sp. *lentis* (Fol) pathogen

Sl. No.	Pseudomonad isolates	Colony diameter *(mm)	Percentage inhibition of radial growth (PIRG)* (Fol)
1.	PGP1	63.67	29.26
2.	PGP2	62.67	30.37
3.	PGP3	60.67	32.59
4.	PGP4	90.00	0.00
5.	PGP5	66.67	25.93
6.	PGP6	34.33	61.85
7.	PGP7	52.00	42.22
8.	PGP8	48.33	46.30
9.	PGP9	48.00	46.67
10.	PGP10	46.00	48.89
11.	PGP11	38.00	57.78
12.	PGP12	50.67	43.70
13.	PGP13	46.00	48.89
14.	PGP14	46.00	48.89
15.	PGP15	40.00	55.56
16.	PGP16	34.67	61.48
17.	PGP17	36.33	59.63
18.	PGP18	29.33	67.41
19.	PGP19	47.67	47.04
20.	PGP20	44.00	51.11
	Control (KBs)	90.00	
	C.D.	3.84	3.32
	SEm (±)	1.34	1.15
	C.V.	4.68	4.26

*Mean values of three replication




Fig. 1: Screening of native PGPRs (Pseudomonad isolates) against *F. oxysporum* f.sp. *lentis* (Fol) pathogen

Conclusion

The dual culture test demonstrated that most of the Pseudomonad isolates (PGPRs) were capable of inhibiting the growth of *F. oxysporum* f.sp. *lentis* (Fol), except for isolate PGPR4 (as shown in Fig.1 & Table 1). In this study, out of 20 PGPR isolates, PGP18 was found to be the most effective in inhibiting mycelial growth with 67.41% inhibition. PGP6, PGP16, and PGP17 also showed significant inhibition with 61.85%, 61.48%, and 59.63% respectively. PGP11, PGP15, and PGP20 showed moderate inhibition with 57.78%, 55.56%, and 51.11% respectively. Conversely, PGPR4 exhibited the least inhibition with 0.00% compared to the control. The results suggest that the varying levels of inhibition displayed by the PGPR pseudomonads against *F. oxysporum* f.sp. *lentis* could be attributed to their differing antifungal abilities.

Comment [H6]: delete

References

1. Abbouni, B., Ghanem, M., Benine, M. L., Labdi, M., Reguig, M., Benali, M., &. (2018). Isolation, Screening, Characterization of PGPR of Lentils *Lens culinaris*. *Der Pharmacia Lettre*, 10 (1), 91-104
2. Adhikari, P., Jain, R., Sharma, A., & Pandey, A. (2021). Plant growth promotion at low temperature by phosphate-solubilizing *Pseudomonas* spp. isolated from high-altitude Himalayan soil. *Microbial ecology*, 82(3), 677-687.
3. Babalola, O. O., Emmanuel, O. C., Adeleke, B. S., Odelade, K. A., Nwachukwu, B. C., Ayiti, O. E., & Igiehon, N. O. (2021). Rhizosphere microbiome cooperations: Strategies for sustainable crop production. *Current Microbiology*, 78(4), 1069-1085.
4. Belabid, L., & Fortas, Z. (2002). Virulence and vegetative compatibility of Algerian isolates of *Fusarium oxysporum* f. sp. lentis. *Phytopathologia Mediterranea*, 41(3), 179-187.
5. Blanco-Vargas, A., Rodríguez-Gacha, L. M., Sánchez-Castro, N., Garzón-Jaramillo, R., Pedroza-Camacho, L. D., Poutou-Piñales, R. A., ... & Pedroza-Rodríguez, A. M. (2020). Phosphate-solubilizing *Pseudomonas* sp., and *Serratia* sp., co-culture for *Allium cepa* L. growth promotion. *Heliyon*, 6(10), e05218.
6. Dubey, K. and Pandey, V. (2020). Infection, disease and management of Fusarium wilt of lentil (*Fusarium oxysporum* f. Sp. Lentis). *International Journal of Creative Research Thoughts*, 8(3), 2638-2648.
7. Dubey, K. and Singh, S. K. (2018) Study Cultural, Morphological and Pathogenic Variation among Different Isolates of *Fusarium oxysporum* f. sp. lentis. *Int.J.Curr.Microbiol.App.Sci*, 7(9), 170-175.
8. Eldeeb, A. M., Farag, A. A. G., Al-Harbi, M. S., Kesba, H., Sayed, S., Elesawy, A. E., ... & Aioub, A. A. (2022). Controlling of *Meloidogyne incognita* (Tylenchida: Heteroderidae) using nematicides, *Linum usitatissimum* extract and certain organic acids on four peppers cultivars under greenhouse conditions. *Saudi Journal of Biological Sciences*, 29(5), 3107-3113.
9. Godbole, S., Sambherao, A., & Choubey, S. (2021). Isolation and characterization of plant growth promoting rhizobacteria (PGPR) from different crops in junnar tehsil. *International Journal of Recent Scientific Research*, 12, (03), 41186-41190.
10. Habib, S. H., Kausar, H., Saud, H. M., Ismail, M. R., & Othman, R. (2016). Molecular characterization of stress tolerant plant growth promoting rhizobacteria (PGPR) for growth enhancement of rice. *Int. J. Agric. Biol*, 18, 184-191.
11. Hisamuddin, A. A., Robab, M.I. and Sharf, R. A. (2012). Plant growth promoting Rhizobacteria: An overview. *J. Nat. Prod. Plant Resour.* 2 (1), 19-31.
12. Kalantari, S., Marefat, A., Naseri, B., & Hemmati, R. (2018). Improvement of bean yield and Fusarium root rot biocontrol using mixtures of *Bacillus*, *Pseudomonas* and *Rhizobium*. *Tropical Plant Pathology*, 43(6), 499-505.
13. Kumar, S., Sahni, S., Kumar, B., & Prasad, S. S. (2018). Multifarious antagonistic potentials of native *Pseudomonad* isolate from rhizosphere as biocontrol agents for the management of chickpea wilt. *CJAST*, 31, 1-13.

14. Mondal, A., Mahapatra, S., Chakraborty, S., Debnath, D., Das, T., & Samanta, M. (2021). Eco-friendly Management of Collar Rot of Lentil by Introduced Native Rhizobacterial Candidates. *Indian Journal of Agricultural Research*, 1(8).
15. Muehlbauer, G. J., Staswick, P. E., Specht, J. E., Graef, G. L., Shoemaker, R. C., & Keim, P. (1991). RFLP mapping using near-isogenic lines in the soybean [*Glycine max* (L.) Merr.]. *Theoretical and Applied Genetics*, 81(2), 189-198.
16. Nadia, B., Fatima, G., Rachid, M., & Mona, T. (2019). The genetic potential of moroccan lentil landraces. *International Journal of Agricultural Science and Research*, 9(3), 291-306.
17. Sachdev, S., & Singh, R. P. (2018). Isolation, characterisation and screening of native microbial isolates for biocontrol of fungal pathogens of tomato. *Clim Change Environ Sustain*, 6(1), 46-58.
18. Sahu, S., Sahu, M. K., Tiwari, R. K. S., & Khare, N. (2017). In-vitro Evaluation of Antagonistic Potential of Fluorescent *Pseudomonas* and *Trichoderma* against *Fusarium oxysporum* f. sp. *udum* inciting wilt in Pigeonpea (*Cajanus cajan* L. Mill. Sp). *Research Journal of Agricultural Sciences*, 8(4), 832-837.
19. Tiwari, N., Ahmed, S., & Sarker, A. (2018). Fusarium wilt: a killer disease of lentil. *Fusarium-Plant Diseases, Pathogen Diversity, Genetic Diversity, Resistance and Molecular Markers*.119-139. <http://dx.doi.org/10.5772/intechopen.72508>
20. Tsegaye, Z., Gizaw, B., Tefera, G., Feleke, A., Chaniyalew, S., Alemu, T., & Assefa, F. (2019). Isolation and biochemical characterization of Plant Growth Promoting (PGP) bacteria colonizing the rhizosphere of Tef crop during the seedling stage. *J. Plant Sci. Phytopathol*, 3(1), 013-027.
21. Turan, M., Arjumend, T., Argin, S., Yıldırım, E., Katırcıoğlu, H., Gürkan, B., & Bolouri, P. (2021). Plant root enhancement by plant growth promoting rhizobacteria. *Plant Roots*, 2021.
22. Harsha, M. K., Daunde, A. T., Bhalerao, P. B., & Sakhare, S. S. (2023). In vitro evaluation of native bacterial antagonists against *Fusarium oxysporum* f. sp. *ciceri* causing wilt of chickpea.

Comment [H7]: Transfer to No.10