

Original Research Article

Effect of nitrogen and phosphorus fertilizer doses on agro morphological parameters of wheat varieties in vertisol region in India

1. ABSTRACT

Wheat is the most important food grain, with more than a third of the world's population eating it as a staple diet for the optimum use of different nutrient. A field experiment was conducted in ICAR-IISS Bhopal. In this experiment we studied the different morphological parameters in nine wheat varieties i.e. plant height, plants dry weight, biological yield, grain yield. Some wheat varieties perform well in limited nutrient condition and some perform well in optimum nutrient doses condition. So those varieties which are perform well in nutrient limited condition grown efficiently in nutrient limited area of different region in India.

Keywords: Wheat, staple diet, soil fertility, morphological parameters

1. INTRODUCTION

Nitrogen is the fourth most prevalent element in living creatures, and it is utilized to make essential biological components including amino acids and nucleic acids (Schuur, 2011).

Phosphorus (P) is one of the most important nutrients for plant growth and survival. It is essential for cellular bioenergetics and metabolic pathways within the plant body. The primary function of mineral fertilizers is to increase crop yields, but the biggest impediment to realizing known crop potential is the low use of fertilizers, notably P and N (Irfan, 2018).

Batte (1992) also reported that increased nitrogen fertilisation rates significantly increased total dry matter production, leaf area index, plant height, and yield components (number of effective tillers m⁻², spikelets spike⁻¹, grains spike⁻¹, thousand grain weight, and grain yield) in wheat. It was determined that a two-split application of 90 kg ha⁻¹ of N and 20 kg ha⁻¹ of N at crop emergence can meet the wheat crop's N requirements and, as a result, appropriate growth and development was ensured, resulting in increased wheat grain production.

Kaur (2015) conducted two-year field experiment to screen twelve wheat genotypes for optimal nitrogen usage efficiency under four levels of nitrogen management (recommended dose of nitrogen; RDN, RDN-50 percent, RDN-25 percent, and RDN+25 percent). The result indicated that plant height, tiller number, spikelet number, grain yield, thousand grain weight, and biomass all increased with increase in the nitrogen dose).

Wheat growth, N uptake, and yield characteristics were all significantly greater with split (two or three) applications of N than with a single application (Bhardwaj, 2009).

Comment [W11]: 1. No Numbered
2. Describe the purpose of the work, techniques and methods used, major findings with important data and conclusions

Comment [W12]: In the text, citations should be indicated by the reference number in brackets

Comment [W13]: Irfan et al., 2018
More than 1 author

Comment [W14]: 1.not listed in the reference
2.use the latest reference, please

Comment [W15]: Kaur et al., 2015
More than 1 author

Comment [W16]: 1.Bhardwaj et al., 2018; More than 1 author
2.use the latest reference, please

Patel (2012) observed that in a wheat field experiment under three nitrogen levels (0, 60, and 120 kg N/ha) the higher values for growth attributes such as plant height and number of leaves/plants; yield attributes such as number of grains/spikes, spike length grain and straw yield were reported in the highest nitrogen fertilized plot (120 kg N/ha). Among the four wheat genotypes evaluated, two genotypes nitrogen consumption efficiency (agronomic efficiency) at 60 kg nitrogen application was significantly higher than the other genotypes at 120 kg N/ha. In a study, the effects of three N, P, and K levels (35-25-25, 70-50-50, and 105-75-75 kg ha⁻¹) on the growth, yield, and quality of three wheat types were investigated by Hussain *et al.* (2020). Plant height, number of viable tillers, 1000-grain weight, grain yield, and grain protein content of wheat were all altered by different NPK levels. The application of 105-75-75 kg NPK ha⁻¹ resulted in the maximum grain production (4.99 t ha⁻¹).

Comment [W17]: Patel et al., 2012; more than 1 author

In an experiment, the most important durum wheat varieties were examined with four N fertilization levels (0, 60, 120, and 180 kg N ha⁻¹). The results showed that at 120 kg N ha⁻¹ of fertilizer ensured a fair balance of yield and nitrogen use efficiency. In fact, the yield (3.01 and 3.07 t ha⁻¹, respectively) and protein content (13.7 and 13.5 percent) at 120 and 180 kg N ha⁻¹ level were identical. Pre-anthesis N uptake accounted for roughly 67.5 percent of total N uptake and it was strongly and positively linked with wheat production. The importance of late N supply in grain quality was confirmed by a positive connection between post-anthesis N uptake and grain protein concentration (Montemurro, 2007).

Comment [W18]: in reference to 2002 not 2022

The effects of various nitrogen fertilization levels (0, 60, 120, and 180 kg/ha N) on the agronomic performance of six wheat (*Triticum aestivum* L.) cultivars were investigated by Benin (2012). The result indicated that in the cultivars tested, there was genetic diversity in response to nitrogen fertilization. Higher nitrogen fertilizer levels were linked to better yield component performance and a more suited water regime resulted with the greatest yield gains.

Comment [W19]: 1. Montemurro et al., 2007; More than 1 author
2. use the latest reference, please

Tahir (2020) studied 12 bread wheat cultivars released for a heat-stressed environment on yield performance for two seasons at four nitrogen levels (0 (N₀), 43 (N₄₃), 86 (N₈₆), and 129 (N₁₂₉) kg/ha). Increasing N levels from N₀ to N₄₃, N₈₆, and N₁₂₉ resulted in yield increases of 4-45 percent, 13-69 percent, and 34-87 percent at N₄₃, N₈₆, and N₁₂₉, respectively, when averaged across the seasons. These increases in crop yield were linked to increases in biomass growth (r = 0.86).

Comment [W110]: Benin et al., 2012; More than 1 author

The efficiency of wheat crops that utilizes nitrogen effectively is determined by genetic and environmental factors. Kraisig *et al.* (2021) evaluated the efficiency of nitrogen consumption by wheat genotypes as a measure of biomass, productivity, and grain quality parameters. The result indicated that there were reported genetic differences in nitrogen usage efficiency, productivity (yield and biomass), and wheat quality. With an increase in N-fertilizer, wheat crop production and biomass daily rate⁻¹ are affected significantly (Brezolin, 2016).

Comment [W111]: not listed in the reference

Patel (2012) observed that in a wheat field experiment under three nitrogen levels (0, 60, and 120 kg N/ha) the higher values for growth attributes such as plant height and the number of leaves/plants; yield attributes such as the number of grains/spikes, spike length grain and straw yield were reported in the highest nitrogen fertilized plot (120 kg N/ha). Among the four wheat genotypes

Comment [W112]: Brezolin et al., 2016; More than 1 author

Comment [W113]: Patel et al., 2012; more than 1 author

evaluated, two genotypes' nitrogen consumption efficiency (agronomic efficiency) at 60 kg nitrogen application was significantly higher than the other genotypes at 120 kg N/ha. In a study, the effects of three N, P, and K levels (35-25-25, 70-50-50, and 105-75-75 kg ha⁻¹) on the growth, yield, and quality of three wheat types were investigated by Hussain *et al.* (2020) Plant height, the number of viable tillers, 1000-grain weight, grain yield, and grain protein content of wheat were all altered by different NPK levels. The application of 105-75-75 kg NPK ha⁻¹ resulted in maximum grain production (4.99 t ha⁻¹).

The effects of various nitrogen fertilization levels (0, 60, 120, and 180 kg/ ha N) on the agronomic performance of six wheat (*Triticum aestivum* L.) cultivars were investigated by Benin (2012) the result indicated that in the cultivars tested, there was genetic diversity in response to nitrogen fertilization. Higher nitrogen fertilizer levels were linked to better yield component performance and a more suited water regime resulted in the greatest yield gains.

Comment [W114]: Benin et al., 2012; More than 1 author

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The field experiment was conducted during the *rabi* season of 2020-21 at the research farm, ICAR-IISS, Bhopal and Madhya Pradesh. It comes under a semi-arid and sub-tropical zone and is characterized by hot summer and cold winter. Bhopal comes in Vindhyan Plateau Agro-climatic Zone. Mean annual precipitation is about 1100 mm, most of which is received during the monsoon period of July to September. The average maximum temperature during summer is 35-40°C, while the average minimum temperature during winter is 2-9°C.

Table 1 Weather data

	November	December	January	February	March	April	May
Temperature °C	22.4 °C	18.8 °C	20.0°C	21.1 °C	26 °C	26 °C	31.3 °C
Precipitation / Rainfall (mm)	303.8	254	203.2	304.8	228.6	228.6	101.6
Humidity (%)	0.47	0.5	0.5	0.42	0.29	0.29	0.21
Rainy days (d)	1	1	1	1	1	1	1
Avg. Sun hours (hours)	9.7	9.4	9.3	10	10.8	10.8	11.4

The wheat types were manually planted in lined furrows at a rate of 100 kg/ha on November 25, 2020, about 3 cm deep, with a row to row distance of 22.5 cm and a plant to plant distance of 5 cm

Nine varieties of wheat (*T. aestivum* and *T. durum*) crop were selected and grown as test crops in the current investigation adopting a split-plot design replicated thrice with nutrient dose as the main plot and varieties as subplot treatment there are 36 plots in a block (9 variety x 4 fertilizer N and P treatments). The area of each plot is 2m x 2m.

With three replications, the experiment was set up in a split-plot design. There were 36 treatments in total, including three nitrogen levels and three P levels with nine genotypes, in each replication. In each replication, all of the treatments were split separately.

In all treatments except control, full-recommended doses of P and K fertilizers @ 60 kg P₂O₅ & 40 kg K₂O per acre were treated as a basal in SSP and MOP, respectively. N, P, and K were not included in the control plots (T₁). Normal dose treatment (T₂) received the authorized amount of N (120 kg ha⁻¹) and P (60 kg/ha) while T₃ received half of the prescribed quantity of N (60 kg/ha) and a full dose of P (60 kg/ha). The T₄ treatment received the full dose of N and half dose of P fertilizers. 50 percent of the applied N is provided as basal, while the remaining N was top-dressed in equal splits at 25 and 45 DAS, respectively. The wheat crops were cultivated following standard agronomic practices and five irrigations (5 cm) and one manual weeding at 30 DAS was done.

The leaf area of plant samples was measured with the LICOR Leaf area meter (model 1100) and expressed in cm². Leaf area was assessed by choosing three plants at random in each treatment plot at 50, 65 and 85 DAS.

Plant height was taken at the physiological maturity stage of the crop in each replication comprising 5 plants and it was expressed in cm.

Plant samples were collected from the treated plots and control plots. The whole plant was divided into above-ground biomass of the samples were dried at 65°C in an oven for 72 hrs or till constant weight. The dry weight of the plant sample was expressed in grams.

Plant samples were collected from the treated plots and control plots. The whole plant was divided into above ground biomass of the samples were dried at 65°C in oven for 72 hrs or till constant weight. The dry weight of plant sample was expressed in gram.

The crop was harvested after it reached maturity, when the foliage had faded and the ears had assumed a yellowish hue. To eliminate the border impact, one border row from both sides and 30 cm from either end of each plot were collected. The remaining experimental plots were harvested with sickles, with ears detached and plant biomass wrapped into bundles. Each treatment's plant biomass (kg/ha), grain yield (kg/ha), and straw yield (kg/ha) were measured individually and weighed.

3. RESULTS AND DISCUSSION

The crop was harvested after it reached the maturity when the foliage had faded and the ears had assumed a yellowish hue. To eliminate the border impact, one border row from both sides and 30 cm from either end of each plot were collected. The remaining experimental plots were harvested with sickles, with ears detached and plant biomass wrapped into bundles. Each treatment's plant biomass (kg/ha), grain yield (kg/ha), and straw yield (kg/ha) were measured individually and weighed.

Comment [W115]: It is suggested that the discussion about the tables and figures should appear in the text **before the appearance of the respective tables and figures**

Table 2 Results of different wheat varieties

wheat varieties	Dry matter			Days to 50% flowering	Days to maturity	Leaf area (cm ²)			Plant Biomass kg/ha.	Grain yield kg/ha.
	50 DAS	65 DAS	85 DAS			50 DAS	65 DAS	85 DAS		
	Mean A	Mean A	Mean A	Mean A	Mean A	Mean A	Mean A	Mean A	Mean A	Mean A
HI8663	71.38	1.55	1.51	1.46	66.25	105.00	163.20	149.40	154.10	6345.81
HI8737	71.75	1.52	1.58	1.52	67.88	107.75	160.40	152.50	157.40	7754.31
HI8713	83.00	1.98	1.76	1.65	68.88	119.38	166.40	154.80	152.00	8409.38
HI1563,	78.88	1.48	1.33	1.23	70.75	116.63	168.40	157.30	148.60	8281.22
HI1544	87.63	1.67	1.64	1.50	67.13	116.75	153.80	165.10	144.50	8151.13
HI1531	79.63	1.48	1.49	1.33	71.75	125.25	146.50	171.10	136.80	7962.24
GW366	87.13	1.52	1.48	1.33	76.38	118.13	146.40	161.10	139.40	8605.75
LOK1	84.63	2.22	2.08	2.04	74.75	124.50	172.00	164.80	158.30	8738.28
NARMADA14	83.75	2.85	2.54	2.35	67.38	128.13	182.20	160.50	169.40	9109.13
Mean B	80.86	1.81	1.71	1.60	70.13	117.95	162.14	159.62	151.17	8150.81
Factors	C.D.									
Factor(A)	0.37	0.19	0.13	0.06	1.85	1.51	4.11	11.19	5.72	182.07
Factor(B)	3.05	0.10	0.16	0.06	0.99	1.79	4.45	9.34	7.01	163.38
Factor(B) at same level of A	NS	0.24	0.32	0.13	2.31	3.75	9.36	20.22	14.60	350.44
Factor(A) at same level of B	NS	0.26	0.31	0.13	2.56	3.66	9.23	20.49	14.25	351.94

Where Factor A= Treatment, Factor B = Varieties




Fig. 1. Effect of N & P on biomass yield of wheat genotypes




Fig. 2. Effect of N & P on grain yields of different wheat genotypes

There was a strong influence of the three nitrogen and three phosphorus dosages. Plant height grew as the nitrogen level increased from zero to 120 kg/ha. When the nitrogen dose was 120 kg/ha, the maximum plant height (91 and 90 cm) was recorded, whereas the minimum plant height (65 cm) was reported in the control. The average plant height for the nine cultivars over nitrogen dosages revealed that HI1544 had the highest plant height (87.63 cm) and HI8663 had the lowest plant height (71.38 cm). It was discovered that there was a considerable relationship between nitrogen, phosphorus, and cultivars. However, when the nitrogen dose was 120 kg/ha and the P dose was 30 kg, GW366 had the highest plant height (91 cm), but HI8737 had a minimum plant height (65cm) with 0 kg/ha N and P fertilizers. The findings of Saleem (1987) and Khan *et al.* (2000), who found that increasing nitrogen levels increased plant height, corroborate these findings. Plant height was shown to be higher with increased nitrogen and phosphorus administration in the current study. Ullah *et al.* (2018) found a

comparable increase in plant height with increased fertilizer dose, with higher plant height being recorded with higher Nitrogen @203 kg/ha dosages. (Gwal *et al.*, 1999, Jan *et al.*, 2002, Liaqat *et al.*, 2003, Rakaby *et al.*, 2016) found a similar rise in plant height with additional nitrogen fertilizer. Because nitrogen is the most important nutrient for plant growth and development, proper nitrogen dosage is critical for maintaining plant growth processes that lead to increased plant height. According to Lenka and Behera, (1967) increased N levels significantly increased plant height (1967).

There were significant differences in the number of days attaining 50% flowering among varieties of wheat and fertilizer treatments. The number of days required to get 50% flowering was the lowest in HI8663 (62.5 days) and the highest in GW366 (80 days) (Table 2). Among the treatments, days to 50% flowering was the higher in control plots followed by, reduced phosphatic fertilizer dose treatment, reduced nitrogen dose fertilizer treatment and the lowest in 100% NP dose treatment. In this study, early flowering was observed with nitrogen fertilization, while it was delayed by a few days in the control conditions. The control conditions in the current study might have induced nitrogen deficit, causing a delay in leaf development and leading to a delay in flowering by a few days. Several studies have shown that N deficiency/starvation causes delayed flowering in Arabidopsis, Kalanchoe, and Lemna (Castro Marn *et al.*, 2011; Dickens, 1988; Tanaka, 1986; Yuan *et al.*, 2016). Lin and Tsay (2017) proposed that Arabidopsis has a Nitrogen regulated U-shaped flowering response. To see if this reaction can be extended to other plants and crops, a wide range of N concentrations must be explored systematically with a variety of species.

There were significant variations in days to maturity observed in terms of varieties of wheat and fertilizer treatments. Selected varieties matured between 100.5 (HI8663) to 135 days (Narmada14) among the nutrient treatments (Table 3). Varieties took a longer time to mature in control plots followed by, reduced nitrogen dose fertilizer treatment, reduced phosphatic fertilizer dose treatment and normal NP treatment. In a full dose of N & P treatment, the highest number of days taken for maturity was found in Narmada14 (125 days) and the lowest number of maturity days was found in HI8663 (100.5 days).

Wheat varieties and NP treatments had a significant effect on the leaf area of wheat at 50 DAS (Table 4). Leaf area was found between 122.18 (GW366) to 191.67 cm²/per plant (NARMADA14) among all the treatments (Table 4). The mean leaf area of the wheat plant was higher in 100% NP treatment (T₂) than in reduced phosphatic fertilizer dose treatment (T₄) and reduced nitrogen dose fertilizer treatment (T₃), and the lowest was in control plots. In T₂, the highest Leaf area was recorded in Narmada14 (191.67 cm²/per plant) and the lowest Leaf area was found in HI1531 (162.53 cm²/per plant). The highest leaf area at every stage of sampling was obtained in T₂ (100%NPK) and the lowest was in T₁ (control). Not much difference was observed between 50 DAS and 65 DAS, whereas it was lowest in 85 DAS. Amongst the varieties, the leaf area of Narmada 14 was the highest at 50 DAS and 85 DAS, whereas HI1531 recorded the highest leaf area at 65 DAS (Table 4). The lowest leaf area was recorded in GW 366 at 50 DAS; however, it was HI8663 and HI 1531 at 65 and 85 DAS, respectively. Different nitrogen and phosphorus application doses resulted in significant differences in the leaf area (Table 3). At all growth stages of measurements, the normal

dose treatment produced the highest leaf area in NARMADA 14(191.67 cm²) and Lok1 (182.97 cm²) while the treatment got no nitrogen and phosphorus (T₁) produced the lowest leaf area (GW 366 and HI1531). The present results conformed with the findings of Bali *et al.* (1991) wherein a maximum leaf area with increased N fertilization was reported. In the presence of increased N, the rise in leaf area is generated by an increase in the number of tillers and the size of subsequent leaves. Nitrogen enrichment has been shown in numerous studies to enhance leaf area, flag leaf area, and plant development (Khurshheed & Mohammed,2015).

Comment [W116]: 3.not listed in the reference
4.use the latest reference

Comment [W117]: Not listed in the reference

The effect of different nutrient treatments on total biomass yield and grain yield of nine wheat varieties are given in Table 2. There were significant differences in plant biomass observed between varieties of wheat and fertilizer treatments (Table 5). Among all the treatments the range of Plant biomass was found in between 5101.50 kg/ha (HI8663) to 10377.25 kg/ha (NARMADA14). The mean Plant biomass was higher in Normal dose treatment followed by, reduced phosphatic fertilizer dose treatment, reduced nitrogen dose fertilizer treatment and lower in control plots. Among the varieties grown in a full dose of N & P treatment, the highest Plant biomass was found in NARMADA14 (10377.25 kg/ha) and the lowest Plant biomass was found in HI8737 (8050.25 kg/ha). Among the varieties grown in a half dose of N fertilizer treatment the highest Plant biomass was observed in LOK1 (8781.25 kg/ha) followed by GW366 (8751.25 kg/ha) and among the varieties grown in a half dose of P, the highest plant biomass was found in NARMADA14 (9670.25kg/ha) followed by HI1563 (9657.75kg/ha). Total biomass yield is the total dry matter generated by a plant as a result of photosynthesis and nutrient uptake after accounting for the losses during respiration (Shah, 1994). The selected nine wheat cultivars have shown considerable variations in biological yield (t/ha). There was a lot of interaction between the nine cultivars and the four levels of nutrients (nitrogen and phosphorus treatments). As the amount of nitrogen and phosphorus application was increased from the control level to 60 kg/ha and 120 kg/ha, the biological yield increased. Normal dose treatment yielded the highest biological output (10.37 t/ha).

Summery and Conclusion

Comment [W118]: a. summery???
b. 4.Conclusion
c. In conclusion: briefly state the major findings of the study

Normal RDF of nitrogen and phosphorus increased the plant height of wheat, but low N conditions resulted in the lowest plant height. The wheat genotype HI 1544(87.63 cm) had a greater mean plant height whereas HI8663 (71.38cm) had a lower mean plant height. However, varieties GW366(91cm) grown under optimum N with sub-optimum P dose also exhibited higher plant height similar to an optimum dose of both N & P. Normal RDF of nitrogen and phosphorus increased the dry matter weight of the plant at all three stages of growth, 50, 65, and 85 DAS. At 85 DAS, NARMADA14 (3.06 g/plant) had the highest dry matter weight, followed by lok1 (2.83 g/plant).

Early flowering was observed with nitrogen and phosphorus fertilization, but it was delayed by a few days in control plots. In the current study, the control condition may have resulted in nitrogen deficiency, which may have caused flowering to be delayed by a few days. Varieties grown under normal RDF (T₂) had a higher leaf area. HI 1531(171.13 cm²) at 65 DAS and NARMADA14 (169.38 cm²) at 85 DAS exhibited the highest mean leaf area among all the varieties.

Under normal, reduced N dose reduced P dose condition and control fertilizer doses, the variety HI8713 produced considerably larger plant biomass of 8383.00 kg/ha, 9,393.00 kg/ha, 7,913.25 kg/ha, 7,948.25 kg/ha. The same variety produced considerably higher grain yields of 3,035.13 kg/ha, 4,580.88 kg/ha, 3,996.63 kg/ha, 4190.88 kg/ha respectively under similar fertilizer doses.

Of the four nutrient treatments, the recommended dose of N, P & K T₂ performed the best in terms of yield attributes, grain yield and different use efficiency terms in all the nine selected varieties of wheat, while no nutrient treatment T₁ was the least. Among all the wheat varieties tested the variety HI 8713 had considerably higher dry weight, leaf area, biomass, grain yield, lower days to 50% flowering, between sub-optimal doses of N and P (T₃ and T₄) T₃ produced higher biomass yield than T₄ averaging all the selected nine varieties of wheat, whereas T₄ had higher grain yield than T₃. In terms of mean grain yield cutting across all the treatments, the selected nine varieties followed the trend

HI8713>HI1531>NARMADA 14>GW366>HI1563>HI8737>HI8663>HI1544>LOK1

References

- Belete, F., Dechassa, N., Molla, A., & Tana, T. (2018). Effect of nitrogen fertilizer rates on grain yield and nitrogen uptake and use efficiency of bread wheat (*Triticum aestivum* L.) varieties on the Vertisols of central highlands of Ethiopia. *Agriculture & Food Security*, **7**(1), 1-12.
- Benin, G., Bornhofen, E., Beche, E., Pagliosa, E. S., Silva, C. L. D. and Pinnow, C. (2012). Agronomic performance of wheat cultivars in response to nitrogen fertilization levels. *Acta Scientiarum. Agronomy* **34**(3), 275-283.
- Bhardwaj, V., Yadav, V. and Chauhan, B. S. (2010). Effect of nitrogen application timings and varieties on growth and yield of wheat grown on raised beds. *Archives of Agronomy and Soil Science* **56**(2), 211-222.
- Brezolin, A. P., da Silva, J. E. A. G., Frantz, F. C. R., Binelo, M. O., Valdiero, A. C., Zimmer, C. M., & Mazurkiewicz, G. (2016). The efficiency of wheat yields by nitrogen dose and fractionation. *African Journal of Agricultural Research*, **11**(36), 3440-3449.
- Dhull, S., Goyal, S., Kapoor, K., & Mundra, M. (2004). Microbial biomass carbon and microbial activities of soils receiving chemical fertilizers and organic amendments. *Archives of Agronomy and Soil Science*, **50**(6), 641-647.
- Hussain, I., Khan, M. A. and Khan, E. A. (2006). Bread wheat varieties as influenced by different nitrogen levels. *Journal of Zhejiang University Science B* **7**(1), 70-78.
- Hussain, M. I., Shah, S. H., Hussain, S. A. J. J. A. D. and Iqbal, K. H. A. L. I. D. (2002). Growth, yield and quality response of three wheat (*Triticum aestivum* L.) varieties to different levels of N, P and K. *International Journal of Agriculture and Biology* **4**(3), 362-364.

Comment [W119]: 1. Use reference style for author guideline
2. Use latest reference [not more than 10 years]

Irfan, M., Abbas, M., Shah, J. A., & Memon, M. Y. (2018). Internal and external phosphorus requirements for optimum grain yield are associated with P-utilization efficiency of wheat cultivars. *Communications in Soil Science and Plant Analysis*, **49**(22), 2843-2853.

Kaur, G., Asthir, B. and Bains, N. S. (2015). Nitrogen levels effect on wheat nitrogen use efficiency and yield under field conditions. *African Journal of Agricultural Research* **10**(23), 2372-2377.

Montemurro, F., Convertini, G., & Ferri, D. (2007). Nitrogen application in winter wheat grown in Mediterranean conditions: effects on nitrogen uptake, utilization efficiency, and soil nitrogen deficit. *Journal of plant nutrition*, **30**(10), 1681-1703.

Nasri, R., Kashani, A., Barary, M., Paknejad, F. and Vazan, S. (2014). Nitrogen uptake and utilization efficiency and the productivity of wheat in double cropping system under different rates of nitrogen. *International Journal of Agriculture and Biosciences* **4**(4), 184-193.

Patel, C. B., Singh, R. S., Yadav, M. K., Singh, S. K., Singh, M. K., Singh, K. K., & Mall, R. K. (2012). Response of different wheat (*Triticum aestivum* L. emend Fiori&Paol.) genotypes to various nitrogen levels under late sown conditions of Eastern Uttar Pradesh. *Ecology* **30**(3C), 1192-1196.

Patel, C. B., Singh, R. S., Yadav, M. K., Singh, S. K., Singh, M. K., Singh, K. K., & Mall, R. K. (2012). Response of different wheat (*Triticum aestivum* L. emend Fiori&Paol.) genotypes to various nitrogen levels under late sown conditions of Eastern Uttar Pradesh. *Ecology*, **30**(3C), 1192-1196.

Comment [W120]: Doble

Schuur, E. A. (2011). Nitrogen from the deep. *Nature* **477**(7362), 39-40.

Tahir, I. S. A., Elbashier, E. M. E., Ibrahim, M. A. S., Saad, A. S. I., & Abdalla, O. S. (2020). Genetic gain in wheat grain yield and nitrogen use efficiency at different nitrogen levels in an irrigated hot environment. *International Journal of Agronomy*, **2020**.

Comment [W121]: Vol Number???