

## Effect of plant spacing and manganese on Yield, Economics and Quality of sesame (*Sesamum Indicum L.*)

### Abstract

A field experiment was conducted during *Zaid* season (2022) at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P.), India, to study the influence of spacing and manganese on yield and economics and quality of sesame. The treatments consisted of 3 levels of spacing (30×10cm, 40×10cm, 50×10cm) and 3 levels of manganese (1.5, 3.0 and 4.5 kg/ha). The experiment was laid out in Randomized Block Design with 10 treatments and replicated thrice. The results recorded significantly higher in yield and yield attributes viz. number of capsules/plant (52.47), number of seeds per capsules (74.50), test weight (3.37 g), seed yield (1.19 t/ha), stover yield (2.67 t/ha) and oil content (48.10%). Maximum Gross returns (113050.00 (Indian Rupee) INR/ha), Net returns (76944.15 (Indian Rupee) INR/ha) and Benefit cost ratio (2.13) was found in treatment (T<sub>9</sub>) with the application of 50×10cm + manganese sulfate (MnSO<sub>4</sub>) 4.5 Kg/ha.

**Keywords:** - Spacing, Yield, Manganese, Zaid, Economics.

### 1. INTRODUCTION

Sesame (*Sesamum indicum L.*) is an important oil seed crop which plays a crucial role in Indian agricultural economy both in area as well as production. It is a drought resistant crop and it can be easily survive or grown under rain-fed conditions. Sesame is mostly grown in summer season and also in Kharif season. Sesame is the oldest known oil seed crop and it is popular worldwide. It is commonly known as Till, Gingelly, Simsim. Sesame has the highest protein contents and oil content. (Raja *et al.*, 2007). Oil content of sesame varies from 46-52%. Sesame seeds are rich in protein, energy, dietary fiber, carbohydrate and also contain riboflavin, thiamine. Proper spacing supplies sufficient light, efficient absorption of nutrients and water from soil and at optimum spacing avoids intra space competition. Spacing has

effect on yield attributes viz., number of capsules/ plant, number of seeds/ capsules (2) (Subrahmaniyan *et al* .,) Optimum plant spacing starts the sesame plant to grow properly both its aerial and underground parts through utilizing radiant energy more to encourage crop production. (3) (Shinde *et al* .,)spacing will bring a positive effect on yield.It is most important component in farming system in intensive method.Micro nutrients bring positive effect on yield parameters. Manganese in plants is a constituent and activator of enzymes involved in protein synthesis,lipid metabolism and photosynthesis. Manganese deficiency mostly leads to  $\text{NO}_3^-$  - N in plant tissues. The signs of manganese deficiency in plants include decrease in number of lower per plant, infertility of pollen. Keeping these points in view an experiment was conducted to achieve yield with profitable investment by treatment combinations of different spacing and also manganese levels.

## 2. MATERIALS AND METHODS:

A field experiment was carried out during *Zaid* season of 2022. The experiment was conducted in Randomized Block Design and it consists of ten treatment combinations with three replications and was laid out variously with different treatments assigned randomly in each replication. The soil in experimental field was sandy loam texture, having alkaline reaction (pH 7.1) with very low organic carbon (0.28%), available higher n (225 Kg/ha), p (19.50 kg/ha) and higher level of potassium (92.00 kg/ha). Treatment combinations were T<sub>1</sub> - 30×10 cm + Manganese sulfate (MnSO<sub>4</sub>)1.5 kg/ha, T<sub>2</sub> - 40×10cm + MnSO<sub>4</sub> 1.5 kg/ha, T<sub>3</sub> - 50×10cm + MnSO<sub>4</sub> 1.5 kg/ha, T<sub>4</sub> - 30×10cm + MnSO<sub>4</sub> 3.0 kg/ha, T<sub>5</sub> - 40×10 cm+ MnSO<sub>4</sub> 3.0kg/ha, T<sub>6</sub> - 50×10 cm+ MnSO<sub>4</sub> 3.0kg /ha, T<sub>7</sub> - 30×10 cm + MnSO<sub>4</sub> 4.5 kg/ha, T<sub>8</sub> - 40×10cm+ MnSO<sub>4</sub> 4.5 kg/ha, T<sub>9</sub> - 50× 10cm + MnSO<sub>4</sub> 4.5 Kg/ha, T<sub>10</sub> - Control ( RDF) . The observations were recorded on yield and yield parameters i.e. Number of capsules/plant, number of seeds/ capsule, test weight, grain yield and stover yield, oil content, gross returns, net returns, Benefit- Cost Ratio (BCR)

## 3. RESULTS AND DISCUSSION

### A. Yield:

Treatment with the application of 50×10 cm + (Manganese sulfate) MnSO<sub>4</sub> 4.5kg/ha was reported maximum number of capsules/ plant (52.47) which was significantly superior over all from other and treatment with application of 50×10cm +MnSO<sub>4</sub> 3.0 Kg/ha (50.53) which was

statistically at par with the treatment with the application of 50×10cm+ MnSO<sub>4</sub> 4.5 Kg/ha. Treatment with application of 50×10cm+ MnSO<sub>4</sub> 4.5 Kg/ha was recorded maximum number of seeds per capsules (74.50) which was significantly superior over all other and treatment with application of 30×10 cm + MnSO<sub>4</sub> 4.5 Kg/ha (73.07) and 50×10cm + MnSO<sub>4</sub> 1.5 Kg/ha (72.67) which was statistically at par to the treatment with application of 50×10cm + MnSO<sub>4</sub> 4.5 Kg/ha. Treatment with application of 50×10cm+ MnSO<sub>4</sub> 4.5 Kg/ha was recorded maximum test weight (3.37g) which was significantly superior over all other and treatment with application of 50×10cm + MnSO<sub>4</sub> 3.0 Kg/ha (3.20) which was statistically at par to the treatment with application of 50×10cm + MnSO<sub>4</sub> 4.5 Kg/ha. Treatment with application of 50×10cm + MnSO<sub>4</sub> 4.5 Kg/ha was recorded maximum seed yield (1.19 t/ha) which was significantly high over all other and treatment with application of 40×10cm +MnSO<sub>4</sub> 4.5 Kg/ha (1.16) which was statistically at par with the treatment with application of 50×10 cm + MnSO<sub>4</sub> 4.5 Kg/ha. Treatment with application of 50×10cm +MnSO<sub>4</sub> 4.5 Kg/ha was recorded maximum stover yield (2.67t/ha) which was significantly more from all other and treatment with application of 50×10cm + MnSO<sub>4</sub> 3.0 Kg/ha (2.66 t/ha) which was statistically at par to treatment application of 50×10cm +MnSO<sub>4</sub> 4.5 Kg/ha. Treatment with application of 50×10 cm + MnSO<sub>4</sub> 4.5 Kg/ha was observed maximum harvest index (30.82 %) and minimum with application of 40×10 cm + MnSO<sub>4</sub> 4.5 kg/ha (30.36%). (Elayaraja *et al* 2019).

### **B. Oil Content :-**

With treatment application of 50×10cm + MnSO<sub>4</sub> 4.5kg/ha found more oil content (48.10%) and higher compared to all other treatments. With treatment of 50×10 cm +MnSo<sub>4</sub>3.0kg/ha observed (47.54%) and it was statistically at par with 50 ×10cm + MnSO<sub>4</sub> 4.5kg/ha.

### **C. Economics:**

Economic viability and efficiency of crop cultivation are mostly outcome of crop production with less production cost. Higher cost of cultivation ( 36,105.85)was observed in (T<sub>9</sub>) 50× 10cm + MnSO<sub>4</sub> 4.5kg/ha. Maximum Gross returns ( 113050.00₹/ha), net returns ( 76944.15₹/ha ), Benefit -cost ratio (BCR) ( 2.13) was reported in (T<sub>9</sub>) 50× 10 cm + MnSO<sub>4</sub> 4.5kg/ha.

## **4. CONCLUSION**

Based on conclusions of one season experimentation it can be concluded that application of 50×10cm+MnSO<sub>4</sub> 4.5 kg/ha was found more productive (1.19 t/ha) and it can be recommended to farmers after further trails.

### **FUTURE SCOPE**

As there was less research happened in this field, further research should be done to obtain better results and help farmers obtaining better yield. Since findings are based on one season further trails are needed to confirm the results of this experiment.

### **ACKNOWLEDGEMENT**

The authors are thankful to Dr. Rajesh Singh Associate Professor Department of Agronomy , SHUATS, Prayagraj,U.P. India. for providing us necessary facilities to undertake the studies.

**References.**

1. Bhaduria , .S., Nidhi nagar and Sohan , Krishan mudgal (2012). Effect of micro nutrient cations on yield, quality and their uptake by mustard in alluvial soil. *Annals of plant and soil research* . **14**(2) : 130-132.
2. Elayaraja, D., Sathiyamurthi S. and Kamalakannam P. (2019). Study on influence of organics and micro nutrients for increasing sesame (*Sesamum indicum L.*) production and sustainable soil fertility in coastal and soils. *Journal of Pharmacognosy and Phytochemistry* **2**: 483-487.
3. Jan, A., Ali, S., Adail, M. and khan, A. (2014) Growth and yield components of sesame (*Sesamum indicum L.*) as influenced by phosphorous levels under different row spacing. *Journal of Environment and Earth Science*. **4** (22), 2224-3216.
4. Nadeem, A., Kashani, S., Ahmed, N., Mahmooda, B., Saeed, Z., Fateh, M. and Ahmed, S. (2015). Growth and yield of sesame (*Sesamum indicum L.*) under the influence of planting geometry and irrigation regimes. *American Journal of Plant Science*, **6**: 980- 986.
5. Richa ,S.A, Rajshree , P.and K., N,Namdeo.(2019). Effect of plant spacing on growth, yield and quality of mustard ( *Brassica juncea L.*) genotypes. *Annals of plant and soil research* (2019) **21**(2): 172-176.
6. Saini, J.S., Tripathi, H.P., Devivedi, R.S. and Randhauwa, M.S. (1975). Effect of micronutrients on the yield and quality of groundnut. *Pumjab journal research*. **12**(3): 224-227.
7. Shah S.N., Patel K.H., Patel P.A. (2020) Effect of Spacing and Topping on yield of summer sesame (*Sesamum indicum L.*). *International journal of current Microbiology and Applied sciences*. **9** (5) 2312-2319.
8. Shinde, S.D.; Raskar, B.S. and Tamboli, B.D. (2011). Effect of spacing and sulphur levels on productivity of sesame (*Sesamum indicum L.*) under summer condition. *Journal of Maharashtra Agricultural University*, **36**(1):28 – 31.

9. Sivagamy, k. and Rammohan, J. (2013). Effect of sowing data and crop spacing on growth, yield attributes and quality of sesame. *IOSR journal of Agriculture and Veterinary Science* **5**(2): 38-40.
10. Subramaniyan K, Arulmozhi N, Kalaiselven P. Influence of plant density and NPK levels on the growth and yield of sesame ( *Sesamum indicum* L.) genotype. *Griculture science digest* (2001); **21** (3): 208-209.
11. Vahid , G., Ghalavand, A., Ali Soroosh, Z., Alireza, P.(2010). Effectt of iron , Zinc, Manganese on quality and quantity of Soya bean seed.*Journal of phytology*. **2** (11): 73.
12. Yadav, R. A.; Mishra, A. and Singh, D. (2007). Response of sesame cultivars under various plant spacing and seed rates. *Plant Archives*, **7**(1): 287 – 288.
13. Yilmaz, A., Kiz,H.E., Torun,B., Gulkin,I.,Karaniks, Bagci,A.and Cakmak,I. ( 1997). Effect of different zinc application methods on grain yield and zinc concentration in wheat cultivars grown on Zinc deficient calcareous soils. *Journal of plant nutirtion*. **20**: 461-471.

Table.1 Effect of plant spacing and manganese on yield and yield attributes of sesame.

Treatments	No. of capsules per plant	No. of seeds per capsules	Test weight (g)	Seed yield (t/ha)	Stover yield (t/ha)	Harvest index (%)
30×10 cm + MnSO <sub>4</sub> 1.5 kg/ha	47.03	68.87	2.10	1.00	2.54	29.37
40 ×10cm+ MnSO <sub>4</sub> 1.5 kg/ha	45.17	69.10	2.63	1.04	2.57	28.80
50×10cm+ MnSO <sub>4</sub> 1.5 kg/ha	46.77	72.67	3.07	1.07	2.59	29.23
30×10cm + MnSO <sub>4</sub> 3.0 kg/ha	46.47	68.17	2.40	1.10	2.60	29.72
40 ×10cm + MnSO <sub>4</sub> 3.0 kg/ha	45.27	72.60	2.77	1.11	2.62	29.75
50×10 cm + MnSO <sub>4</sub> 3.0 kg /ha	50.53	74.43	3.20	1.17	2.64	30.70
30×10 cm + MnSO <sub>4</sub> 4. 5 kg/ha	47.60	73.07	2.43	1.13	2.61	30.21
40 ×10cm + MnSO <sub>4</sub> 4.5 kg/ha	48.40	72.07	2.90	1.16	2.66	30.36
50×10 cm + MnSO <sub>4</sub> 4.5 Kg/ha	52.47	74.50	3.37	1.19	2.67	30.82
Control ( RDF)	46.97	65.03	2.07	0.90	2.50	26.47
SEm (±)	1.02	1.75	0.09	0.04	0.01	0.02
CD (5%)	3.05	5.22	0.28	0.14	0.03	0.06

Table.2 Effect of plant spacing and manganese on economic indicators

S.no	Treatments	Total cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C ratio
1.	30×10 cm + MnSO <sub>4</sub> 1.5 kg/ha	35,930.55	94683.33	58752.78	1.64
2.	40 ×10cm+ MnSO <sub>4</sub> 1.5 kg/ha	35,830.55	98800.00	62969.45	1.76
3.	50×10cm+ MnSO <sub>4</sub> 1.5 kg/ha	35,630.55	101650.00	66019.45	1.85
4.	30×10cm + MnSO <sub>4</sub> 3.0 kg/ha	36,116.85	104183.33	68066.48	1.88
5.	40 ×10cm + MnSO <sub>4</sub> 3.0 kg/ha	36,016.85	105766.67	69749.82	1.94
6.	50×10 cm + MnSO <sub>4</sub> 3.0 kg /ha	35,916.85	111150.00	75233.15	2.09
7.	30×10 cm + MnSO <sub>4</sub> 4.5 kg/ha	36,305.85	107666.67	71360.82	1.97
8.	40 ×10cm + MnSO <sub>4</sub> 4.5 kg/ha	36,205.85	110200.00	73994.15	2.04
9.	50×10 cm + MnSO <sub>4</sub> 4.5 Kg/ha	36,105.85	113050.00	76944.15	2.13
10.	Control ( RDF)	35,743.35	85183.33	49439.98	1.38

Table .3 Effect of plant spacing and manganese on oil content of sesame

<b>S.no</b>	<b>Treatments</b>	<b>Oil content (%)</b>
<b>1.</b>	<b>30×10 cm + MnSO<sub>4</sub> 1.5 kg/ha</b>	45.20
<b>2.</b>	<b>40 ×10cm+ MnSO<sub>4</sub> 1.5 kg/ha</b>	45.48
<b>3.</b>	<b>50×10cm+ MnSO<sub>4</sub> 1.5 kg/ha</b>	46.02
<b>4.</b>	<b>30×10cm + MnSO<sub>4</sub> 3.0 kg/ha</b>	46.14
<b>5.</b>	<b>40 ×10cm + MnSO<sub>4</sub> 3.0 kg/ha</b>	46.10
<b>6.</b>	<b>50×10 cm + MnSO<sub>4</sub> 3.0 kg /ha</b>	45.70
<b>7.</b>	<b>30×10 cm + MnSO<sub>4</sub> 4. 5 kg/ha</b>	47.54
<b>8.</b>	<b>40 ×10cm + MnSO<sub>4</sub> 4.5 kg/ha</b>	46.45
<b>9.</b>	<b>50 ×10cm + MnSO<sub>4</sub> 4.5kg/ha</b>	47.15
<b>10.</b>	<b>Control ( RDF)</b>	48.10
	SEm (±)	S
	<b>CD (5%)</b>	0.98