

Determinants of uptake of Mass Drug Administration for Schistosomiasis Control in Butiaba Sub-county, Uganda

Abstract.

Introduction: Schistosomiasis is one of the Neglected Tropical Diseases (NTDs) targeted for elimination in Uganda by 2025 through Mass Drug Administration (MDA) using praziquantel. To achieve this, WHO estimates indicate that MDA coverage and uptake of 75% is required. However, coverage remains suboptimal with insufficient knowledge and inadequate drug supply often cited as key reasons. There is need to add to the body of knowledge in various settings to enable more robust mitigation measures. This study aimed to assess the uptake of praziquantel for MDA and associated factors in Butiaba sub-county along the shores of Lake Albert in Uganda.

Methods: A cross-sectional study was conducted in five randomly selected villages within Butiaba sub-county between July and September 2021 using quantitative and qualitative approaches. Semi-structured questionnaires were administered to 450 adults, with additional two Focus Group Discussions and Key Informant interviews held with implementation structures from village to district level.

Results: Self-reported uptake of praziquantel within twelve months of the most recent MDA exercise was 71.56% (95% CI: 67.14 – 75.68). Of all the participants, 5.78% reported having never swallowed praziquantel in their lifetime and 75% (96/128) of participants who didn't swallow praziquantel in the last twelve months reported having at least swallowed the drug in the last ten years. Respondents were less likely to have swallowed praziquantel if they had no knowledge about schistosomiasis signs (AOR=0.18, 95% CI: 0.08–0.39) and more likely if they were between the ages 30–39 years (AOR=2.31, 95% CI: 1.35–3.95) or 40 years and above (AOR= 2.86, 95% CI: 1.45–4.95). Operational challenges such as inadequate supply of praziquantel and financial constraints also influence uptake of praziquantel during MDA in Butiaba sub-county.

Conclusion: Uptake of praziquantel during MDA in Butiaba sub-county was high but still below the WHO target of 75%. People with limited knowledge on schistosomiasis symptoms and those aged 18 – 29 years were less likely to take Praziquantel. Irregular drug supply was also a key challenge.

Recommendation: Rigorous health education and ensuring continuous supply of Praziquantel are key in improving MDA uptake.

Keywords: Schistosomiasis, Mass Drug Administration, Uganda, Sub-Saharan Africa, Uptake, Praziquantel.

1. Introduction

Schistosomiasis, also known as bilharzia, is a neglected tropical parasitic disease transmitted through fresh water snails currently estimated to affect 240 million people worldwide with approximately 800 million people at risk due to poor water, sanitation and hygiene (Organization, 2014). Along with geohelminths, schistosome infections have been estimated to account for 40% of the global tropical disease burden excluding malaria (Olveda et al., 2013). The disease burden is highest in Africa with statistics reporting schistosomiasis as the second most common parasitic disease on the continent putting strain on economies and governments (Adenowo, Oyinloye, Ogunyinka, & Kappo, 2015). Furthermore, the 2022 WHO schistosomiasis fact sheet revealed that at least 90% of those requiring treatment for the disease globally were in Africa (WHO, 2022).

In Uganda, *Schistosoma mansoni* is the most common species causing intestinal schistosomiasis (Emmanuel, Aginya, & Doehring, 2008) which is endemic in 73 districts (Loewenberg, 2014) with a national prevalence of 25.6% and estimates of more than 4 million people infected and 55% of the population at risk (Exum et al., 2019). The disease is particularly of great concern among fishing communities (Parker et al., 2012) such as those along the Lake Albert shores in Butiaba sub-county where the only available data on statistics registered a prevalence as high as 72% in 2004 (Kabatereine et al., 2005).

In a bid to combat this high prevalence, the chief strategy endorsed by the World Health Organization (WHO) is Preventive Chemotherapy (PC) through Mass Drug Administration (periodic large scale population treatment) with Praziquantel as the drug of choice (Organization, 2012) aiming at a target coverage of 75%. As of 2016 however, global coverage rate stood at

14.3% as per the epidemiological report from WHO department of control of neglected diseases and a compliance rate of less than 50% thus an actual level of reach of only about 5% of the intended population(Ross et al., 2017).

In Uganda, the National Bilharzia and worm Control Programme is implemented at the district level. This is done through the District Health Office (DHO) by the District Vector Control Officer (DVCO) and the District Health Educator (DHE)(Government_of_Uganda_Ministry_of_Health, 2010). Praziquantel distributions are done both in schools and in the community. In the communities, trained volunteer drug distributors administer the drug to registered community members at particular sites and door to door visits. Drug distribution is done twice each year in regular intervals to endemic populations because of susceptibility of reinfection.

Despite the use of a variety of avenues to reach out to the endemic communities and the benefits such as reduced schistosomiasis morbidity, MDA uptake is still below target coverage with statistics revealing fluctuating levels of coverage from 64.7% when the national control programme was initially launched in 2003(Narcis B. Kabatereine et al., 2006) to percentages as low as 50% in 2012(Tuhebwe et al., 2015) and 48.8% in 2016(Adriko et al., 2018). Uptake of MDA among adults varies across different communities and is tied to local dynamics as demonstrated by 84.5% coverage registered in Bugiri versus 48.8% coverage in Arua in the 2003/2004 drug distribution(Parker, Allen, & Hastings, 2008a). Low uptake jeopardizes achievement of major goals set by the World Health Organization in eliminating schistosomiasis. These findings/statistics demonstrate that if schistosomiasis elimination is to be achieved through Mass Drug Administration, then strategies to increase treatment uptake need to be tailored to the

specific community dynamics. This study aimed to assess the level as well as the determinants of MDA uptake in Butiaba sub-county, Buliisa District. Findings from this study could yield critical insight into strategies to tailor MDA interventions in similar settings.

2. Methods

2.1 Study design and setting

A cross sectional study was conducted in which quantitative and qualitative data was collected from adults (18 years and above) in Butiaba sub-county in September 2021. Butiaba is a sub-county within Buliisa district in which schistosomiasis is endemic; the most predominant species being *Schistosoma mansoni*. It is located right at the shores of Lake Albert and the predominant economic activity is fishing. The four parishes that constitute the sub-county have twenty-one villages and an estimated population of 29,181 people as per the 2014 census report from Uganda Bureau of Statistics (Government of Uganda Bureau of Statistics, 2014). However, as of 2020, 11 villages experienced flooding, leading to migration of residents.

MDA campaigns for praziquantel in Butiaba sub-county usually occur biannually but occurred once in 2020 because of the COVID pandemic. The most recent MDA campaign in the sub county had happened in October/November 2020 hence this study happened within one year after this. This study specifically sought to estimate uptake of PZQ among adults during the most recent MDA exercise, identify sociodemographic, knowledge and attitudinal factors associated with this uptake.

2.2 Sampling and Participant selection

Five villages were picked by random selection from a list of the 10 sub county villages using STATA with stratification to ensure equitable representation.

Multistage Probability sampling was used to select households within the identified villages using the sub county registers. Given the targeted sample size of 450 participants, it was determined that 90 adults needed to be interviewed per village (one adult was interviewed per household to allow equal/wider representation.)

For selection of households to visit, a sampling interval was selected by dividing total number of households per village (as reported on village registers provided by the LC1 Chairman) by intended number of households/respondents to be interviewed per village (which was 90). Consecutive numbers were generated using the sample interval and these were used to identify households from the register to visit for interviews. Numbers allocated in village registers were used for this exercise.

Eligible respondents were adults who had lived in Butiaba for 12 months or more and consented to participate in the study. At household level, adults were listed on separate papers and selection was done randomly from a bowl. If a household had no one who met our inclusion criteria, the neighboring household was selected.

The four key informants were however purposively selected with representation from the local leadership, the district vector control office and the MDA parish supervisors to seek for expert opinion on MDA uptake in Butiaba.

2.3 Variable measurement

Both qualitative and quantitative data was collected. Quantitative data was collected as illustrated in table 1 below

Table 1: Study variables

Objective	Variable	Indicator	Means of data collection
To determine the uptake of Mass Drug Administration for schistosomiasis control among adults in Butiaba, Uganda	Uptake of Mass drug administration (dependent)	Self-reported swallowing of praziquantel in last Oct/Nov 2020 MDA campaign	Semi structured questionnaire administered through face to face interviews
To assess the association between socio-demographic determinants and uptake of Mass Drug Administration for schistosomiasis control among adults in Butiaba, Uganda.	Sociodemographic determinants for uptake of MDA for schistosomiasis control (independent)	Age Sex Tribe Occupation Level of education	Semi structured questionnaire administered through face to face interviews
To assess the association between knowledge and uptake of Mass Drug Administration for schistosomiasis control among adults in Butiaba, Uganda.	Knowledge of schistosomiasis and MDA (independent)	Schistosomiasis symptoms schistosomiasis control measures Benefits of MDA	Semi structured questionnaire administered through face to face interviews
To assess the association between attitudes and uptake of Mass Drug Administration for schistosomiasis control among adults in Butiaba, Uganda.	Attitudes towards Mass Drug Administration (independent)	Exposure risk Motivation to take praziquantel Agreement with benefits of MDA Outcome expectancy	Semi structured questionnaire administered through face to face interviews

The semi structured questionnaire (File S1) used was developed based on recommendations from WHO 2016 field guide for implementation of coverage evaluation and pretested (with a Content Validity Index of 0.81). The questionnaire was modified from related literature (Tuhebwe et al., 2015) with minor changes to fit the study's objectives. It was prepared in English and later translated and administered to the respondents in their local language, Alur by trained local interviewers.

The dependent variable was Mass Drug Administration uptake that was determined using self-

reported swallowing of praziquantel distributed in the last October/November 2020 MDA campaign. The percentage uptake represented the fraction of the population sampled that reported having swallowed praziquantel in the most recent October/November 2020 MDA campaign in the sub-county.

The independent variables were knowledge, attitudes and socio-demographic determinants (age, sex, occupation, tribe, educational level) of MDA uptake. Association of attitudes with MDA uptake was determined using questions with answers measured on a five-point Likert scale with values between 1-3 representing negative attitudes and values 4 and 5 representing positive attitudes. Each question had response categories ranging from (1) strongly disagree to (5) strongly agree.

Similarly, association of MDA uptake with knowledge was assessed using a score with 6 variables. Values between 0 and 3 represented low knowledge and 4-6, high levels of knowledge.

For the qualitative data, Focus Group Discussions and Key Informant Interviews were used to identify further hindering factors.

2.4 Data Management and statistical analysis

The required sample size was estimated based on the study first objective which was to determine the uptake of praziquantel during mass drug administration among adults in Butiaba sub-county. We employed a formula by Kish Leslie (Kish, 1965) with an assumption of coverage of approximately 17%, and a design effect of 2 (as recommended by WHO field guide for implementation of PC coverage evaluation surveys, 2016). The sample size attained was

434 participants to estimate uptake with a precision of 5%. We sampled 450 respondents to cater for non-response.

Quantitative data was entered in Epi Data and analyzed in STATA version 12. Data analysis was done at three stages: univariate, bivariate and multivariate and for all statistical analyses, a statistical significance level (α) or p-value of 0.05 was used (95% CI). All the variables considered were informed by our conceptual framework developed based on the modified Anderson model of health service utilization (Andersen, 1995) and are shown in the tables of bivariate analysis.

In univariate analysis, descriptive statistics were obtained for all study variables and presented using standard statistical parameters (frequencies, percentages, means and standard deviations). MDA uptake was computed as the number of respondents who reported swallowing praziquantel in the last October/November 2020 MDA campaign divided by the total number of respondents who answered this question.

In bivariate analysis, chi-square test was used (as is the case with binary and categorical outcomes) to explore association of the respondents' sociodemographic characteristics, knowledge, and attitudes to schistosomiasis and MDA uptake/likelihood of swallowing praziquantel. Interaction effects between the sociodemographic and knowledge variables were also analyzed using the Chi-square test.

In Multivariate Analysis, the binary logistic regression model was used to examine the relationship between MDA uptake and the independent variables. This method was used because the dependent variable is binary in nature (yes/no). Stepwise backwards regression was used to eliminate factors that are not significantly associated with swallowing praziquantel (with a cut off for statistical significance of $P \leq 0.05$) until all independent variables within the model had a

p-value less than 0.05. Crude and adjusted odds ratios with their corresponding confidence intervals and p-values were used to determine association.

Qualitative data was analyzed through thematic content analysis. The data from key informant interviews and the focus group discussions collected using audio recordings was transcribed verbatim and translated to English. To this end, transcripts were read back and forth several times and then coded. Key themes and subthemes related to study objectives were organized in a matrix and discussed as a team for appropriate interpretation. Key quotations that epitomized central themes relating to the health system were identified. Furthermore, analysis of qualitative interview data was done in accordance with the approaches suggested by Krefting (Krefting, 1991) to ensure credibility, applicability, dependability and confirmability of the data.

2.5 Ethical Considerations

Approval for the study was sought and granted by the Institutional Review Board of Clarke International University- Research Ethics Committee (Approval Number: CLARKE-2021-115). Permission to conduct study was sought from Uganda National Council for Science and Technology (Approval Number: SS1083ES).

Written informed consent was sought from each person before the questionnaire or interview was administered. No personal identifiers were collected and furthermore, the study involved people 18 years (the age of consent in Uganda) and above with the ability to exercise free will.

3. Results

A total of 450 respondents sampled from five villages in Butiaba sub-county were interviewed. Three key informant interviews and two focus group discussions (one per parish) were conducted. The most recent MDA campaign in the sub-county had happened in October/November

2020 hence this study happened within one year after this.

3.1 Baseline characteristics of study participants

The socio-demographic characteristics of the respondents that participated in the questionnaire interviews are shown in Table 2. Over half (52.7%) of the respondents were female. Respondents' age ranged from 18 to 80 with a mean age of 34 years (SD = ± 11.4). Most respondents (60.89%) had received at least a primary education with majority (37.56%) being business people and the second most common occupation being fishing.

Table 2: Socio-demographic characteristics of 450 adults from Butiaba Sub-county, Buliisa District surveyed in September 2021

Variable	Number (N=450)	Percentage (%)
Age group(years)		
18-24	106	23.56
25-29	75	16.67
30-39	153	34.00
40+	116	25.78
<i>Total</i>	450	100
Sex		
Male	213	47.33
Female	237	52.67
<i>Total</i>	450	100
Tribe		
Alur	366	81.33
Bagungu	43	9.55
Banyoro	22	4.89
Others	19	4.21
<i>Total</i>	450	100
Duration of residence at current address		
1-10	124	27.56
11-20	131	29.11
>20	195	43.33
<i>Total</i>	450	100
Highest level of education		
None	85	18.89
Primary	274	60.89
Lower secondary	75	16.67
Upper secondary	13	2.89
University	3	0.67

<i>Total</i>	450	100
--------------	-----	-----

As shown in table 3, majority of the respondents (96%) reported using latrines as place of ease with these being majorly shared/neighborhood latrines (47.98%). The most common source of water reported was the lake at 82.44%.

Table 3: Sanitation characteristics of 450 adults from Butiaba Sub-county, Buliisa District surveyed in September 2021

Variable	Number (N=450)	Percentage (%)
Place of ease		
Latrine	432	96.00
Bush	15	3.33
Lake	3	0.67
<i>Total</i>	450	100
Location of toilet used		
Household	209	46.44
Shared	216	47.98
Public	11	2.44
<i>Total</i>	450	100
Main source of water		
Lake	371	82.44
Tap	73	16.22
Rainfall	2	0.44
Borehole	2	0.44
Other	2	0.44
<i>Total</i>	450	100

Uptake of praziquantel during MDA for schistosomiasis control

In all the five sampled villages, distribution of praziquantel was conducted in the 2020 MDA exercise. Of the 450 respondents, 322 respondents i.e., 71.56% (95% CI: 67.14 – 75.68) reported that they had swallowed the drug in the 2020 MDA exercise. 5.78 % (26/450) of the participants reported having never swallowed praziquantel in their lifetime and 21.3% (96/450) of the

participants hadn't swallowed praziquantel within the last 10 years. Overall, 84.73% of respondents reported that they were willing to swallow praziquantel in the next MDA exercise with majority (34.65%) citing schistosomiasis prevention as reason.

Factors associated with reported uptake of praziquantel in the bivariate analysis are shown in Tables 4,5 and 6. Female respondents were less likely to swallow Praziquantel (OR=0.52, 95% CI: 0.34–0.79) and being 30 years and above had a positive association with uptake of mass treatment: 30-39 years (OR= 2.31, 95% CI: 1.35-3.95), 40+ years (OR= 2.86, 95% CI: 1.58-5.22). Being in the fishing occupation also had a positive association with uptake of praziquantel during MDA (OR =2.72, 95% CI: 1.41-5.24).

Table 4: Bivariate Association between Socio-demographic characteristics and uptake of praziquantel in 450 participants from Butiaba sub-county, Buliisa District

Variable	Uptake of MDA		Unadjusted OR 95% CI	P-value
	Yes (n=322) N(%)	No (n=128) N(%)		
Age group(years)				
18-24	62(13.8)	44(9.8)	1.00	
25-29	50(11.1)	25(5.6)	1.42(0.77-2.63)	0.265
30-39	117(26)	36(8)	2.31(1.35-3.95)	0.002
40+	93(20.7)	23(5.1)	2.86(1.58-5.22)	0.001
<i>Total</i>	322(71.56)	128(28.4)		
Sex				
Male	167(37.1)	46(10.2)	1.00	
Female	155(34.4)	82(18.2)	0.52(0.34-0.79)	0.002
<i>Total</i>	322(71.56)	128(28.4)		
Length of stay in Butiaba				
1-10	75(16.7)	49(10.9)	1.00	
11-20	90(20)	41(9.9)	1.43(0.86-2.4)	0.17
>20	157(34.9)	38(8.4)	2.69(1.63-4.47)	0.00
<i>Total</i>	322(71.56)	128(28.4)		
Highest level of education				
None	61(13.6)	24(5.3)	1.00	
Primary	200(44.4)	74(16.4)	1.06(0.62-1.83)	0.82
Lower secondary	50(11.1)	25(5.5)	0.79(0.40-1.54)	0.49
Upper secondary	10(2.2)	3(0.7)	1.31(0.33-5.18)	0.70
University	1(0.2)	2(0.4)	0.20(0.02-2.27)	0.19
<i>Total</i>	322(71.56)	128(28.4)		

Major occupation				
Unemployed	35(7.8)	24(5.3)	1.00	
Fishing	119(26.4)	30(7.1)	2.72(1.41-5.24)	0.003
Office	4(0.9)	1(0.2)	2.74(0.29-26.08)	0.38
Business	119(26.4)	50(11.1)	1.63(0.88-3.02)	0.12
Farming	15(3.75)	4(0.9)	2.57(0.76-8.7)	0.13
Other	30(6.7)	19(4.2)	1.08(0.49-2.35)	0.84
<i>Total</i>	322(71.56)	128(28.4)		

As shown in table 5, knowledge about Bilharzia was positively associated with uptake of MDA. Respondents who were not knowledgeable about bilharzia cause (OR= 0.19, 95% CI: 0.09-0.41), signs (OR= 0.18, 95% CI: 0.08-0.39), and control measures (OR= 0.27, 95% CI: 0.08-0.86), were less likely to swallow Praziquantel during MDA exercise.

With regard to attitude related factors (table 6), bivariate analysis revealed that respondents who perceived a high personal risk of catching Bilharzia (OR=2.02, 95% CI:1.21-3.38) & agreed that MDA improves one's health (OR=2.51, 95% CI:0.99–6.36) were more likely to swallow PZQ.

Table 5: Bivariate Association between knowledge related factors and uptake of praziquantel of 450 adults from Butiaba Sub-county, Buliisa District surveyed in September 2021

Variable	Uptake of MDA		Unadjusted OR 95% CI	P-value
	Yes(n=322) N(%)	No(n=128) N(%)		
Knowledge of cause				
Ways one can get Bilharzia				
Knowledgeable	310(68.9)	107(23.8)	1.00	
Not knowledgeable	12(2.7)	21(4.7)	0.19(0.09-0.41)	0.00
<i>Total</i>	322(71.56)	128(28.4)		
Knowledge of signs				
Bilharzia signs				
Knowledgeable	311(69.1)	107(23.8)	1.00	
Not knowledgeable	11(2.4)	21(4.7)	0.18(0.08-0.39)	0.00
<i>Total</i>	322(71.56)	128(28.4)		
Knowledge of control measures				
Bilharzia control measures				
Knowledgeable	317(70.4)	121(26.9)	1.00	
Not knowledgeable	5(1.1)	7(1.6)	0.27(0.08-0.86)	0.029

<i>Total</i>	322(71.56)	128(28.4)		
Drug control of Bilharzia				
Swallowing MDA PZQ controls Bilharzia				
Knowledgeable	245(54.4)	93(20.7)	1.00	
Not knowledgeable	77(17.1)	35(7.8)	0.83(0.52-1.33)	0.45
<i>Total</i>	322(71.56)	128(28.4)		

Table 6: Bivariate Association between attitudinal factors and uptake of praziquantel in 450 adults from Butiaba Sub-county, Buliisa District surveyed in September 2021

Variable	Uptake of MDA		Unadjusted OR 95% CI	P-value
	Yes (n=322) N(%)	No (n=128) N(%)		
Attitude towards risk				
Personal risk of catching Bilharzia				
Low	44(9.8)	31(6.9)	1.00	
High	278(61.8)	97(21.5)	2.02(1.21-3.38)	0.007
<i>Total</i>	322(71.56)	128(28.4)		
Attitude towards severity				
Bilharzia can cause death				
Disagree	4(0.9)	1(0.2)	1.00	
I don't know	9(2)	8(1.8)	0.28(0.02-3.07)	0.298
Agree	309(68.7)	119(26.4)	0.65(0.07-5.87)	0.7
<i>Total</i>	322(71.56)	128(28.4)		
Attitude towards MDA need				
Everyone should take PZQ in MDA				
Disagree	2(0.4)	1(0.2)	1.00	
I don't know	16(3.6)	12(2.7)	0.67(0.05-8.23)	0.75
Agree	304(67.6)	115(25.6)	1.32(0.12-4.71)	0.82
<i>Total</i>	322(71.56)	128(28.4)		
Perceived benefits				
MDA controls Bilharzia effectively				
Disagree	32(7.1)	7(1.6)	1.00	
I don't know	11(2.4)	13(2.9)	0.19(0.05-0.58)	0.004
Agree	279(62)	108(24)	0.57(0.24-1.31)	0.187
<i>Total</i>	322(71.56)	128(28.4)		
MDA improves one's health				
Disagree	10(2.2)	9(2)	1.00	

I don't know	13(2.9)	12(2.7)	0.98(0.29-3.22)	0.97
Agree	299(66.4)	107(23.8)	2.51(0.99-6.36)	0.05
<i>Total</i>	322(71.56)	128(28.4)		
Perceived barriers				
PZQ tastes bad				
Disagree	139(30.9)	61(13.5)	1.00	
I don't know	16(3.6)	13(2.9)	0.54(0.24-1.19)	0.13
Agree	167(37.1)	54(12)	1.36(0.88-2.08)	0.17
MDA can cause death or bad effects				
Disagree	197(43.8)	80(17.8)	1.00	
I don't know	13(2.9)	8(1.8)	0.66(0.26-1.66)	0.38
Agree	112(24.9)	40(8.9)	1.14(0.73-1.77)	0.57
Medicine isn't enough during MDA				
Disagree	80(17.8)	35(7.8)	1.00	
I don't know	42(9.3)	33(7.3)	0.56(0.3-1.01)	0.06
Agree	200(44.4)	60(13.3)	1.46(0.89-2.38)	0.13
<i>Total</i>	322(71.56)	128(28.4)		

In multivariate analysis using the Logistic regression model, variables that were considered included age, sex, length of stay, occupation, knowledge of bilharzia signs, causes, control measures, perceived bilharzia risk, effectiveness of MDA in controlling Bilharzia & improving health, taste of praziquantel and sufficiency of praziquantel during MDA campaigns (as indicated in the tables of bivariate analysis above).

Of those (table 7), only age and knowledge about schistosomiasis symptoms were found to be significant predictors for the uptake of Praziquantel during MDA. Belonging to the age group 30 years & above and having knowledge about signs of schistosomiasis were positively associated with uptake of praziquantel during MDA.

Table 7: Significant factors associated with uptake of praziquantelin 450 adults from Butiaba Sub-county, Buliisa District surveyed in September 2021

Variable	Uptake of MDA		Unadjusted OR 95% CI	Adjusted OR 95% CI	P-value AOR
	Yes (n=322) N(%)	No (n=128) N(%)			

Age group(years)					
18-24	62(13.8)	44(9.8)	1.00		
30-39	117(26)	36(8)	2.31(1.35-3.95)	2.35(1.34-4.09)	0.003
40+	93(20.7)	23(5.1)	2.86(1.58-5.22)	2.68(1.45-4.95)	0.002
<i>Total</i>	322(71.56)	128(28.4)			
Knowledge of signs					
Bilharzia signs					
Knowledgeable	311(69.1)	107(23.8)	1.00		
Not knowledgeable	11(2.4)	21(4.7)	0.18(0.08-0.39)	0.00	0.00
<i>Total</i>	322(71.56)	128(28.4)			

3.2 Hindering factors associated with uptake of praziquantel

Hindering factors were further highlighted through the Focus group discussions and Key Informant Interviews. All the participants of the Focus Group discussions from the two parishes had been residents of Butiaba for at least 2 years and all the 3 key informants had participated in conducting MDA exercises within Butiaba sub county.

The major themes cited in both Focus Group Discussions included: matters relating to drug acceptability (such as side effects, size & smell), poor sensitization, poor mobilization, irregular supply and inadequacy of tablets during the MDA exercises. *“From my experience, the side effects of the drug are too severe moreover the tablets smell so bad and require one to eat heavily before swallowing. Another reason is inadequacy of tablets for example I missed in the last MDA exercise. It demoralizes me to participate because I know that they always come in small quantities so I may miss again. It makes me lose interest in taking the tablet.”* (Bugoigo FGD participant)

Of the three key informants, two cited financial constraints as a major challenge in delivering MDA for Bilharzia in the sub county. One key informant said:

“There’s a lot of resistance from the VHTs because of lack of facilitation vis-a-vis the large amount of work. We pull ropes with them and keep on persuading with a few incentives, but the

major issue is lack of facilitation to do the work.” (KI: district vector control office)

Another challenge that was cited was the inadequacy of drugs. “*The amount of medicine brought to the community is low compared to the data that we present on the population.*” (KI: parish supervisor). This was highlighted as a demoralizer to participate in Bilharzia MDA within the community.

Additionally, infective sensitization was cited as a major problem in delivery of MDA for Bilharzia services. “*The community lacks knowledge on MDA. Local Leaders and elders need to be utilized as well to deliver the right message to the community.*” (KI: local leader)

Another challenge identified was constant migration. “*The community at the landing site is mostly a moving community: today someone is in Sonsio, tomorrow Runga. People are registered for praziquantel in one area and are not in the same place when the MDA exercise starts. This hinders uptake of Praziquantel.*” (KI: parish supervisor)

As illustrated above, 5 major themes were cited in the qualitative interviews including: financial constraint, poor sensitization & mobilization, irregular and inadequate drug supply, constant migration, and matters relating to drug acceptability (such as side effects, size & smell).

4. Discussion

4.1 Uptake of MDA

Although uptake of praziquantel during MDA in Butiabasub county was high at 71.56%, it was still below the WHO target of 75% (Organization, 2014). According to the World Health Organization, it is important to attain at least the target of 75% coverage so as to avoid high intensity of infection and eliminate schistosomiasis as a public health threat by

2025(Organization, 2014). With uptake below the target, this cannot be achieved. Lower than recommended drug uptake rates have in the past been reported in different parts of Uganda including Koomes Island(Tuhebwe et al., 2015) and Mayuge(Adriko et al., 2018). Such low drug uptake as reported could considerably prolong the time needed to reach the goal of schistosomiasis transmission elimination. It is thus important to identify barriers to optimal drug uptake rates so as to improve them and also to identify enabling factors that could further be strengthened to ensure successful schistosomiasis elimination.

Analysis of data in this study also revealed that 5.78% of the participants reported having never swallowed praziquantel in their lifetime and 21.3% of participants reporting having not swallowed the drug in the last 10years. A number of studies have in the past focused on recent treatment coverage(Chami et al., 2017; Knopp et al., 2016; Tuhebwe et al., 2015), but have not looked at lifetime treatment data and the important issue of systematic non-treatment rates(Shuford, Turner, & Anderson, 2016; Turner et al., 2017). It is good to note that the greater percentage of participants in this study had been treated at least once in 18 years of MDA. Using such data to develop models could have a significant effect on the predicted duration that repeated MDA is needed to control morbidity. Using this study's noncompliance rate of 21.3% within the last 10 years, the sensitivity analyses by Turner and colleagues (2017) indicate that MDA is needed for 6–9 years with a 20% systematic non-compliance(Turner et al., 2017). If these lower than target annual treatment rates and high lifetime MDA non-treatment rates continue, MDA programs in districts such as this may potentially never attain the goals of schistosomiasis elimination. The closeness of percentages obtained in this study to the set targets is however encouraging that once efforts are geared in the right direction, goals could be achieved.

4.2 Socio demographic factors associated with uptake of praziquantel during MDA

There was a strong association between age and uptake of praziquantel during MDA. This is in agreement with other studies that young adults are less likely to swallow praziquantel during MDA exercises. (Adriko et al., 2018; Abdoulaye Dabo, Boubacar Bary, Bourema Kouriba, Oumar Sankaré, & Ogobara Doumbo, 2013; D. W. Njomo, Kibe, et al., 2020; Parker, Allen, & Hastings, 2008b; Tuhebwe et al., 2015). In a fishing community like Butiaba, this is possibly because younger adults spend more time at the lake shores and outdoors and are therefore more likely to miss out on door to door and even centralized drug distributions. This similarity suggests a need for further research to understand the reasons behind this trend revealed by the different studies. Additionally, if younger adults are more likely to reject free treatment than their older counterparts, then there is a need to tailor MDA exercises targeted at involving them more.

This study did not find a significant association between occupation and uptake of praziquantel during MDA as demonstrated by some studies (Tuhebwe et al., 2015). A study done on a lake Victoria island linked being a fisher folk to a greater likelihood to swallow praziquantel but that wasn't the case in this study. It is possible that trends differ along different lake shores.

4.3 Knowledge related factors associated with uptake of praziquantel

In line with knowledge, the results show that study participants who were knowledgeable about signs of schistosomiasis were more likely to swallow praziquantel. This observation correlates with other studies (Adeneye et al., 2007; Fleming et al., 2009; Odhiambo, Musuva, Odiere, & Mwinzi, 2016) that had similar findings. Knowledge of schistosomiasis symptoms possibly implies that the residents have had prior experience with the disease in their community and are hence more aware of their exposure risk and more willing to swallow the drug to protect themselves. However, it should be noted that knowledge collectively doesn't always positively

correlate with increased uptake of the drug as shown by some studies (Muhumuza, Olsen, Nuwaha, & Katahoire, 2015; Parker et al., 2008b) because of barriers like fear of praziquantel side effects.

A majority of the participants had heard about schistosomiasis as is expected in highly endemic areas that have been treated repeatedly for a number of years. Similarly, in studies done in endemic parts of Africa, more than 90% of the study populations have been observed to have an awareness of schistosomiasis (Maseko, Mkhonta, Masuku, Dlamini, & Fan, 2018; Mwai, Njenga, & Barasa, 2016; Sady et al., 2015). However, awareness is often limited to just being familiar with the disease name. In this study, a strong correlation was found between knowledge of symptoms and uptake of the drug. Poor awareness of signs and symptoms was also observed to impact praziquantel uptake in other studies (Acka et al., 2010; Dawaki et al., 2015) which could in turn deter elimination of schistosomiasis.

Furthermore, although greater than 50% of the respondents were knowledgeable on the cause, signs and control measures of schistosomiasis, the remaining unknowledgeable population evidences a need for intensive health education about schistosomiasis and its control. Health education is a major control strategy acting as a founding block on which other strategies can flourish. If health education is inadequate, this may result in rejection of free drugs during mass drug administration (Hastings, 2016; Tilak Chandra Nath et al., 2020; D. W. Njomo, Kimani, et al., 2020). Health education on both schistosomiasis and the importance of praziquantel during MDAs is beneficial in enabling community participation and changing perceptions & beliefs of the community towards given control measures (Wang et al., 2018).

4.3 Attitudinal factors associated with uptake of praziquantel

There was no significant association between attitude towards personal risk of exposure to schistosomiasis or its severity and the uptake of praziquantel. This contradicts with the health belief model which suggests that when a person perceives themselves to be at risk of acquiring the disease and the benefits of the health intervention are known, the intervention will be complied with (Champion & Skinner, 2008). Furthermore, perception that MDA improves health and controls schistosomiasis effectively did not significantly impact uptake of praziquantel during MDA. This could possibly be because of the absence of a major spiritual belief in the community that could drive negative attitudes towards the disease and MDA exercise.

In spite of the bad taste and severe side effects of praziquantel, personal attitudes towards this were not negatively associated with people swallowing the drug. Our results, similar to other studies (A. Dabo, B. Bary, B. Kouriba, O. Sankaré, & O. Doumbo, 2013; Knopp et al., 2016), show that the personal perceptions and fear of side effects or taste of praziquantel may not necessarily influence uptake of the drug. This could be because the negative barriers of taste and side effects are negligible in comparison with the drug benefits (Putri, Diana, Daris, Afriana, & Hidayat, 2021). In conflict with these findings, other studies (Koudessi, 2020; Muhumuza, Olsen, Katahoire, & Nuwaha, 2013; Omedo et al., 2012) show that awareness of side effects and bad taste is negatively associated with uptake of drugs during MDA exercise. However, given that this was a cross-sectional study, it is possible that this finding was because people who had swallowed the drugs had potentially experienced the side effects and the bad taste and hence were more knowledgeable about them. It is beneficial to work towards alleviating praziquantel side effects when possible.

4.4 Hindering factors associated with uptake of praziquantel

Additional hindering factors included operational and service delivery challenges highlighted through the qualitative interviews such as financial constraint hence low Village Health Team (VHT) motivation, poor sensitization & mobilization, irregular and inadequate drug supply that in turn affected uptake of praziquantel during MDA in this study.

VHTs and local leaders participated in the MDA exercises within the sampled Butiavillages which illustrates positive community involvement. This is very vital in ensuring that the larger community supports the MDA exercise and hence positively impacts on the drug uptake (Mo, 2010; Ssali et al., 2021). However, as shown in the results, uptake was lower than recommended by WHO perhaps because of low motivation amongst the VHTs in the absence of facilitation. Furthermore, VHTs lacked training and adequate supplies to execute their duties.

VHTs that are not facilitated to execute their duties may lack the motivation to accomplish their tasks. Additionally, absence of training results in lack the knowledge and skills to adequately sensitize the community. Together, these can result in failure to achieve set goals with drug uptake during MDA (D. Njomo, Amuyunzu-Nyamongo, Mukoko, Magambo, & Njenga, 2012; Tuhebwe et al., 2015). There is therefore a need to train VHTs, provide adequate facilitation and supplies for them to execute their duties as per the Uganda Health Sector Strategic Plan.

Other challenges highlighted included poor sensitization, irregular and inadequate drug supply. These need to be solved to enable the Uganda national bilharzia control program (UNBCP) to achieve the objective of creating a local demand for mass treatment (Narcis B Kabatereine et al., 2006). Drug supply that is inadequate and irregular has been shown to negatively impact on the uptake of MDA programs (Krentel, Fischer, & Weil, 2013; Tilak C Nath, Padmawati,

&Murhandarwati, 2019)by demoralizing people from participating in the MDA exercises hence effort needs to be geared towards availing a constant and adequate supply.

Conclusively, the results show that sociodemographic factors likeage and enabling factors like knowledge about schistosomiasis increase the likelihood of praziquantel uptake during MDA in Butiaba. This information could be used to improve the national bilharzia control programin implementing effective mass drug administration.

4.5 Limitations of the study

Two major limitations were incurred. Firstly, there is inability to generalize findings since the study only looked at participants from one area (sub county) along Lake Albert shores.

Secondly, this was a cross-sectional study hence is not able to determine causality between identified factors and uptake of praziquantel.

5. Conclusions and Recommendations

5.1 Conclusions

Uptake of praziquantel during MDA in Butiaba was high but still below the target rate of 75% set by WHO. Limited knowledge on schistosomiasis symptoms, being in a young adult age group (18-29) and irregular and/or inadequate drug supply are some of the factors associated with uptake being below target.

5.2 Recommendations

- Rigorous health educationsupervised by local leaders and the Ministry of Health needs to be done in order to share information on schistosomiasis and MDA.
- Adequate and regular praziquantel supply by the Ministry of Health as inadequacy of the

drug was cited as a major demoralizer to participate in MDA exercise.

- A strategic approach to drug distribution involving distribution of praziquantel at youth hot spots (such as lake shores, dancing halls) with prior mobilization done by fellow youth leaders to target the young adults. In addition, the proposed strategy should constantly monitor the program performance and make the required changes as guided by the Quality of Care model (Organization, 2006) developed by the World Health Organization. This model provides a simple framework for engaging the various stakeholders, including representatives from different parts of the community, in the planning and implementation of an intervention whilst ensuring a predefined level of quality. By using such systematic approach and by involving local decision makers, program planners, drug distributors and community members, the MDA programs targeting schistosomiasis in Butiaba should be able to increase praziquantel uptake rates and hence attain the target of elimination of the disease as a public health problem.
- Further research is needed to understand why the young adults are less likely to swallow praziquantel during MDA exercise.

References

- Acka, C. A., Raso, G., N'goran, E. K., Tschannen, A. B., Bogoch, I. I., Séraphin, E., . . . Utzinger, J. (2010). Parasitic worms: knowledge, attitudes, and practices in western Côte d'Ivoire with implications for integrated control. *PLoS neglected tropical diseases*, 4(12), e910.
- Adeneye, A., Akinwale, O., Idowu, E., Adewale, B., Manafa, O., Sulyman, M., . . . Appelt, B. (2007). Sociocultural aspects of mass delivery of praziquantel in schistosomiasis control: The Abeokuta experience. *Research in Social and Administrative Pharmacy*, 3(2), 183-198.
- Adenowo, A. F., Oyinloye, B. E., Ogunyinka, B. I., & Kappo, A. P. (2015). Impact of human schistosomiasis in sub-Saharan Africa. *The Brazilian Journal of Infectious Diseases*, 19(2), 196-205. doi:<https://doi.org/10.1016/j.bjid.2014.11.004>
- Adriko, M., Faust, C. L., Carruthers, L. V., Moses, A., Tukahebwa, E. M., & Lambertson, P. H. (2018). Low praziquantel treatment coverage for *Schistosoma mansoni* in Mayuge District, Uganda, due to the absence of treatment opportunities, rather than systematic non-compliance. *Tropical medicine and infectious disease*, 3(4), 111.
- Andersen, R. M. (1995). Revisiting the behavioral model and access to medical care: does it matter? *Journal of health and social behavior*, 1-10.
- Chami, G. F., Kontoleon, A. A., Bulte, E., Fenwick, A., Kabatereine, N. B., Tukahebwa, E. M., & Dunne, D. W. (2017). Community-directed mass drug administration is undermined by status seeking in friendship networks and inadequate trust in health advice networks. *Soc Sci Med*, 183, 37-47. doi:10.1016/j.socscimed.2017.04.009
- Champion, V. L., & Skinner, C. S. (2008). The health belief model. *Health behavior and health education: Theory, research, and practice*, 4, 45-65.
- Dabo, A., Bary, B., Kouriba, B., Sankaré, O., & Doumbo, O. (2013). Factors associated with coverage of praziquantel for schistosomiasis control in the community-direct intervention (CDI) approach in Mali (West Africa). *Infect Dis Poverty*, 2(1), 11. doi:10.1186/2049-9957-2-11
- Dabo, A., Bary, B., Kouriba, B., Sankaré, O., & Doumbo, O. (2013). Factors associated with coverage of praziquantel for schistosomiasis control in the community-direct intervention (CDI) approach in Mali (West Africa). *Infectious diseases of poverty*, 2(1), 1-11.
- Dawaki, S., Al-Mekhlafi, H. M., Ithoi, I., Ibrahim, J., Abdulsalam, A. M., Ahmed, A., . . . Atroosh, W. M. (2015). The menace of schistosomiasis in Nigeria: knowledge, attitude, and practices regarding schistosomiasis among rural communities in Kano state. *PLoS One*, 10(11), e0143667.
- Emmanuel, I., Aginya, O., & Doehring, E. (2008). Epidemiology, of bilharzias (schistosomiasis) in Uganda from 1902 until 2005. *African health sciences*, 8(4), 239-243.
- Exum, N. G., Kibira, S. P., Ssenyonga, R., Nobili, J., Shannon, A. K., Ssempebwa, J. C., . . . Makumbi, F. E. (2019). The prevalence of schistosomiasis in Uganda: A nationally representative population estimate to inform control programs and water and sanitation interventions. *PLoS neglected tropical diseases*, 13(8), e0007617.
- Fleming, F., Fenwick, A., Tukahebwa, E., Lubanga, R., Namwangye, H., Zaramba, S., & Kabatereine, N. (2009). Process evaluation of schistosomiasis control in Uganda, 2003 to 2006: perceptions, attitudes and constraints of a national programme. *Parasitology*, 136(13), 1759-1769.
- Government_of_Uganda_Bureau_of_Statistics. (2014). National Population and Housing Census

2014. Retrieved from https://www.ubos.org/wp-content/uploads/publications/03_20182014_National_Census_Main_Report.pdf
- Government_of_Uganda_Ministry_of_Health. (2010). Health Sector Strategic Plan III 2010/11-2014/15. Retrieved from https://www.health.go.ug/docs/HSSP_III_2010.pdf
- Hastings, J. (2016). Rumours, riots and the rejection of mass drug administration for the treatment of schistosomiasis in Morogoro, Tanzania. *Journal of biosocial science*, 48(S1), S16-S39.
- Kabatereine, N., Kemijumbi, J., Ouma, J. H., Kariuki, H., Richter, J., Kadzo, H., . . . Vennervald, B. (2005). Epidemiology and morbidity of *Schistosoma mansoni* infection in a fishing community along Lake Albert in Uganda. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 98, 711-718. doi:10.1016/j.trstmh.2004.06.003
- Kabatereine, N. B., Tukahebwa, E., Kazibwe, F., Namwangye, H., Zaramba, S., Brooker, S., . . . Webster, J. P. (2006). Progress towards countrywide control of schistosomiasis and soil-transmitted helminthiasis in Uganda. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100(3), 208-215.
- Kabatereine, N. B., Tukahebwa, E., Kazibwe, F., Namwangye, H., Zaramba, S., Brooker, S., . . . Fenwick, A. (2006). Progress towards countrywide control of schistosomiasis and soil-transmitted helminthiasis in Uganda. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100(3), 208-215. doi:10.1016/j.trstmh.2005.03.015
- Kish, L. (1965). *Survey sampling*.
- Knopp, S., Person, B., Ame, S. M., Ali, S. M., Muhsin, J., Juma, S., . . . Rollinson, D. (2016). Praziquantel coverage in schools and communities targeted for the elimination of urogenital schistosomiasis in Zanzibar: a cross-sectional survey. *Parasit Vectors*, 9, 5. doi:10.1186/s13071-015-1244-0
- Koudessi, C. F. E. (2020). *Uptake of Praziquantel Mass-Drug Administration for Schistosomiasis Control Among School-Age Children in Kpando Municipality in the Volta Region of Ghana*. University of Ghana,
- Krefting, L. (1991). Rigor in qualitative research: The assessment of trustworthiness. *American journal of occupational therapy*, 45(3), 214-222.
- Krentel, A., Fischer, P. U., & Weil, G. J. (2013). A review of factors that influence individual compliance with mass drug administration for elimination of lymphatic filariasis. *PLoS neglected tropical diseases*, 7(11), e2447.
- Loewenberg, S. (2014). Uganda's struggle with schistosomiasis. *The Lancet*, 383(9930), 1707-1708.
- Maseko, T. S., Mkhonta, N. R., Masuku, S. K., Dlamini, S. V., & Fan, C.-K. (2018). Schistosomiasis knowledge, attitude, practices, and associated factors among primary school children in the Siphofaneni area in the Lowveld of Swaziland. *Journal of microbiology, immunology and infection*, 51(1), 103-109.
- Mo, H. (2010). Village Health Team: Strategy and Operational Guidelines. *Kampala, Uganda: Uganda Ministry of Health*.
- Muhumuza, S., Olsen, A., Katahoire, A., & Nuwaha, F. (2013). Uptake of preventive treatment for intestinal schistosomiasis among school children in Jinja district, Uganda: a cross sectional study. *PLoS One*, 8(5), e63438.
- Muhumuza, S., Olsen, A., Nuwaha, F., & Katahoire, A. (2015). Understanding low uptake of mass treatment for intestinal schistosomiasis among school children: a qualitative study in Jinja district, Uganda. *Journal of biosocial science*, 47(4), 505-520.

- Mwai, J., Njenga, S., & Barasa, M. (2016). RETRACTED ARTICLE: Knowledge, attitude and practices in relation to prevention and control of schistosomiasis infection in Mwea Kirinyaga county, Kenya. *BMC public health*, *16*(1), 1-11.
- Nath, T. C., Adnan, M. R., Sultana, N., Husna, A., Ndossi, B. A., Kang, Y., . . . Lee, D. (2020). Integration of health education intervention to improve the compliance to mass drug administration for soil-transmitted helminths infection in Bangladesh: An implementation research. *Parasite Epidemiology and Control*, *11*, e00165.
- Nath, T. C., Padmawati, R. S., & Murhandarwati, E. H. (2019). Barriers and gaps in utilization and coverage of mass drug administration program against soil-transmitted helminth infection in Bangladesh: An implementation research. *Journal of infection and public health*, *12*(2), 205-212.
- Njomo, D., Amuyunzu-Nyamongo, M., Mukoko, D., Magambo, J., & Njenga, S. (2012). Social mobilization and compliance with mass treatment for lymphatic filariasis elimination in Kenya. *African Journal of Health Sciences*, *20*(1-2), 42-49.
- Njomo, D. W., Kibe, L. W., Kimani, B. W., Okoyo, C., Omondi, W. P., & Sultani, H. M. (2020). Addressing barriers of community participation and access to mass drug administration for lymphatic filariasis elimination in Coastal Kenya using a participatory approach. *PLoS neglected tropical diseases*, *14*(9), e0008499.
- Njomo, D. W., Kimani, B. W., Kibe, L. W., Okoyo, C., Omondi, W. P., & Sultani, H. M. (2020). Implementation challenges and opportunities for improved mass treatment uptake for lymphatic filariasis elimination: Perceptions and experiences of community drug distributors of coastal Kenya. *PLoS neglected tropical diseases*, *14*(12), e0009012.
- Odhiambo, G. O., Musuva, R. M., Odiere, M. R., & Mwinzi, P. N. (2016). Experiences and perspectives of community health workers from implementing treatment for schistosomiasis using the community directed intervention strategy in an informal settlement in Kisumu City, western Kenya. *BMC public health*, *16*(1), 1-12.
- Olveda, D. U., Li, Y., Olveda, R. M., Lam, A. K., Chau, T. N., Harn, D. A., . . . Ross, A. G. (2013). Bilharzia: pathology, diagnosis, management and control. *Tropical medicine & surgery*, *1*(4).
- Omedo, M. O., Matey, E. J., Awiti, A., Ogutu, M., Alaii, J., Karanja, D. M., . . . Mwinzi, P. N. (2012). Community health workers' experiences and perspectives on mass drug administration for schistosomiasis control in western Kenya: the SCORE Project. *The American Journal of Tropical Medicine and Hygiene*, *87*(6), 1065.
- Organization, W. H. (2006). *Quality of care: a process for making strategic choices in health systems*: World Health Organization.
- Organization, W. H. (2012). *Accelerating work to overcome the global impact of neglected tropical diseases: a roadmap for implementation*. Retrieved from
- Organization, W. H. (2014). Schistosomiasis, fact sheet no 115. *Geneva: World Health Organization*.
- Parker, M., Allen, T., & Hastings, J. (2008a). Resisting control of neglected tropical diseases: dilemmas in the mass treatment of schistosomiasis and soil-transmitted helminths in north-west Uganda.
- Parker, M., Allen, T., & Hastings, J. (2008b). Resisting control of neglected tropical diseases: dilemmas in the mass treatment of schistosomiasis and soil-transmitted helminths in north-west Uganda. *Journal of biosocial science*, *40*(2), 161-181.
- Parker, M., Allen, T., Pearson, G., Peach, N., Flynn, R., & Rees, N. (2012). Border parasites:

- schistosomiasis control among Uganda's fisherfolk. *Journal of Eastern African Studies*, 6(1), 98-123.
- Putri, A. S. D., Diana, T. V., Daris, R., Afriana, F., & Hidayat, S. H. (2021). Does the presence of praziquantel-related adverse events affect the health community's perception toward the mass chemopreventive program in the highest prevalence area of Schistosomiasis in Indonesia? Study in Dodolo village, Napu Valley, Central Sulawesi. *Gaceta Sanitaria*, 35, S487-S490.
- Ross, A. G. P., Chau, T. N., Inobaya, M. T., Olveda, R. M., Li, Y., & Harn, D. A. (2017). A new global strategy for the elimination of schistosomiasis. *International Journal of Infectious Diseases*, 54, 130-137. doi:<https://doi.org/10.1016/j.ijid.2016.09.023>
- Sady, H., Al-Mekhlafi, H. M., Atroosh, W. M., Al-Delaimy, A. K., Nasr, N. A., Dawaki, S., . . . Chua, K. H. (2015). Knowledge, attitude, and practices towards schistosomiasis among rural population in Yemen. *Parasites & vectors*, 8(1), 1-13.
- Shuford, K. V., Turner, H. C., & Anderson, R. M. (2016). Compliance with anthelmintic treatment in the neglected tropical diseases control programmes: a systematic review. *Parasit Vectors*, 9, 29. doi:10.1186/s13071-016-1311-1
- Ssali, A., Pickering, L., Nalwadda, E., Mujumbusi, L., Seeley, J., & Lamberton, P. H. (2021). Schistosomiasis messaging in endemic communities: Lessons and implications for interventions from rural Uganda, a rapid ethnographic assessment study. *PLoS neglected tropical diseases*, 15(10), e0009893.
- Tuhebwe, D., Bagonza, J., Kiracho, E. E., Yeka, A., Elliott, A. M., & Nuwaha, F. (2015). Uptake of mass drug administration programme for schistosomiasis control in Koome Islands, Central Uganda. *PLoS One*, 10(4), e0123673.
- Turner, H. C., Truscott, J. E., Bettis, A. A., Farrell, S. H., Deol, A. K., Whitton, J. M., . . . Anderson, R. M. (2017). Evaluating the variation in the projected benefit of community-wide mass treatment for schistosomiasis: Implications for future economic evaluations. *Parasit Vectors*, 10(1), 213. doi:10.1186/s13071-017-2141-5
- Wang, M., Han, X., Fang, H., Xu, C., Lin, X., Xia, S., . . . Tao, H. (2018). Impact of health education on knowledge and behaviors toward infectious diseases among students in Gansu Province, China. *BioMed research international*, 2018.
- WHO. (2022). Schistosomiasis Factsheet. Retrieved from <https://www.who.int/news-room/factsheets/detail/schistosomiasis>

WHO/Department of control of neglected tropical diseases (2017). Schistosomiasis and soil-transmitted helminthiasis: number of people treated in 2016. [Online] Available at: [https://www.who.int/neglected_diseases/resources/who_wer9249/en/#:~:text=In%202016%2C%20%3E89.2%20million%20individuals,million%20women%20of%20reproductive%20age\).](https://www.who.int/neglected_diseases/resources/who_wer9249/en/#:~:text=In%202016%2C%20%3E89.2%20million%20individuals,million%20women%20of%20reproductive%20age).) (Accessed 12th May 2021)

World Health Organization (2017) 'Preventive Chemotherapy and Transmission Control Databank.' [Online]. Available at: http://www.who.int/neglected_diseases/preventive_chemotherapy/databank/en/index.html; (Accessed 13th May 2021)

World Health Organization (2016). 'Coverage Evaluation Surveys for Preventive Chemotherapy: Field guide for Implementation' [Online]. Available at: <https://www.ntdsupport.org/sites/default/files/uploads/docs/resources/Coverage%20Evaluation>

UNDER PEER REVIEW