

## Original Research Article

### PROFILE OF ANXIETY AND DEPRESSION SYMPTOMS ASSOCIATED WITH SLEEP QUALITY OF MEDICAL STUDENTS IN THE EASTERN AMAZON

#### ABSTRACT

**Objective:** Associate the prevalence of the main symptoms of anxiety and depression with sleep quality in medical students at a Brazilian university center.

**Methods:** This is a cross-sectional, descriptive, and analytical study, with data obtained through online questionnaires answered by medical students, during the year 2022. To assess symptoms of depression and anxiety, the Hospital Anxiety and Depression Scale (HADS) was used, while, to assess the quality of sleep, the Pittsburgh Sleep Quality Index (PSQI).

**Results:** 229 students participated in the study. Of these, 51% had anxiety symptoms. As for depression, 27.5% had depression symptoms. With regard to sleep quality, 28% had poor sleep quality, and 71% had good quality. There was a significant association between sleep quality and anxiety and depression symptoms ( $p < 0.001$ ): of the individuals with good sleep quality, 51% had possible and probably anxiety and 21.2% were classified as possible and probably depression. Furthermore, among the students with poor sleep quality: 80% were possibly and probably anxious students and 43.8% were possibly and probably depressive.

**Conclusions:** The study documents a statistically significant association between anxiety and depression symptoms and poor sleep quality. Further multicentric analyzes are needed to understand this scenario.

**Keywords:** anxiety, depression, sleep, students, medical.

## INTRODUCTION

Medical students have always been a target of several studies that evaluate mental health, due to the great amount of content to which they are exposed and, consequently, to greater pressure and time dedicated to studies<sup>1</sup>. Thus, such students have a higher prevalence of mental disorders, such as depression and anxiety, when compared to university students from other areas. Their quality of life was also considered worse than that of individuals in the same age group<sup>2</sup>.

Furthermore, sleep is one of the most relevant topics nowadays, as there is consistent evidence that deprivation influences metabolic and inflammatory processes with significant negative impacts on human health<sup>3,4</sup>, and may also act as an early marker of anxiety and depression<sup>5,6</sup>. In general, medical students present susceptibility to unsatisfactory sleep, reducing their sleep time to support their academic obligations, corroborating that poor sleep has a higher prevalence in them compared to the general population<sup>6,7</sup>.

Considering that research indicates that sleep disorders tend to occur simultaneously with the increase in mental health disorders in university students, little is known about the relationship between sleep quality and mental health specifically in medical students<sup>8</sup>.

Therefore, given the need to understand this scenario, the present study aimed to establish the prevalence and relationship between anxiety and depression symptoms and sleep quality in medical students at a university center in northern Brazil.

## METHODS

### Study Design

This work is a cross-sectional, analytical, and observational study to obtain data through the questionnaires answered by students enrolled in the Medicine Course of the University Center of the State of Pará-Brazil, in the year 2022. From this, it was analyzed data on the prevalence of anxiety and depression symptoms in medical students and compared ~~them~~ with sleep quality scores, seeking to identify possible relationships between these conditions.

Data collection through questionnaires was authorized by the center, through approval by the Institution's Research Ethics Committee. In addition, data collection was authorized by the students, by signing the Informed Consent Form contained on the first page of the questionnaire.

### Sample Selection

This analysis included all medical students from the university center, from the 1st to the 12th semester, who agreed to answer the questionnaires and who signed the Informed Consent Form. Exclusion criteria were not adopted.

In the second section of the questionnaire, there were sociodemographic questions, such as age, gender, and stage in the medical school. The students were considered per biennium, as the medical education in Brazil is divided into stages, with the first two years considered the "basic cycle", years 3 and 4 the "clinical cycle", and years 5 and 6 "internship", usually composed exclusively of internships and practices.

### Data Collection and Analysis

To subjectively assess sleep quality, an excellent recognized tool is the "Pittsburgh Sleep Quality Index (PSQI)" of 1989<sup>9</sup>. The questionnaire evaluates seven sleep components: subjective quality of sleep, sleep latency, sleep duration, sleep efficiency, sleep disorders, medication use, and daytime dysfunction. The score can range from 0 to 3 for each component, with a maximum total score of 21 points. Scores below 5 indicate good quality of sleep and above that indicate poor quality of sleep. This index was validated for the Portuguese language<sup>10</sup> and is considered accessible to identify sleep quality, identifying whether individuals sleep well, poorly, or have sleep disorders<sup>11</sup>. The PSQI showed strong reliability, with moderate structural validity in different samples, suggesting that the tool fulfills its intended utility<sup>12</sup>.

To assess symptoms of anxiety and depression, the Hospital Anxiety and Depression Scale (HADS)<sup>13</sup> validated for the Portuguese language<sup>14</sup> was used. The scale evaluates subjective symptoms of anxiety and depression and does not confirm these diseases or even replace the evaluation of a specialist professional. The HADS consists of 14 items divided into two subscales, 7 questions to assess depression and 7 to assess anxiety. Each item is scored from 0 to 3, with a maximum score of 21 for anxiety or depression. The recommended scores according to Zigmond and Snaith<sup>13</sup> are 8 to 10 for possible cases and  $\geq 11$  for probable cases. Scores less than 7 indicate non-cases.

The choice to use the HADS scale was due to its easy completion, which allowed for quick and reliable data collection, and the great advantage of the absence of confusing symptoms common between clinical illnesses and depression or anxiety, such as fatigue, loss of appetite, and sleep disturbances<sup>15</sup>. Furthermore, the scale proved to be potentially effective and applicable for the screening and detection of mild anxiety and depression disorders<sup>16,24,25</sup>.

Data were collected through online questionnaires after Institution's Research Ethics Committee approval, from June to August 2022. They were organized in Microsoft Excel 2010. Graphs and tables were built with the tools available in Microsoft Word, Excel, and Bioestat 5.5 programs.

All tests were performed using the Bioestat 5.5 software. Quantitative variables were described by frequency and percentage. 95% confidence intervals were calculated for the proportion to infer how the prevalences behave in relation to the population from which they were obtained. The independence or association between two categorical variables was tested using the chi-square or Fisher's exact test, depending on the case, and significant associations were detailed using the analysis of standardized residues, to identify the categories that most contributed to the result. Results with  $p \leq 0.05$  were considered statistically significant.

## RESULTS

A total of 229 students were included in the study, and 7 students refused to participate. Most of whom them (142 or 62%) were female and 37.6% were male. Most (142 or 62%) were aged between 21 and 25 years and 25.3% of individuals were aged between 17 and 20 years (Table 1).

Considering the students per biennium, 37.6% of the individuals were from the 9th to the 12th semester, 75 individuals (32.8%) were from the 5th to the 8th semester and 86 (37.6%) were from the 9th to the 12th semester (Table 1).

**Table 14-** Sociodemographic characteristics of the students, evaluated from June to August 2022, Belém-Pará.

Variable	Frequency	Percentage
<b>Sex</b>		
Female	142	62.0
Masculine	86	37.6
Other	1	0.4
<b>Age</b>		
From 17 to 20 years old	58	25.3
From 21 to 25 years old	142	62.0
From 26 to 30 years old	20	8.7
<b>Biennium in the medical school</b>		
1st to 2nd year	68	29.7
3rd to 4th year	75	32.8
5th to 6th year	86	37.6

Percentages are relative to the total number of participants (n=229).

As for the classification of anxiety and depression symptoms, 94 individuals (41%) had an improbable classification of anxiety and 37.6% of individuals had possible anxiety. Most (166 or 72.5%) had unlikely depression, 49 subjects (21.4%) had possible depression, and 6.1% (14 subjects) had probable depression. 64 (27.9%) had poor quality sleep. The narrower this interval is, the

Formatted: Font: Bold, Do not check spelling or grammar

greater the certainty related to that prevalence in the population from which this sample was obtained (Table 2).

**Table 2-** Classification regarding anxiety and depression (HADS) and sleep quality (PSQI) of the students, evaluated from June to August 2022, Belém-Pará.

Variable	Frequency	Percentage	CI95%
<b>Anxiety Classification (HADS)</b>			
Unlikely	94	41.0	34.7 - 47.7
Possible	86	37.6	31.3 - 44.2
Likely	49	21.4	16.4 - 27.4
<b>Depression Classification (HADS)</b>			
Unlikely	166	72.5	66.1 - 78.1
Possible	49	21.4	16.4 - 27.4
Likely	14	6.1	3.5 - 10.3
<b>Sleep Quality (PSQI)</b>			
Good sleep	165	72.1	65.7 - 77.7
bad sleep	64	27.9	22.3 - 34.3

Percentages are relative to the total number of participants (n=229). 95% CI: 95% confidence interval for the prevalence.

Table 3 showed the association between sleep quality (PSQI), age, gender and stage in the medical school. In individuals with good sleep quality, most (104 or 63%) were aged 21 to 25 years and 26.1% of individuals were aged 17 to 20 years. In the group with poor sleep quality, more than half (38 or 59.4%) were aged 21 to 25 years and 23.4% of subjects were aged 17 to 20 years. However, these observed differences did not reach statistical significance ( $p=0.353$ ). Sleep quality and sex were significantly associated ( $p=0.007$ ): of the 64 individuals with poor sleep quality, 78.1% were female, in a higher proportion than expected; of the individuals with good sleep quality, 43.6% were male, a higher proportion than expected by the statistical test. That is, the quality of sleep among women was significantly worse. Furthermore, according to the data obtained, sleep quality and stage in the medical school were not significantly associated ( $p=0.064$ ).

**Table 3-** Association between age, gender and stage in the medical school with sleep quality of the students, evaluated from June to August 2022, Belém-Pará.

Variable	Good Sleep (n=165)	Bad Sleep (n=64)	p-value
<b>Age</b>			0.353
From 17 to 20 years old	43 (26.1)	15 (23.4)	
From 21 to 25 years old	104 (63.0)	38 (59.4)	
From 26 to 30 years old	11 (6.7)	9 (14.1)	

From 31 to 39 years old	7 (4.2)	2 (3.1)	
<b>Sex</b>			0.007
Female	92 (55.8)*	50 (78.1)†	
Masculine	72 (43.6)†	14 (21.9)*	
<b>Biennium in the medical school</b>			0,064
1st to 2nd year	56 (33,9)	12 (18,8)	
3rd to 4th year	49 (29,7)	26 (40,6)	
5th to 6th year	60 (36,4)	26 (40,6)	

Categorical variables are displayed as n (%). The percentages are relative to the total of each column. In all cases, the chi-square was used. \*: this frequency was lower than what would be expected by chance. †: this frequency was higher than expected.

Table 4 showed the association between sleep quality, anxiety classification, and depression classification. There was a significant association between sleep quality and anxiety ( $p < 0.001$ ): of the individuals with good sleep quality, 81 (49.1%) had improbable anxiety, this proportion being higher than expected by the statistical test; of the 64 individuals with poor sleep quality, 24 (37.5%) were likely, this proportion being higher than expected by the statistical test. There was also a significant association between sleep quality and depression ( $p < 0.001$ ): of the individuals with good sleep quality, 130 (78.8%) were classified as unlikely depression, in a higher proportion than expected; in terms of poor sleep quality, 9 (14.1%) were likely to have depression, with this proportion being higher (†) than expected.

**Table 4-** Association between anxiety classification (HADS), depression classification (HADS) and sleep quality of the students, evaluated from June to August 2022, Belém-Pará.

Variable	Good Sleep (n=165)	Bad Sleep (n=64)	p-value
<b>Anxiety Classification (HADS)</b>			<0.001
Unlikely	81 (49.1)†	13 (20.3)*	
Possible	59 (35.8)	27 (42.2)	
Likely	25 (15.2)*	24 (37.5)†	
<b>Depression Classification (HADS)</b>			<0.001
Unlikely	130 (78.8)†	36 (56.3)*	
Possible	30 (18.2)	19 (29.7)	
Likely	5 (3.0)*	9 (14.1)†	

Categorical variables are displayed as n (%). The percentages are relative to the total of each column. In all cases, the chi-square was used. \*: this frequency was lower than what would be expected by chance. †: this frequency was higher than expected.

## DISCUSSION

In this study, it was identified that 37% of students had possible anxiety, while 21% had probable anxiety<sup>17</sup>. As for depression, 21.4% of individuals had possible depression and 6.1% probable depression. In the same way – through the HADS - and with a similar sample, Vasconcelos<sup>18</sup> identified that 34.3% of the students presented a classification of possible anxiety, and 19.7% presented probable anxiety. With regard to depression, 19.3% had symptoms of possible depression and 5.6% manifested probable symptoms of the disorder. Furthermore, another study showed, using the HADS, that medical students had symptoms of anxiety and depression at a prevalence of 20% and 7% respectively<sup>17,19</sup>, which is compatible with the results of this study.

In recent meta-analyses, it was identified an overall prevalence rate of anxiety among medical students of 33.8% (95% confidence interval: 29.2-38.7%)<sup>20</sup> and a global prevalence of depression of 28.0% (confidence interval of 95% [CI] 24.2-32.1%)<sup>21</sup>. Thus, although the sample of the present study is small, of a unicentric nature, and there are different screening tools for these conditions, it was possible to identify similarity with the values obtained in relation to the classification of possible anxiety and depression, guaranteeing reliability in the screening of the symptoms of these conditions through HADS.

When considering the extensive workload demanded in the medical course, many students find the availability to study at night, thus reducing the number of hours reserved for sleep<sup>6,22</sup>. In addition, the impairment of the quality of this sleep due to the stressors of medical graduation, such as study overload, high competitiveness, responsibility, and expectations associated with the physician's role, is analyzed<sup>6,22-24</sup>. Such contexts become alarming and are connected, since sleep disorders increase the incidence of psychiatric disorders, in addition to reducing the professional and academic performance of individuals<sup>24</sup>.

Thus, anxiety symptoms are associated with more sleep disorders and the use of sleeping pills<sup>25-29</sup>, with proven polysomnographic changes in these individuals<sup>26</sup>. In the present study, there was a significant association between anxiety symptoms and poor sleep quality ( $p < 0.001$ ), in which 37.5% were probable and 42.2% possible anxious individuals with poor sleep quality. With that in mind, adding these values, about 80% of medical students with anxiety symptoms had poor sleep quality.

In other studies<sup>27-29</sup>, poor sleep quality was associated with anxiety and depression. In the present study, which also showed a relevant association ( $p < 0.001$ ), 29.7% had probable depression and 14.1% possible, totaling 43.8% of the total number of students with depressive symptoms and poor sleep quality.

In this sense, understanding that insomnia can behave as a predictor of anxiety<sup>30</sup> and depression<sup>30-32</sup> and that it can also be a symptom of both disorders - according to the DSM-5<sup>33</sup> - it becomes difficult to establish a direct causal relationship. However, it is proven that cognitive-behavioral therapy for sleep

disorders improves mental health and the severity of symptoms in patients with mental disorders, including depression<sup>34</sup>; making sleep treatment an important front not only for the prevention but also for the treatment of these disorders.

Although the study did not identify a significant association between sleep quality and stage in the course or age, sleep quality and gender were significantly associated ( $p=0.007$ ): of the 64 individuals with poor sleep quality, 78.1% were from ~~the~~ women. Other studies point to this same factor, in which they confirm that sleep disturbances, mainly insomnia, are more prevalent among women due to behavioral aspects and biological constitution<sup>27,35-37</sup>.

Other markers of mental suffering, such as burnout syndrome, also deserve attention in the medical field. This syndrome occurs not only in students, but also in medical residents, who perpetuate these disorders. A recent Brazilian study with resident physicians exposed ~~the~~ relevant prevalence of anxiety, depression, and burnout, identifying similar patterns and individual and environmental common factors, such as high workloads, ~~the~~ occurrence of psychological abuse, poor supervision of the faculty, unsatisfactory learning experience and conflicts between work and personal life<sup>38,39,40</sup>.

As for the limitations of the study, the use of scales is listed, in which both the HADS - although it suggests, does not confirm anxiety or depression - and the PSQI are subjective tools and can be influenced by memory, inattention or even dishonesty of the participants. In addition, the different scales and other ways of evaluating these disorders used in the studies end up making a completely reliable comparative analysis difficult. It is also possible to point out the failure to establish causality and other confounders associated with sleep quality and/or symptoms of anxiety and depression. Finally, the unicentric character of the study may limit the interpretation of the results<sup>38,39,40</sup>.

## CONCLUSION

In summary, the data from this study contemplate what was outlined as an objective, significantly supporting the harmful relationship between mental health disorders, depression and anxiety, and poor sleep quality. With regard to sociodemographic characteristics, the study also showed a statistically significant association between poor sleep quality and ~~the~~ female gender.

The data obtained are relevant for the definition of intervention models, applying them after recognizing the variables that interfere ~~in-with~~ the connection between mental health disorders and, consequently, bad sleep quality. Further multicenter analyzes are needed to understand this scenario in medical schools around the world.

## REFERENCES

1. Oliveira LM. et al. Medical student's quality of life. Goiania. Goias Federal University. 2015.
2. Barbosa-Medeiros MR, Caldeira AP. Mental health of medical students: longitudinal study. *Rev. bras. educare med.* 2021, v. 45(3). DOI: 10.1590/1981-5271v45.3-20190285.
3. Barros MBA, Lima MG, Ceolim MF, Zancanella, E, Cardoso TAMO. Quality of sleep, health and well-being in a population-based study. *Rev. of Public Health.* 2019.v. 53, 82. DOI:10.11606/s1518-8787.2019053001067.
4. Cappuccio FP. et al. Sleep Duration and All-Cause Mortality: A Systematic Review and Meta-Analysis of Prospective Studies. *Sleep.* 2010. v. 33(1), 585-592.
5. Almojali AI. et al. Prevalence and association of stress with sleep quality in medical students. *J for Epidemiol. and Global Health.* 2017. v. 7, no. 3, 169-174.
6. Azad MC, Fraser K, Rumana N, Abdullah AF, Shahana N, Hanly PJ, Turin TC. Sleep disturbances among medical students: a global perspective. *J Clin Sleep Med.* 2015 Jan 15;11(1):69-74. doi: 10.5664/jcsm.4370. PMID: 25515274; PMCID: PMC4265662.
7. From LD Light. et al. Assessment of the Pittsburgh Sleep Quality Index in medical students: An integrative literature review. *Research Soc. and Dev.* 2021. v. 10, no. 8, E52410817530.
8. Milojevich HM, Lukowski AF. Sleep and Mental Health in Undergraduate Students with Generally Healthy Sleep Habits. *PLoS One.* 2016 Jun 9;11(6):e0156372. doi: 10.1371/journal.pone.0156372.
9. Bugueño M. et al. Sleep quality and academic performance in high school students. *Rev. Med. from Chile.* 2017. v. 145, no. 9, 1106-1114.
10. Bertolazi NA. Translation, cultural adaptation and validation of two sleep assessment instruments: EPWORTH Sleepiness Scale and Pittsburgh Sleep Quality Index. Porto Alegre. Federal University of Rio Grande do Sul. 2008.
11. Escobar-Córdoba F, Eslava-Schmalbach J. Colombian Validation of the Pittsburgh Sleep Quality Index. *Rev de Neurol.* 2005. 40(3), 150-155.
12. Mollayeva T, Thurairajah P, Burton K, Mollayeva S, Shapiro CM, Colantonio A. The Pittsburgh sleep quality index as a screening tool for sleep dysfunction in clinical and non-clinical samples: A systematic review and meta-analysis. *Sleep Med Rev.* 2016, 25. 52-73. DOI: 10.1016/j.smrv.2015.01.009
13. Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand.* 1983. 67(6). 361-370. DOI:10.1111/j.1600-0447.1983.tb09716.x
14. Apostolo JLA, Mendes AC, Azeredo ZA. Adaptation to Portuguese of the Depression, Anxiety and Stress Scales (DASS). *Rev Lat Am Nursing.* 2006.
15. Castro MMC, Quarantini L, Batista-Neves S, Kraychete DC, Daltro C, Miranda-Scippa Â. Validity of the hospital anxiety and depression scale in patients with chronic pain. *Rev Bras Anesthesiol.* 2006. 56(5). 470-477. DOI: 10.1590/S0034-70942006000500005

16. Botega NJ, Bio MR, Zomignani MA, Garcia Jr C, Pereira WAB. Mood disorders in a medical clinic ward and validation of an anxiety and depression measurement scale (HAD). *Rev Saude Publica*. 1995. 29(5). 359-363. DOI: 10.1590/S0034-89101995000500004
17. Faro A. Confirmatory Factor Analysis and Normalization of the Hospital Anxiety and Depression Scale (HADS). *Psicol Teor e Pesqui*. 2015. 31(3). 349-353. DOI:10.1590/0102-37722015032072349353
18. Vasconcelos TC. et al. Prevalence of anxiety and depression symptoms in medical students. *Rev Bras de Educ Med*. 2015.v. 39, p. 135-142.
19. Bertani DE. et al. Anxiety, depression and personality traits in Italian medical students. *Rev Psiquiatr*. 2020. 55(6). 342-348. DOI: 10.1708/3503.34892.
20. Tian-Ci QT, Tam WSX, Tran B, Zhang M, Zhang Z, Su-Hui HC, Chun-Man HR. The global prevalence of anxiety among medical students: a meta-analysis. *Int J of Envir Res and Pub Health*. 2019.v. 16 (15) 2735.
21. Puthran R. et al. Prevalence of depression amongst medical students: A meta-analysis. *Med education* 2016. v. 50, (4). 456-468.
22. Almojali AI, Almaliki SA, Alothman AS, Masuadi EM, Alaqeel MK. The prevalence and association of stress with sleep quality among medical students. *J Epidemiol Glob Health*. 2017 Sep;7(3):169-174. doi: 10.1016/j.jegh.2017.04.005. Epub 2017 May 5.
23. Thomas M, Bigatti S. Perfectionism, impostor phenomenon, and mental health in medicine: a literature review. *Int J Med Educ*. 2020 Sep 28;11:201-213. doi: 10.5116/ijme.5f54.c8f8.
24. Ludwig AB, Burton W, Weingarten J, Milan F, Myers DC, Kligler B. Depression and stress amongst undergraduate medical students. *BMC MedEduc*. 2015 Aug 27;15:141. doi: 10.1186/s12909-015-0425-z.
25. Becker SP. et al. Sleep in a large, multi-university sample of college students: sleep problem prevalence, sex differences, and mental health correlates. *Sleep health*. 2018. v. 4(2). 174-181.
26. Baglioni C, Nanovska S, Regen W, et al. Sleep and mental disorders: A meta-analysis of polysomnographic research. *Psychol Bull*. 2016. 142(9). 969-990. DOI:10.1037/bul0000053.
27. Vilchez-Cornejo J. et al. Mental health and sleep quality en estudiantes de ocho facultades de medicina humana del Perú. *Rev. chil. neuro-psiquiatr*. 2016. v. 54(4). 272-281. DOI:http://dx.doi.org/10.4067/S0717-92272016000400002.
28. Al-Khani AM. et al. A cross-sectional survey on sleep quality, mental health, and academic performance among medical students in Saudi Arabia. *BMC research notes*. 2019. v. 12(1). 1-5.
29. Joo HJ, Kwon KA, Shin J, Park S, Jang SI. Association between sleep quality and depressive symptoms. *J Affect Disord*. 2022. 3(10). 258-265. DOI:10.1016/j.jad.2022.05.004
30. Hertenstein E, Feige B, Gmeiner T, et al. Insomnia as a predictor of mental disorders: A systematic review and meta-analysis. *Sleep Med Rev*. 2019. 4(3). 96-105. DOI:10.1016/j.smr.2018.10.006

31. 31. Li L, Wu C, Gan Y, Qu X, Lu Z. Insomnia and the risk of depression: a meta-analysis of prospective cohort studies. *BMC Psychiatry*. 2016. 16(1). 375. DOI:10.1186/s12888-016-1075-3
32. 32. Baglioni C, Battagliese G, Feige B, et al. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord*. 2011. 135(3).10-19. DOI:10.1016/j.jad.2011.01.011
33. 33. Nascimento MIC. et al. Manual Diagnóstico e Estatístico de Transtornos Mentais: DSM-5. 5ª ed. Porto Alegre. Artmed, 2014. Accessed on: 10 Jan 2023. Available at: <http://www.institutopebioetica.com.br/documentos/manual-diagnostico-e-estatistico-de-transtornos-mentais-dsm-5.pdf>.
34. 34. Hertenstein E, Trinca E, Wunderlin M, et al. Cognitive behavioral therapy for insomnia in patients with mental disorders and comorbid insomnia: A systematic review and meta-analysis. *Sleep Med Rev*. 2022. 62.101-597. DOI:10.1016/j.smrv.2022.101597
35. 35. Hur S, Oh B, Kim H, Kwon O. Associations of Diet Quality and Sleep Quality with Obesity. *Nutrients*. 2021. 13(9). 3181. DOI:10.3390/nu13093181
36. 36. Mehta N, Shafi F, Bhat A. Unique Aspects of Sleep in Women. *Mo Med*. 2015. 112(6). 430-434.
37. 37. Krishnan V, Collop NA. Gender differences in sleep disorders. *Curr Opin Pulm Med*. 2006.12(6). 383-389. DOI:10.1097/01.mcp.0000245705.69440.6a
38. 38. Gil-Calderón J, Alonso-Molero J, Dierssen-Sotos T, Gómez-Acebo I, Llorca J. Burnout syndrome in Spanish medical students. *BMC Med Educ*. 2021 Apr 22;21(1):231. doi: 10.1186/s12909-021-02661-4.
39. 39. Bhugra D, Molodynski A, Ventriglio A. Well-being and burnout in medical students. *Ind Psychiatry J*. 2021 Jul-Dec;30(2):193-197. doi: 10.4103/ipj.ipj\_224\_21. Epub 2021 Nov 23.
40. 40. De Mélo SJML, Valença MM, Rocha-Filho PAS. Individual and residency program factors related to depression, anxiety and burnout in physician residents - a Brazilian survey. *BMC Psychiatry*. 2022 Apr 19;22(1):272. doi: 10.1186/s12888-022-03916-0.

Comment [null1]: Double numbering