

Impacts of pollutants (phosphorus, nitrogen and potassium) from agricultural activities on the soils and waters of Toho Lake (Benin)

ABSTRACT

We evaluated the impacts of agricultural activities on the cultivated soils around Lake Toho as well as the waters from the streams through phosphorus, nitrogen and potassium pollution of the soils and waters from the streams heading towards Lake. For soil, pH_{water} and pH_{KCl} organic matter, moisture total phosphorus and its fractionation were determined. The various supernatants obtained are analyzed using a 1600PC UV spectrophotometer as well as the waters for the determination of ammonium, nitrite, nitrates and phosphate. In soils, the determination of ammonium, nitrite, nitrate, phosphate and potassium was carried out by the DR 5000 spectrometry method of the MP-AES brand. As a result, the soils around this lake are weakly acidic with an average water pH of 6.91. Humidity increases from surface to depth (8% to 28%), while organic matter (11% on average) and organic carbon decrease from surface to depth. These soils are rich in phosphorus with an overall average concentration of 10.10 mg/g. The following forms of phosphorus are present in soil in the order $P\text{-residual} > P\text{-org\&Al} > P\text{-Ca} > P\text{-Fe} > P\text{-Labile}$. The physico-chemical analyzes of the waters show that these waters are moderately loaded and present an increased risk of eutrophication with multifaceted consequences. In soils, nitrate is the dominant form of nitrogen. Cultivated soils provide an abundant source of nitrogen and phosphorus nutrients to Lake Toho via its recharge sources.

Keywords: Lake Toho, phosphorus, nitrogen, potassium, stream waters, fractionation.

1. INTRODUCTION

Since its existence and for its survival, man has always used the environment for its various needs. And since that time man has used the resources of this environment for his provision and development. With the demographic explosion, these needs have increased exponentially, leading man to overexploit the environment; especially for agriculture, construction, energy and transport by means of the industrial revolution. All these areas have led to an imbalance of the environment through the destruction of plant cover, atmospheric emissions, the use of chemicals leading to pollution of soil and water resources. In 1840, [1] showed that phosphorus inputs were necessary to increase crop yields. This has led to agricultural automation through the production of chemical fertilizers. The nitrogen and phosphorus contained in these chemical fertilizers are at the root of environmental problems, such as eutrophication and the toxicity of surface and ground water [2]. Modern agriculture nowadays requires the use of chemical fertilizers to increase production yields. Thus, we are witnessing an uncontrolled use of these inputs which are nowadays the main source of pollution of soils and water resources. This phenomenon intensifies with the non-respect of the technical itineraries of agricultural production and the uncontrolled production of chemical fertilizers.

The mobility of phosphorus, nitrogen and potassium in soils and their transfer to surface waters depends on the different forms available. The presence of phosphorus in surface waters causes eutrophication of aquatic environments and that of nitrite disrupts the quality and functions of water in a damaging way [3]. This contamination leads to the loss of aquatic

species. Thus, Lake Toho located in southern Benin has experienced the death of fish several times in 2012, 2018, and in 2021. In 2012, according to information from PNE-Benin, the death of fish occurred following a stormy rain which changed the turbidity of the water. This phenomenon is a source of deoxygenation of the aqueous medium and has cost the lives of these fish. In 2018 and 2021, the same drama happened on the same lake. For this time if the main cause of this phenomenon in 2018 and 2021 is linked to the spill of harmful substances from unknown sources with change of color PNE Benin. Studies have shown that the lake is subject to anthropogenic pollution [4,5]. Since agriculture is the main activity that develops around the lake, it is therefore essential to study the mobility and bioavailability of phosphorus, nitrogen and potassium, which constitute the essential elements of the chemical fertilizers used in the area.

2. MATERIAL AND METHODS

2.1 Study area

The study was carried out around Lake Toho which is located between 6°35 to 6°40 north latitude and 1°45 to 1°50. It covers an area of 9.6 km² in the dry season and 15 km² in the rainy season [6]. Lake Toho is surrounded by several cultivable soils where samples have been taken.

2.2 Sampling

A total of thirty (36) samples were taken, including 9 water samples on three sites and 27 soil samples on nine sites along the three horizons [0, 30]; [30, 60]; [60, 90] cm




Figure 1: Sampling map

Table 1: Characteristics of sampling stations

SITES	Geographical coordinates	Characteristic	Samples
-------	--------------------------	----------------	---------

S1 (HOUNKPOT ONNOU)	6°35'53.35"N 1°47'11.84"E	-Next to a usable borehole -Surrounded by cornfields - Presence of public toilets nearby - Presence of an old abandoned borehole	Soil
S2 (TOHONOU DADAHOUE)	6°36'34.66"N 1°47'33.33"E	-Fields of corn, okra, teak and banana trees. -Use of herbicides on site soil -Proximity to houses not far from the lake	Soil
S3 (TOHONOU FIFADJI)	6°37'7.40"N 1°47'22.10"E	-activities carried out fishing, agriculture, - Laundry	Water Soil Sediments
S4 (KPINNOU)	6°35'5.12"N 1°45'59.17"E	-Houses not far from the lake - fields of crincrin not far from the lake - fishing on the lake	Water Soil Sediments
S5 (AGONTIME)	6°36'16.35"N 1°45'6.06"E	-Large fields of horsehair - use of herbicides -Presence of stream	Water Soil Sediments
S6 (KPONNOU)	6°36'36.49"N 1°45'25.13"E	-Immediate proximity to the lake -laundry and fishing on the lake -Cornfield -corn fields by the lake	Soil
S7 (LOGBO)	6°37'0.65"N 1°46'0.42"E	-Laundry by the lake -sin - discharge of fish waste into the lake	Soil
S8 (HOUIN AGADJI)	6°38'9.14"N 1°45'53.54"E	- large corn fields -palmerais	Soil
S9 (DUIME)	6°38'23.33"N 1°46'54.10"E	- corn fields not far from the lake Presence of fish farm	Soil

2.3 Analysis methods

The soil samples were dried in an oven at 40°C for 24 hours without inducing any significant change in their characteristics. The pH_{water} and the pH_{KCl} were measured with the HANNA brand multi-parameter conductivity meter according to the NF X 31-103 standard. The humidity was determined according to the Standard AFNOR X31-102, [7]. As for the organic matter, it was determined according to the Walkley-Black method [8]. The determination of total phosphorus was carried out by mineralization with potassium persulfate in an acid medium (H₂SO₄) at 120° C for 2 hours [9]. As for the fractionation, it was carried out according to the protocol of Rydin and Welch [10]. The extractions were analyzed by a UV-1600 PC spectrophotometer. As phosphate and potassium in the soil were made using

the DR 5000 MP-AES spectrometry method. The determinations of nitrite, nitrate, ammonium, orthophosphates and then potassium were carried out by the colorimetric method using a UV-1600 PC spectrophotometer. The Ifremer eutrophication risk diagnostic grid [11] is used to assess the eutrophication risks of the soils and water analyzed.

3 RESULTS

o Physicochemical characterization of soils

The results of the physicochemical parameters of the soils showed that the water pH value varies from one site to another and from one horizon to another and is between 5.23 and 7.89 with an average of 6.91. These soils are moderately acidic. (Figure 2).

The recorded pH_{KCl} value varies from 4.2 to 7.09 with an average of 5.73 (Figure 3). At the level of the sites (S1, S2, S4, S7, S8 and S9) the value decreases starting from the surface towards the depth while at the level of the other sites, it increases.

The water content (moisture) of the soil around Lake Toho decreases from surface to depth at sites (S2, S4, S5, S7 and S9), increases at sites (S3, S6 and S8) and remains constant at the site (S1) Figure 4.

The organic matter content in the soils varies between 3% and 11%, the highest content is recorded at site S2 on the surface (horizon 30). The lowest grade is observed at horizon 90 at the sites (S2, S3, S5 and S7). The upper layer shows the high organic matter content at the sites (S2, S3, S5 and S7) compared to the other sites (Figure 5).




Figure 2: Variations in soil of pH_{water}




Figure 3: Variations in soil of pH_{KCl}




Figure 4: Soil humidity content




Figure 5: Soil organic matter content

○ **Phosphorus and its derivatives in soils**

According to Figure 6, the fraction of total phosphorus predominates at horizon H30 on all the sites, the concentration of total phosphorus increases going from the surface towards the depth at the level of site S3, it decreases going from the surface towards the depth at sites (S1, S4, S6, S7 and S8), with the exception of sites (S2, S5 and S9) where the opposite is observed. This phosphorus present in the different layers of the soil is bound to several particles or elements of the soil in several forms.

• **The variation of each form of phosphorus according to soil profiles**

According to Figure 7, the fraction of labile phosphorus predominates at horizon H30 on all sites, the concentration of phosphorus decreases from the surface to the depth at the levels of (S1, S6, S7 and S8) with the exception of sites (S2, S3, S4, S5 and S9) where the opposite is observed.

According to Figure 8, the fraction of phosphorus bound to iron (P-Fe) predominates at the H30 horizon on all the sites, the concentration of phosphorus bound to iron decreases going from the surface towards the depth at the level sites (S1 and S8) with the exception of sites (S2, S3, S4, S5, S6, S7 and S9) where the opposite is observed.

According to figure 9, the fraction of phosphorus bound to organic matter predominates at the H90 horizon on all the sites, the concentration of phosphorus bound to organic matter increases going from the surface towards the depth at the level of the sites (S2 and S5), it decreases from the surface to the depth at site S9 with the exception of sites (S1, S3, S4, S6, S7 and S8) where the opposite is observed.

According to figure 10, the fraction of phosphorus bound to calcium predominates at the H30 horizon on all the sites, the concentration of phosphorus bound to calcium decreases going from the surface to the depth at the level of the sites (S1 and S7), it increases going from the surface to the depth at the level of the S9 site, with the exception of the sites (S2, S3, S4, S5, S6 and S8) where the opposite is observed.

According to figure 11, the fraction of residual phosphorus predominates at the H30 horizon on all the sites, the concentration of residual phosphorus increases going from the surface towards the depth at the level of the sites (S5 and S7), it decreases from the surface to the depth at site S8, with the exception of sites (S1, S2, S3, S4, S6 and S9) where the opposite is observed.




Figure 6: Content of the total fraction




Figure 7: Content of the labile fraction




Figure 8: Content of iron-bound fraction




Figure 9: Content of the fraction bound to organic matter




Figure 10: Content of the fraction bound to calcium

Figure 11: Content of the residual fraction

- The predominance of different forms by Horizon

According to figure 12 at horizon [0 - 30]. The labile fraction is low on all the sites except site S7. The residual fraction (P-residual) predominates over the others at the H30 horizon at the site levels (S1 and S2). On the other hand, at the level of the sites (S3, S4, S8 and S9) the organic fraction (P-Org) predominates over the other fractions, at the level of the S6 site the fraction bound to iron predominates, and at the site S5 the fraction bound to calcium predominates. Of all the fractions, the organic fraction predominates at horizon H30.

According to figure 13 at horizon [30-60] the labile fraction (P-L) is low on all the sites. The residual fraction (P-r) predominates at the sites (S1, S4, S5, and S6). On the sites (S7, S8 and S9) it is the organic fraction which predominates while on the sites (S2 and S3) it is the fraction linked to calcium which predominates. At horizon H60 after the residual fraction, the organic fraction takes the second position.

According to Figure 14 at horizon [60-90], labile phosphorus (P-L) is low at all sites. Residual phosphorus (P-r) predominates at site levels (S1, S2 and S7). At the level of the sites (S3, S4, S5, S6, S8, and S9) it is the organic fraction (P-org) which predominates.




Figure 12: the cumulative value (a) and proportions (b) of the phosphorus fractions at H 30 cm




Figure 13: Cumulative value (a) and proportions (b) of phosphorus fractions at H 60 cm




Figure 14: the cumulative value (a) and proportions (b) of the phosphorus fractions at H 90 cm

- The different forms of nitrogen in the soil

The analysis of Figure 15 shows that the ammonium content in the soil at 30cm varies from 2.747mg/kg at site (S5) to 7.771mg/kg at site (S1), while at 60cm it varies from 2.014 mg/kg at the site (S1) to 6.255 mg/kg at the site (S4), while at 90 cm the same content varies from 3.649 mg/kg at the site (S7) to 6.823 mg/kg at site (S1). Ammonium is more concentrated on the surface horizon with the exception of site S2 and S5 where the ammonium content at horizon 90 is slightly high

According to Figure 16, the nitrite content in the soil at 30 cm ranges from 5.878 mg/kg at site level (S5) to 21.058mg/kg at site level (S1), while at 60 cm it ranges from 5.457mg/kg at site level (S1) to 15.748mg/kg at site level (S5) and at horizon 90 cm, it ranges from 6.937 mg/kg at site level (S4) to 18.489mg/ kg at site level (S1). Nitrite is more concentrated on the surface horizon except for the S2 and S5 site where the nitrite content at horizon 90 is slightly high.

According to Figure 17, the nitrate content in the soil at 30 cm varies from 17.763mg/kg at site level (S8) to 35.155mg/kg at site level (S1), while at 60 cm it is ranging from 9.110mg/kg at the site (S1) to 28.297mg/kg at the site (S4), while at 90 cm it ranges from 16.509mg/kg at the site (S7) to 30.867mg/ kg at site level (S1). The nitrate content decreases with increasing

depth, with the exception of sites (S2, S5 and S9) where the nitrite content at horizon 90 is slightly high.




Figure 15: Ammonium content in soils




Figure 16: Nitrite content in the soil




Figure 17: Nitrate content in the soil

Table 2: Rapport N/P

Sites	S1	S2	S3	S4	S5	S6	S7	S8	S9	
Depth	H30	0,016	0,001	0,007	0,007	0,004	0,002	0,004	0,006	0,010
	H60	0,002	0,005	0,007	0,007	0,005	0,018	0,007	0,005	0,018
	H90	0,021	0,001	0,015	0,007	0,001	0,007	0,006	0,006	0,018

The N/P ratio of the soil nitrogen rate to the calculated soil phosphorus rate (N/P) is much lower than 1 in all the sites, which shows that the phosphorus rate in the soils around Lake Toho predominates on the rate of nitrogen (Table 2).

○ **Assessment of the risk of mobility of soil elements on water resources**

According to the eutrophication risk assessment grid, the cultivated soils around Lake Toho are classified in the group of very poor soils with regard to the risk of eutrophication (Table 3).

Table 3: Positioning of sediment samples (S1 to S9) in the grid of Ifremer's eutrophication risk diagnostic framework (2000).

Parameters	Very good (no risk of eutrophication)	Good (low risk)	Medium (50% risk)	Bad (high risk)	Very bad (very high risk of eutrophication)
MO(%)	3.5	5	7.5	10	S1 to S9
NT (g/kg DW)	1	2	3	4	S1 to S9
PT (mg/kg DW)	400	500	600	700	S1 to S9

Water results

From Figures 18 and 19, nitrite and nitrate are present in the waters of the creek and vary in the same way. The highest value is recorded at site 5 (35.5 mg/L for nitrate and 0.57 mg/L for nitrite) and the lowest value at site 3 (0.55 mg/L for nitrate and 0.112 mg.L-1 mg/L for nitrite).

According to figure 20, we also note the presence of ammonium in the waters of the stream. Compared to nitrite and nitrate, the highest value of ammonium is observed at site S3 (0.825 mg/L) and the lowest value at site S4 (0.407 mg/L).

According to figure 21, the phosphate concentration at these sites varies from 0.063mg/L at site 3 (S3) to 0.393 mg/L at site 4 (S4).




Figure 18: Nitrate concentration in water





Figure 19: Nitrite concentration in water



Assessment of the trophic state of the waters of the stream in contact with Lake Toho
 According to Table 4, the nitrate ions found in the waters of the stream are classified in the wrong class. Similarly, phosphorus classifies 2/3 of the waters of the stream in the bad class and the nitrite ions classify them in the middle class.

Table 4: Diagnostic grid of the trophic state of stream water in contact with Lake Toho

Variables (mg.L-1)	Very good	Good	Average	Poor	Bad		
Nitrate	0	0.43	0.62	1.24	1.86	S3, S4, S5	
Nitrite	0	0.023	0.046	S3, S4	0.23	0.46	S5
Total Phosphorous	0	0.023	0.079	0.119	S5	0.316	S3, S4

Discussion

The state of a soil depends on these physicochemical and chemical characteristics. The behavior of chemical substances present in the soil depends on several factors such as pH, organic matter and inorganic matter.

The analysis of the soils around Lake Toho during dry season (May 2022) shows that the pH of the soils is 6.91 on average. In an acidic medium if the pH is less than 7 it promotes the mobility of inorganic phosphorus. On the other hand, for pH greater than 7, desorption decreases; in an alkaline medium, an increase in pH decreases the mobility of inorganic phosphorus according to Gérard [12]. The pH value of the soils around Lake Toho being close to 7, they are weakly acidic or even neutral. This pH value is within the range of the different values recorded in 2022 [13] in the same environment (6.25-8.3) in the rainy season and slightly higher than the value (6.25) in 2022 [14] around the Mekrou River. The overall average pH_{KCl} is 5.72. Obtained is close to those found by [15, 14].

The humidity of cultivated soils around Lake Toho varies from 8% to 28%, while the maximum value recorded in the rainy season is 38% [13]. This shows that the soils are wetter in the rainy season than in the dry season. As pointed out by [16], permanent exchanges of phosphates take place between the particles and the solution which bathes them. The sites S2, S4, S5 and S9 are far from the shore of the lake which explains the decrease in the humidity rate by evolving in depth, on the other hand, the sites S3, S6 and S8 are more or less close to the lake which explains the increase in humidity.

In the carbon of the soil around Lake Toho the organic part occupies 20% at most and the mineral part takes 35% at most this confirms that these soils are mineralized. Organic matter is one of the essential elements of the soil capable of forming bonds with phosphorus, its proportion in cultivated soils around the lake is 11% on average lower than that found by [17] in Lake Nokoué on the year which varies from 13.8 to 23.9% and maximum values could reach 38.22% dry weight of sediment. Authors find an average value in the sediments of the Okpara dam of around 24.62% on average [18]. Others find 16 to 28% organic matter in the sediments of the Porto-Novo lagoon [19]. We can remember then that the soils are then less rich in organic matter than the reserves towards which the waters which cross them will be stored. The organic matter rate of the soil samples is higher than the [20] value which states that the organic matter rate is normal when it is 2 to 4%. According to the same author when the soil exceeds a rate of 4% it is considered humus. So the collected samples are considered as humus.

The content of total phosphorus determined in our samples varies between 1335.5764 $\mu\text{g/g}$ at the site (S9) at 60 cm and 18644.95 $\mu\text{g/g}$ at the site (S2) at 30 Cm. These concentrations are all above the low environmental risk threshold for phosphorus (500 mg/kg of P_2O_5) established [21]. This means that the soils around Lake Toho in southern Benin are subject to phosphorus pollution. These recorded values are lower than the values recorded [13] at the end of the rainy season in the middle. This difference is due to the period and the mobility of phosphorus which can pass from the solid phase of the soil to the liquid phase of the soil and can infiltrate or run off towards watercourses and bodies of water. But they are higher than those found on the soils of the Mekrou River in northern Benin [14; 22]. We find that soil total phosphorus concentrations generally increase as one approaches Lake Toho. The greatest value is obtained at site S2 at 30 cm in a field close to the lake. This can be explained by the use of chemical fertilizers because [23], the surface application of fertilizers not buried causes an enrichment of the upper layer of the soil.

The fractionation of phosphorus in soil samples revealed the actual presence of all five phosphorus fractions in the soil around Lake Toho. The residual fraction occupies the first position in the soil on all the horizons. This result is consistent with that of 2022 in the same

environment [13] and in cultivated soils around the Mekrou River [14]. After the residual fraction, the phosphorus fraction bound to organic matter predominates. This predominance is due to the high content of organic matter in the surface horizon of the soil. The assimilable P-labile fraction represents the lowest fraction over all the layers (813.65-448.21-451.23 µg/g). On the H30 horizon, the labile fraction predominates compared to the other layers of the soil, this predominance is due to the high content of phosphorus and the humidity responsible for dissolving phosphorus into orthophosphates.

The assessment of soil quality according to the Ifremer grid shows that these soils present a risk of eutrophication. The phosphate ions (HPO_4^{2-}) found in surface waters mainly come from waste water discharges and agricultural spreading [24]. The physicochemical analyzes of agricultural wastewater heading towards Lake Toho show that the water is moderately charged and polluted by nitrogenous elements including nitrite and ammonium which are respectively between 0.112 mg/L at site 3 (S3) at 0.57 mg/L at site 5 (S5) and 0.407mg/L at site 4 (S4) at 0.825mg/L at site 3 (S3) all exceeding the limit recommended by the WFD which is 0.1 mg/L. As for nitrate, these values vary from 8.56 mg/L at the site level (S1) to 35.513mg/L at the site level (S3) and phosphate 0.063 mg/L at the site level (S1) to 0.393 mg /L at the site level (S2). The results of the trophic status diagnosis classify these waters in the wrong class. This leads to the alteration of water quality and reduces its potential for certain uses. However, it has been shown that the presence of a high concentration of nutrients (nitrogen and phosphorus in particular) is mainly responsible for the massive proliferation of phytoplankton (algae and cyanobacteria) [3]. In addition, cyanophyceae secrete toxic substances that affect both zooplankton and higher organisms. Recreational and fish-farming waters are also severely depleted [26]. The very low values of the N/P ratio confirm the table of Thomann and Mueller [27]. According to these authors for small lakes, when the N/P ratio $\ll 10$, nitrogen proves to be a limiting factor in the proliferation of aquatic plants with a call for the intensification of the activities of cyanobacteria. Because these authors summarize the usual values of the N/P ratio of several aquatic environments and show that phosphorus is the most often limiting element, especially in natural areas. When nitrogen becomes the limiting element, certain algae (cyanobacteria) compensate for this deficiency by fixing the nitrogen contained in the atmosphere. The proliferation of aquatic plants, especially macrophytes, is at the origin of the formation of a screen at the interface which constitutes an obstacle to the oxygenation of the environment. Moreover, their decomposition leads to an increase in the quantity of organic matter in the medium, the degradation of which consumes oxygen and makes the medium reduce. The production of sulfide and nitrite (toxic), through the reduction of sulfates and nitrates, inhibits the development of other primary producers and highly sensitive fish. This can cause an

imbalance in the trophic chain (production/consumption) of a body of water and lead to significant adverse ecological consequences. The ecological impact of toxin-producing cyanobacteria is little known. Hepatotoxins, which have the particularity of inhibiting certain protein phosphatases, are potentially capable of affecting a range of organisms in nature. Sometimes cyanobacteria are considered responsible for fish and bird kills. However, in many cases, the direct relationship between mortality and the presence of toxins is difficult to demonstrate. Fish death is more often associated with anoxia caused by bacteria that use compounds from the proliferation of aquatic plants. Fish can be affected by clogging of bronchitis by phytoplankton; which then reduces the efficiency of water filtration at the level of the gills.

This observation is in agreement with the results of the assessment of the risks of soil contamination which classifies these soils in the class of soils at risk of eutrophication. Despite some losses associated with water runoff, the observation is that cultivated soils are enriched in phosphorus, which leads to increased risks of transfers to aquatic ecosystems.

4. CONCLUSION

The soils cultivated around Lake Toho are subject to strong anthropogenic pressure due to the use of fertilizers in agriculture. The results show that the soils cultivated around Lake Toho are weakly acidic or even neutral with an acidic pH_{KCl} subjecting the mobility of the bioavailable forms of phosphorus. This mobility could be increased with the proliferation of cyanobacteria because the N/P ratio is much lower than 10, which draw abundant atmospheric nitrogen by consuming phosphorus. There are high levels of organic matter in the surface layer of the soil, a high content of phosphorus, nitrogen and potassium in cultivated soils. The fractionation of phosphorus in the soils taken around the lake shows that the labile fraction is the lowest. It is important to point out that the residual fraction takes precedence over the other fractions in some sites. A predominance in the organic fraction on the upper soil layer is also noted with the exception of sites located outside crop fields.

Indeed, the high content of phosphorus at the surface horizon of the soil leads to the conclusion that the contamination of the soil by phosphorus, nitrogen and potassium around Lake Toho is an anthropogenic pollution due to the use of chemical fertilizers and pesticides in the 'agriculture. These elements have appeared in the waters of streams flowing into the lake. The phosphorus accumulated in the soils during the period of over-fertilization makes these soils potential sources of continuous pollution.

REFERENCES

- 1 Liebig JV. Chemistry and its application to agriculture and physiology. Taylor and Walton, 1840 London, 4th ed, 352 pp.
- 2 Souidi R, Recovery of phosphorus from wastewater in the form of vivianite using the electrocoagulation method. LAVALE University 2021; 120p.
- 3 Pourriot and Meybeck. General Limnology, 1995 Masson Edition, 956pp
- 4 Kple Evelyne. Contribution to the study of organic and nitrogen pollution of the waters of the Toho in the department of Mono, 2008.
- 5 Gbaguidi GJ., Contribution to the protection against chemical pollution of a lentic system in the tropical region of Lake Toho in South Benin, West Africa 1018; 95p.
- 6 Topanou N, Gbaguidi GJ, Essegbemon A, Djosse GR, Mama D, Aminou T. Assessment of some heavy metals in fresh fish (*Oreochromis aureus*): Case of Toho lake in South Eastern Benin, West Africa. Journal of Geoscience and environment protection 2020 <https://www.sirp.org/journal/GEP>. Pp14.
- 7 AFNOR, 1994. Soil quality, AFNOR Edition, 250 pages.
- 8 Jackson T.F.C., Ivissi A.E., Dervalle F.B. Soil Chem. Analysis, Prentice Hall Inc. Engle Works Cliffs, New Jersey 1984, 498pp.
- 9 Gachter R., Tessier A., Szabo E. and Carignan R. Measurement of total dissolved phosphorus in small volumes of iron rich interstitial water. Aquatics science. 1992; 54p.
- 11 Ifremer, 2000. Update of indicators of the level of eutrophication of Mediterranean lagoon environments – Final report-09-2000, (Volume I – 236p) (Volume II – 196p).
- 12 Gérard F. Clay minerals, iron/aluminum oxides, and their contribution to phosphate sorption in soils_ A myth revisited. Geoderma, 2016, 262, 213-226.
- 14 Ayola PD (2022): Fractionation, bioavailability and mobility of phosphorus in agricultural soils of the Mekrou River watershed.
- 15 Amonmide I., Dagbenonbakin G., Agbangba C.E., Akonkpe P., 2019. Contribution to the assessment of the level of soil fertility in cotton-based cropping systems in Benin. Article INRAB 15p.
- 16 Furumai H. & Ohgaki S. Radiochemical analysis of phosphorus exchange kinetics between sediments and water under aerobic conditions. J. Approx. Qual. 1988 17: 205–212.
- 17 Daouda M., Martin A., Christophe K., Lyde T., Waris C., Odilon C., Véronique D., James B., and Michel B. Phosphorus fractionation of surface sediment of lake Nokoue in Benin J. Appli . Biosci. 2011 45: 3032-3044
- 18 Tometin ASL, Mama D., Zogo ND., Boukari O., Bawa LM. Eutrophication, sediment phosphorus fractionation and short-term mobility study in the surface and under profile sediment of a water dam. (Okparadam, Benin, West Africa). J.Appl. Science. Around. Manage. December 2013 Vol. 17 (4) 517-526
- 19 Chouti K. W., Atchichoe W., Tometin L., Daouda M. Bioavailability and mobility of phosphorus in sediments of the Porto-Novo lagoon. Journal of applied biosciences 2017. <https://dx.doi.org/10.4314/jab.v114i1.1>. 13p.
- 20 Dutheil C. Soil depollution by extraction under reduced pressure: study of some parameters 2003 (Doctoral dissertation, Lyon, INSA).
- 21 Dyer B. On the analytical determination of probably available mineral plant food in soils. J. Chem Soc. 1894; 65, 115-167.
- 22 Anato IF,. Influence of agricultural activities on the chemical quality of cultivated soils in the Benin sub-basin of the Niger River. 2021; Master's thesis. 62p.
- 23 Beaudin, 2006
- 24 Trinquier C. The risk of eutrophication of Mediterranean lagoons: The case of the Thau lagoon (Hérault). 2009 Master 1 professional Management of disasters and natural risks. Paul Valery University
- 25 USEPA (United States Environmental Protection Agency) (1976), Quality criteria for water, U.S. Environ. Protection Agency, Washington, 256 p.

26 Thomann RV, and Mueller J A. Principles of Surface Water Quality Modeling and Control. New York: Harper & Row, 1987; Pub., Inc.

UNDER PEER REVIEW