

Overview of greengram (*Vigna radiata* L.) crop, economic importance, ecological requirements and production constraints in Kenya

Abstract

Greengram (*Vigna radiata* L. Wilczek), also known as mungbean, is an important grain legume containing a high amount of digestible protein, amino acids, sugar, minerals, soluble dietary fibers, and vitamins. In Kenya, greengram production is done mainly by smallholder farmers for food and sale. The crop is mainly grown in arid and semi-arid regions with an altitude of between 50 and 1600m above sea level. It plays an important role towards achieving improved human nutrition and health conditions, reducing poverty through food security and enhancing ecosystem resilience as a source of human food, animal feed, soil nitrogen and soil health. Statistics show that though average area under production has been growing since 1978, average production has been fluctuating and consumption increasing steadily upholding constant deficit catered for through imports. The country's average greengram yield ranges between 0.5-0.6 ton/ha compared to crop potential of 1.5ton/ha and global average yield of 0.73 tons/ha. The production is mainly constrained by myriad of factors such as climate change effects, pest and disease prevalence, poor agronomic practices, land degradation, soil health decline, lack of structured marketing systems and poor research- extension- farmer linkages. Possession of limiting climate smart agriculture knowledge and skills, challenging access to credit facilities and agro processing technologies as well as narrow post-harvest loss management knowledge and skills was identified as other key constraints towards optimal greengram production within the country. It was however noted that the country has potential to achieve optimal greengram production but optimal requires adoption of climate smart technologies, improved flow of information, streamlined government policies, credit facilities as well as structured market system. The current work reviews greengram crop, with emphasis on its biology, economic importance, ecological requirements, current production status in Kenya and production constraints.

Keywords: Green grams, production constraints, arid and semi-arid regions, soil health, climate change.

1.0 Introduction

Green gram (*Vigna radiata* L.) alternatively known as mung beans, green bean, moong bean, golden gram belongs to the family leguminosae family (Udayasri et al., 2022). It is a short duration (65 - 90 days) grain legume grown on more than 6 million hectares globally in the warm areas (Malik et al., 2006; Nair et al., 2012; Hanumantharao et al., 2016). The crop is native to Indian subcontinent where it was domesticated as early as 1500 BC (Hanumantharao et al., 2016) but spread to other parts of the world through migration and trade routes. It is currently cultivated in many parts of the world such as United States of America, South Europe, Pakistan, Bangladesh, Thailand, Indonesia, Malaysia, China and Africa (Mogotsi, 2006; SASOL, 2015; Infonet, 2019). Asian continent stands out as the largest green gram's producer across the globe with India producing more than 50 % and China 19 % of total global production (Alshikh, 2019).

Greengrams or mung beans undergoes epigeal germination that takes place within 4-5 days under favorable growth factors (Sequeros et al., 2021). The plants possess well developed root system (Mogotsi, 2006) that contains many lateral roots and root nodules (Van Damme, 2007). Most varieties have many-branched stems, sometimes twining at the tips with young stems being purple or green and mature stems grayish yellow or brown and are normally covered with short hairs (Mogotsi, 2006). They can be divided into erect cespitose, semi trailing and trailing types with wild types being prostrate while cultivated types are more erect (Lambrides and Godwin, 2007; Van Damme, 2007; Sowmya et al., 2019). The crop has alternate, ovoid or broad ovoid trifoliolate leaves with elliptical to ovate leaflets, 5 -18cm long and 3-15cm wide. The crop is self-pollinated and racemes with papilionaceous pale yellow flowers are developed in the axils and leaf tips containing 4-30 flowers

per pedicel. The flowers are borne in clusters of 12–15 near the top of the plant and are hermaphroditic, have 5 petals and an ovary with one carpel, cavity and style (Sowmya et al., 2019). Depending on the variety, the resultant fruits are hairy elongated cylindrical or flat cylindrical pods, usually 30-50 per plant (Udayasri et al., 2022; Mogotsi; 2006). The pods are normally 5-10cm long and 0.4-0.6 cm wide containing 10-15 septum separated seeds which are either green, yellow, brown or blue and can be cylindrical or spherical in shape (Van Damme, 2007). Seed color and presence or absence of a rough layer usually distinguish different types of mung beans (Lambrides and Godwin, 2007). Cultivated types are generally green or golden and can be shiny or dull depending on the presence of the texture layer (Sowmya et al., 2019). Golden gram has yellow seeds, low seed yield and pods that shatter at maturity and is often grown for forage or green manure. Green grams on the other hand have bright green seeds, more prolific and ripen more uniformly, with a lower tendency for pods to shatter. Seeds are commonly green in color but sometimes yellow, brown, purplish or black depending on the type (Lambrides and Godwin, 2007). The green type is the most commonly grown in Kenya. Maturation is variety specific and most varieties takes between 60-90 days (MALF&C, 2020; Esilaba et al., 2021). A mature crop can grow up to 30 inches or 76 cm tall, having multiple branches with seed pods. Most of the seed pods become brown or darker at maturity, while others remain green.

2.0 Economic Importance of Green Grams

Green gram plays an important role as a food security crop because of its nutritional quality as well as ability to survive in harsh environmental conditions such as arid and semiarid lands (ASALs) (Yvonne et al., 2016). They are mainly grown for Mung beans are mainly grown for human food, in the form of boiled dry beans, stew, flour, sprouts and immature pods as a vegetable. According to Udayasri et a. (2022), green grams are mainly grown for their seeds which are palatable, nutritious, digestible, non-flatulent and rich source of nutrients. The grains contain approximately 25-28% protein, 1.02-1.05% oil. 3.5-4.5% fibre, 4.5-5.5% ash and 60- 65% carbohydrates on dry weight, Greengram also contains vitamin-A ((94mg), Vitamin-C (8mg), iron (7.3mg), calcium (124mg), magnesium (189mg), phosphorus (367mg) potassium (1246mg), zinc (3mg) and foliate (549mg) (Azadi et al., 2013; Udayasri et a., 2022). In addition, proteins from green gram are more easily digestible compared to proteins derived from other legumes and the amino acids profile of green gram is similar to that of soya bean (Patel et al., 2016). Unlike other pulses, it free from flatulent effects in stomach, hence, it is fed to babies and as a coalescent to elders. Green gram grains contain significant quantities of ascorbic acid, riboflavin and cholesterol free sprouts, rich in fibre and high concentration of enzymes that facilitates the digestive process (Udayasri et a., 2022).

Besides being a rich source of protein, green gram roots are important sources of soil nitrogen. The roots develop nodules that help in fixing atmospheric nitrogen in the soil through nitro-bacter bacteria in a process known as biological nitrogen fixation (Marandu et al., 2010; Weil and Brady, 2017). According to Marandu et al., 2010, the crop has the ability to add upto about 30-40kg N/ha in a single season. The vegetative parts, stocks and husks are also useful sources of leguminous fodder for livestock (Malik et al., 2006; Singh and Singh, 2011). The crop also serves as an important cover crop and a rotation crop.

5.0 Green Gram Ecological Requirement

The productivity of green gram is adversely affected by several abiotic and biotic factors such as atmospheric temperature, soil moisture, soil acidity and salinity, drought, water-logging conditions, pest infestations and disease infections that which affect crop growth and development by altering physiological processes and the plant-water relationships (Dreesen et al., 2012; Bitá and Gerats, 2013; Suzuki et al., 2014; Kaur et al., 2015; Zandalinas et al., 2017; Landi et al., 2017). As a heat and drought tolerant annual crop that grow well in low altitude areas of between 50-1600 meters above sea level, higher elevations above 1800m asl are associated with delayed pod and seed set (Mogotsi, 2006; Esilaba et al., 2021). The crop has the capacity to grow and produce in areas receiving lower rainfall ranges but optimal growth and production require well distributed rainfall of between 350-700 mm per annum (Swaminathan et al., 2012; MALF&C 2020; Esilaba et al., 2021). Higher rainfall or humidity levels either promote vegetative growth at the expense of pod formation or post-harvest losses due to delayed maturity or attract disease infestations increasing production cost. Adequate moisture is therefore required during flowering to early and late pod fill period. As a quantitative short- day plant, greengram crop is responsive to day length (Chauhan and Williams, 2018). Short days result in early flowering without adequate vegetative biomass while long days promote delayed flowering. Although different varieties vary in their photoperiod response, the crop thrive most effectively at temperatures between 25°C and 35°C (Malik et al., 2006; Esilaba et al., 2021). Significant flower shedding might occur at temperatures beyond 40°C (Zinn et al., 2010; Sita et al., 2017).

Green gram crop is well adapted to grow in wide range of soils, ranging from red laterite, black cotton and sandy textured soil. It however thrives in deep, fertile, well drained loamy to sandy loam soils with pH 6.0-7.0 (Akpapunam, 1996) while heavy clay soils restrict its root growth (KALRO, 2016; Esilaba et al., 2021). The crop shows limited tolerance to salinity and can show severe iron and chlorosis and other micronutrient deficiencies when grown in alkaline soils (Hanumantharao et al., 2016).

6.0 Green Gram Production in Kenya

Green gram locally known as Ndengu is widely grown in the arid and semi-arid lands (ASALs), either food or sale in local and export markets (ITC, 2016; MoALF, 2017; MALF&C, 2020). It is majorly grown in arid and semiarid parts of Eastern, Coast, Western, Central, Nyanza and Rift valley with Eastern accounting for 95% of national production. Some of the major production areas in this regions are: Taita Taveta, Kilifi and Tana River, Migori, Busia, Homa Bay, Kerio valley, Elgeyo Marakwet Baringo, West Pokot, Kitui, Makueni, Tharaka Nithi, Machakos, Meru and Embu (Kilimo trust, 2017; Esilaba et al., 2021). Other production areas include: Tharaka nithi, Kwale, Meru, Kisumu, Lamu, Tanariver, Kirinyanga, Bungoma and Bomet (Economic survey, 2017; GOK, 2020). Production within this region is carried out by both small scale and large scale farmers (MALF&C (2020). Most largescale farmers establish the crop as a monocrop while small scale farmers establish the crop in either monocrop, intercrop, strip or rotational cropping systems (Lambrides and Godwin, 2007). Intercropping is mainly carried out with grain crops such as sorghum, maize, pearl millet, cassava, citrus and finger millet while strip cropping is done with other legumes such as cowpea, pigeon peas, common beans and ground nuts. Rotations on the other hand is carried out with cereals such as sorghum, maize, pearl millet or finger millet.

Farmers within the country alternate between local and improved varieties as seed stock (Kilimo trust, 2017; GOK, 2020). According to MAF&C, 2020 and Esilaba et al., 2021, the local varieties are well adapted to the environment but are associated with small seeds, varied maturity time, low yields compared to improved varieties, do well in dry areas and some might have many stony seeds, which makes green gram meal difficult to eat. The common improved varieties within the country are N22 or KVR22, N26 (nylon) or KVR26, KS20 (uncle/cotton), Biashara, Karemba and Ndengu tosha (Esilaba et al., 2021). The grain yield potential of each variety per 90kg bag is N22 (4-7bags/acre), N26 (6-8bags/acre), KS20(7-10bags/acre), Biashara (8-9bags/acre), Karemba(8-9bags/acre), Ndengu tosha(8-10bags/acre) (KALRO, 2016; Esilaba et al., 2021). The above improved varieties have been developed and released by KALRO in collaboration with International Centre for Tropical Agriculture (CIAT), International Institute for Tropical Agriculture (IITA) and other partners (MALF&C, 2020).

Varieties KS20 (Nylon) and N26 (Uncle) are commonly categorized as popular varieties with farmers across the country. According to Mugo (2018) and Esilaba et al. (2020), KS20 variety has a dull green color, and its pods turn brown when dry, it is tolerant to aphids, resistant to yellow mosaic virus and performs poorly in dry environments. Variety N26 on the other hand is a high yielding variety with shiny green seed color and pods turn black when dry. It possesses uniform early maturity and does well in dry areas. Both varieties are propagated by seed. For rain fed ecosystems, planting is carried out at onset of rains to avoid crop failure or reduced yield. Depending on variety and environmental conditions, germination occurs within 5-7 days. In areas with higher rainfall, it is recommended to grow green grams on raised beds (Singh and Singh, 2011). For maintenance of soil health and optimal crop yields, application of basal or foliar fertilizers to supplement limiting nutrients is recommended. Because the crop is commonly grown in ASALs where pest are prevalent, integrated pest and disease management practices are recommended. Timely harvesting is also recommended to avoid field losses through shattering of the pods. Depending on the prevailing environmental conditions, harvested seeds are dried for about 2-3 days then thresh and winnow, ready for consumption or storage. Green grams are susceptible to attack by bruchids; storage in airtight containers or gunny bags in a clean ventilated room immediately after drying is recommended for short period storages and seed treatment for long term storage. The general green gram value chain in Kenya normally comprises of Farm to local market -> Local market to wholesaler or miller -> Imports to wholesaler -> Wholesale market miller to retailer -> Retailer to consumer who are end user.

According to GOK, 2020 and KIHBS, 2015-2016, area under production of green grams has been rising steadily from 1978 to 2017, greengram production has been fluctuating while consumption seem to increase steadily over the years. For example: Kenya recorded average production of 121.1MT of green grams in 2014

and 104.9MT in 2015, 115MT in 2016 and 107.6 in 2017 compared to average consumption of 571.9MT in 2014, 591.8MT in 2015, 603.8MT in 2016 and 659.5MT in 2017. This led to an average deficit 450.8MT in 2014, 487MT in 2015, 488.8MT in 2016 and 552.1MT in 2017. The constant production-consumption deficit growth suggests opportunities for enhancing domestic production to substitute imports. According to Kilimo Trust (2017) and MALF&C (2020), the current countries average green gram yield therefore lies within 0.5-0.6 tons/ha which fall far below the countries crop potential and compares unfavorably with the global average yields of 0.73 tons/ha. This production is not able to meet the countries demand and Kenya takes up about 80% of the green grams exported by Uganda and Tanzania.

7.0 Factors Affecting Green Gram Production in Kenya

Despite the growing population and increased demand for climate smart crops such as green grams within the country, green gram industry growth has been constrained over the years. The low greengram industry performance and expansion can be attributed to myriad of factors such as: Pest and disease prevalence, impact of climate change, land degradation, soil health decline, limited access to good quality seeds, erratic rains leading to water scarcity, weed infestation, land tenure systems and land fragmentation, unstructured marketing systems, post-harvest losses, limited primary processing technologies and services, lack of harvesting mechanization technologies as well as poor agronomic technical knowledge (Karanja et al., 2006; KALRO, 2016; MOALFC, 2020; Chabari, 2020; Wambua, 2021).

7.1 Pest and Disease Prevalence

Pest and diseases pose a huge production challenge to small scale farmers because of either lack of management knowledge or capital or low farm gate prices that do not offer farmers incentive to invest in control measures (D'Alessandro et al., 2015; Esilaba et al., 2021). Some of the common greengram pests within the country are: flower thrips (*Megalurothrips sjostedti*), aphids (*Aphis sp.*), pod bugs (*Riptortus pedestri*), white flies (*Bemisia tabaci*), foliage beetles, cutworms (*Agrotis spp*), root-knot nematodes (*Meloidogyne spp*), leaf miner (*Lyriomyza spp*), pod borer (*Etiella zinckenella*), red spider mites (*Tetranychus sp*), Grass blue butterfly (*Euchrysops cnejus spiny*), African boll worm (*Helicoverpa armigera*) and bruchids (Swaminathan et al., 2007; SASOL, 2015; KALRO, 2016; Infonet, 2019). Many of these pest attack greengrams in the field while others such as bruchid cause considerable losses during storage. They pests have been documented to cause between 30-100% losses. Their management also increases cost of production for small scale poor farmers (Swaminathan et al., 2007).

The common greengram diseases within the country include: Anthracnose (*Colletotrichum lindemuthianum*), Bacterial leaf blight (*Xanthomonas phaseoli*), Cercospora leaf spot (*Cercospora canescens*), Leaf crinkle disease (Leaf Curl Virus), Macrophomina blight (*Macrophomina phaseolina*), Yellow mosaic (Mungbean yellow mosaic virus), Powdery mildew (*Erysiphe polygoni*), Root rot (*Macrophomina spp: Fusarium spp*), Leaf blight (*Rhizoctonia solani*) and rust (*Uromyces phaseoli*) (SASOL, 2015; KALRO, 2016; Infonet, 2019). The diseases can have transmitted through contaminated seeds, soil, farm implements or alternate hosts. Loses of up to 10-45% have been reported due to common blight while 80% has been reported due to angular leaf spot disease (Pyndji, 1992; Wortmann et al., 1998; Karanja et a., 2006; Esilaba et al., 2021). Their management also increases cost of production for small scale poor farmers leading to low return to capital (AIC, 2002; KALRO, 2016).

7.2 Climate Variability and Limited Climate Smart Agriculture Knowledge and Skills

Most greengram cultivation areas located in arid and semi-arid areas which are designated to be sensitive to climate variability and highly vulnerable to such as irregular and unpredictable rainfall, frequent droughts during cropping seasons or occasional severe floods (Maliva and Missimer, 2012; IPCC, 2018; Spinoni et al., 2020). Owing to the fact that optimal crop growth and yield require provision of all nutrients in rightful forms and amounts, water plays a major role as a solvent for all soil chemical reactions that eventually determine germination, nutrient uptake, metabolisms, assimilation, photosynthesis and all other plant biochemical processes (Weil and Brady, 2017; Muindi et al., 2022). Most of the greengram production areas within the country have been experiencing erratic rains and frequent drought over the past years leading to up to 79% decrease in production in some seasons (Marshall, 2011; MoALF, 2016). The constant dry spells coupled with poor water conservation technologies and limited climate smart agricultural practices knowledge and adoption among other factors play an important role towards below optimal green gram production within the country (MOALFC, 2020).

7.3 In Adequate Access to Agricultural Information and Extension Services

Access to adequate agricultural information is vital to increased agricultural productivity (Kelil et al., 2020). According to (Yaseen et al., 2016), lack of information sources in rural areas restrains farmer's agricultural production. Public agricultural extension officers play an important role of linking farmers with current technologies and market. The research- extension – farmer link in Kenya has however weakened over the years with the current extension: farmer ratio standing at 1800:1 compared to the FAO recommendation of 400:1 (MOALF, 2020). As a result, many rural smallholder farmers have limited avenues to source for useful and reliable advice and information on soil and water management, plant nutrition, seed quality, agronomic activities, value addition and marketing information leading to low uptake of technologies, poor crop production and resultant yields within the farming communities. Some farmers on the other hand who are able to produce on their own and get good harvest face the challenge of sourcing for readily available reliable markets ending up suffering either post-harvest costs and losses or selling at throw away prices.

7.4. Access to Seeds

Although seed is among the most central production resources that greatly affect productivity, setting up seed production and delivery systems that encourage extensive use of quality seed throughout the marketing chain has been a challenge in Kenya for many years (Muthoni, and Nyamongo, 2008). In an effort to improve the countries agricultural production as part of institutional, policy, and regulatory reforms, the government of Kenya over the years have strived to come up and enact national seed policies and regulation. The policies and regulations include: The National Seed Policy, 2010, Seeds and Plant Varieties (Seeds) Regulations, 2016, Kenya Agricultural and Livestock Research Act of 2013, Crops Act no. 16 of 2013 (revised in 2016) and, Seeds and Plant Varieties Act (Cap 326) (Ayieko *et al.*, 2021). While this national seed policies, regulations and formal seed system exists within the country, a good number of farmers have been reported to rely on informal seed system as seed source (MOALFC, 2020; GOK, 2020; Esilaba et al., 2021). This is typical for most greengram farmers within the country. According to Muthoni and Nyamongo (2008) and MOALFC (2020), many of these farmers prefer growing traditional varieties because the landraces are well adapted to local conditions and have low nutrient requirements. The local varieties are however associated with low yields, long maturity periods (3 to 4 months), small seeds, prone to shattering before harvest thus high postharvest losses, susceptibility to several pest and diseases, and high percentage of stony and hard to cook grains which reduces their preference for consumption (MOALFC, 2020; Esilaba et al., 2021).

7.5 Access to Credit Facilities

According to research carried out by Ndambiri et al. (2012) and Njuguna and Nyairo (2015), there exists a positive relationship between new technologies adoption level and the availability of credit. Availability of credit eases the cash constrictions and enabling farmers to acquisition inputs such as improved seeds, fertilizer, pesticides with ease. Lack of collateral among rural small scale farmers in arid and semiarid areas nonetheless limit their right to benefit from credit facilities (Mbugua, 2013; Musembi, 2019; GOK, 2020). Furthermore, though government policies play a major role in cushioning farmers, and mung bean bill, 2020 exists, the benefits of its implementation is yet to reach many small holder farmers within the country.

7.6 Land Degradation, Access and Use of Fertilizers in Greengram Farming

Land degradation is a major problem facing food production and sustainable development in most arid and semi-arid environments of Sub-Saharan Africa. It is defined by Oluwole and Sikhhalazo 2008; UNCCD, 2013; Mbow et al. 2015 as a complex and multifaceted phenomenon that reduces current or future capacity of either rain-fed cropland, irrigated cropland, range, pasture, forest, and woodlands to produce. The land degradation forms of importance in green gram production ecosystems include: soil erosion, deterioration of soil physical, chemical, biological or economic properties and, long-term loss of natural vegetation and biota among others (UNCCD 2013). According to Weil and Brady (2017), adequate availability of nutrients in rightful proportions is paramount in crop growth and productivity. The seventeen nutrients required by plants include: carbon (C), hydrogen (H), oxygen (O), nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), sulfur (S), zinc (Zn), iron (Fe), copper (Cu), manganese (Mn), chlorine (Cl), boron (Bo) and molybdenum (Mo) (Brady and Weil, 2008; Muindi et al., 2019). Availability of these nutrients for crop uptake is, however, dictated by prevailing soil physical, chemical and biological properties which are often compromised in degraded lands. As a vital soil physical property, moisture which is compromised in degraded lands acts as a solvent for all chemical reactions which eventually influences nutrient sorption and uptake as well as regulating favorable rhizosphere environment. Availability of adequate moisture is however dictated by other physical attributes such as soil texture, structure, consistence, horizonation and bulk density (Weil and Brady, 2017) which are habitually compromised in most degraded lands. Soil pH on the other hand is a chemical property that regulates solubility and availability of all nutrients as well as soil biota population and activity (Brady and Weil, 2018). Nutrients are soluble and available for uptake at pH range 6.0 - 7. 2. At low pH levels of less than 5.5

most macro nutrients such as Ca, Mg, P and K become unavailable for plant uptake. Majority of micronutrients will, however, be abundant, very soluble and toxic at low pH range (Muindi et al., 2015). The opposite is true when pH levels are above 7.2 because most base cations such as calcium, Magnesium and sodium become abundant, soluble and toxic while micronutrients such as Zn, Bo, Mo, and Fe become insoluble and un available (Brady and Weil, 2008). These scenario calls for soil fertility management by supplementation of limiting nutrients through application of either farm yard manure or inorganic fertilizers (Muindi et al., 2019). This is challenging in majority of green gram growing areas because farmers grow greengram as a mono-crop, season after season, without adding amendments for soil improvement (GOK, 2020). This leads to nutrient mining more so for other nutrients apart from Nitrogen which is fixed through biological Nitrogen fixation leading to poor yields and return to capital. Farmers who use commercial fertilizers also often use straight fertilizers that don't provide all limiting nutrients within their soils.

The current withdrawal of Kenyan government from fertilizer market and abandonment of price controls to encourage private investors have led to improved fertilizer distribution but led to increased prices which are unaffordable to most smallholder farmers. Additionally, as a leguminous crop with ability to fix atmospheric nitrogen in soils, green gram farmers are not prioritized in the current subsidized fertilizer scheme operated by the government.

According to research carried by GOK (2020), Fertilizer usage among green gram farmers is relatively low with some farmers not applying fertilizers and others using sole manure. There however exists a relationship between soil moisture availability and mineral fertilizer usage. According to myths from farmers that "fertilizers burn crops, harden soils or dries up crops prematurely". The myths can be scientifically explained by the relationship between fertilizer application timing, prevailing soil moisture levels and soil properties such as organic carbon and water holding capacity. Poor soil fertilization is associated with poor soil health leading to low production.

7.7 Poor Structured Marketing Systems for Green Grams

According to Kihoro, (2016) and MALF&C (2020), there exists no structured marketing system for green grams in Kenya. Farmers marketing channel choices are therefore determined by socio-economic, institutional, and farm level factors. This acts as a setback for small scale poor famers with limited access to credit facilities, market based signals and information.

7.8 Post Harvest Losses and Management

Postharvest loss includes food loss across the food supply chain from crop harvesting until consumption (Aulakh et al., 2013). The losses can be broadly categorized as spoilage, quality loss, nutritional loss, seed viability loss and commercial loss [Boxall, 2001] attributable to lack of knowledge, inadequate technology and/or poor storage infrastructure. They account for between 20% and 40%, of total production and can go up to 80% in severe cases (Fox, 2013; Abass et al., 2014). Within Kenyan green gram sector, post-harvest losses occur during operations such as threshing, winnowing, transportation, processing and storage (Esilaba et al., 2021). The main cause of this postharvest losses is spoilage, due to poor and prolonged storage, spillage during transportation, breakages during threshing, destruction by rodents such as rats and attack by weevils (GOK, 2020). Depending on the levels, post-harvest losses arising from poor storage, handling of produce and poor infrastructure, can make farmers to lose their crops' full value, forcing them to sell the grain throw away price (MALF&C (2020).

7.9 Access to Agro Processing Technologies

According to Kirsten et al. (2013) and MALF&C (2020), farmers access to agro processing technologies is key to ensure profitability of a product after harvesting; something that limits crop production in most African countries. Most greengram farmers within the country either don't have access to agro processing facilities or have access to old and outdated processing methods. This exposes them to high losses leading to low enterprise profitability.

7.10 Weed Management

Weeds are one of the major problems in green gram cultivation, reducing the yield through competition, interference with harvest and harboring especially during periods with favorable growing conditions. Soil moisture availability causes weeds to multiply very fast and outgrow the main crop if intervention is not carried out on time. According to Kumar et al., (2021). weed competition can cause up to 54% total yield losses in green grams.

7.11 Land Fragmentation

Land fragmentation is a universal trait of all agricultural systems which affects farmland productivity and use of machinery. Kenyan agricultural land range from large to small (0.3-3ha) and very small land holdings of less than 0.16 ha. The very small sizes limit green gram production enterprise mechanization and economic viability posing crop productivity intensification challenge.

8.0 Conclusion and Recommendations

The review indicates that green gram plays a significant role achieving improved human nutrition and health conditions, lowering poverty through food security and enhancing ecosystem resilience. They act as source of human food, animal feed, soil nitrogen and soil health. It is also a wonder plant that can grow well in arid and semiarid areas. Statistics show that although area under production has been increasing since 1978, production has been inconsistent while consumption has continued to rise steadily maintaining a deficit that is catered for through imports. Some of the factors that constrain green gram production within Kenya include: climate change effects, pest and diseases prevalence, land degradation and soil health decline due to nutrient mining without adequate replenishment. Other factors such as poor access to agricultural information, credit facilities, poor structured marketing system as well as access to agro processing technologies play a key role in constraining greengram production within the country. Land fragmentation, poor post-harvest losses management and poor seed production and delivery systems were also identified to be key limiting factors. The review revealed that the country has a potential for optimal production and achievement of optimal production potential, that meets increasing demand requires adoption of climate smart technologies, improved flow of information to farmers, government policies, credit facilities as well as structured market system.

9.0 References

- Abass AB, Ndunguru G., Mamiro P, Alenkhe B, Mlingi N, Bekunda M. Post-harvest food losses in a maize-based farming system of semi-arid savannah area of Tanzania. *Journal of Stored Products Research*. 2014;57:49–57.
- AIC. *Field Crops Technical Handbook*. 2002.
- Akapunam M. Mung bean (*Vigna radiata* (L.) Wilczek). In: Nwokolo, E., Smartt, J. (eds) *Food and Feed from Legumes and Oilseeds*. Springer, Boston, 1996: Pgs 209-215.
- Alshikh ZA S. Effect of DAP fertilizer and Seed Rate on Two Varieties of Mung bean (*Vigna radiata* L. Wilczek) (Master's thesis). Sudan University of Science and Technology, Sudan. 2019.
- Aulakh J, Regmi A, Fulton JR, Alexander C. Estimating post-harvest food losses: Developing a consistent global estimation framework; Proceedings of the Agricultural & Applied Economics Association's 2013 AAEE & CAES Joint Annual Meeting; Washington, DC, USA. 4–6 August 2013.
- Ayieko M, Odame H, Olwande J. Strengthening Seed Systems and Market Development in Kenya: Perspectives on Political Economy and Policy Processes. Tegemeo Institute of Agricultural Policy and Development, Nairobi, Kenya. 2021: 51Pgs.
- Azadi E, Rafiee M, Nasrollahi H. The effect of different nitrogen levels on seed yield and morphological characteristic of mungbean in the climate condition of Khorramabad. *Annals of Biological Research*, 2013: 4(2), 51-55.
- Bitá C E, Gerats T. Plant tolerance to high temperature in a changing environment: scientific fundamentals and production of heat tolerance crops. *Frontiers of Plant Sciences* 2013: 4, 273–296.
- Boxall RA. Post-harvest losses to insects—A world review. *Int. Biodeterior. Biodegradation*, 2001: 48:137–152.
- Brady NC, Weil R R. *The nature and properties of soils.*: Prentice Hall. 2008: 5(13), 662-830.
- Chabari K. Socio-Economic Factors Affecting Green Grams Production a Case Study of Small Holder Farmers in Tharaka District. Msc. Thesis. University of Embu, Kenya. 2020.
- Chauhan YS, Williams R. Physiological and Agronomic Strategies to Increase Mungbean Yield in Climatically Variable Environments of Northern Australia. *Agronomy*. 2018; 8(6):83.
- D'Alessandro S, Caballero J, Simpkin S, Lichte J. Kenya Agricultural Risk Assessment. Agriculture Global Practice Note 17. World bank group. 2015: 8pgs.
- Dreesen F E, De Boeck HJ, Janssens IA, Nijs I. Summer heat and drought extremes trigger unexpected changes in productivity of a temperate annual/biannual plant community. *Environ. Exp. Bot.* 2012: 79, 21–30.

Economic survey. Kenya National Bureau of statistics. Nairobi, Kenya. 2017: 333Pgs.

Esilaba AO, Nyongesa D. Okoti M. Otipa M, Wasilwa L. KCEP-CRAL Green Gram Extension Manual. Kenya Agricultural and Livestock Research Organization, Nairobi, Kenya. 2021 58Pgs.

Fox T. Global Food: Waste Not, Want Not. Institution of Mechanical Engineers; Westminster, London, UK: 2013.

GOK Can Green Grams Enhance Food and Nutrition Security in Kenya? Evidence from top eight Green Grams Producing Counties in Kenya. The National treasury and planning state department for planning. 2020: 66 Pgs.

Hanumantharao B, Nair RM, Nayyar H. Salinity and High Temperature Tolerance in Mungbean [*Vigna radiata* (L.) Wilczek] from a physiological perspective. *Frontiers in Plant Science*, 2016: 7, 1-20.

IPCC. Special Report on the Impacts of Global Warming of 1.5°C above Pre-Industrial Levels and Related Global Greenhouse Gas Emission Pathways, in the Context of Strengthening the Global Response to the Threat of Climate Change, Sustainable Development, and Efforts to Eradicate Poverty, World Meteorological Organization, Geneva, Switzerland. 2018.

ITC(International trade Centre). Annual Report.2016.

KALRO. KALRO-KCEP greengram production extension brochure. KALRO, Nairobi Kenya. 2016: 7Pgs.

Karanja DR, Githunguri CM, M'Ragwa L, Mulwa D. Variety, Characteristics and Production Guidelines of Traditional Food Crops. KARI Katumani research centre. 2006:38Pgs.

Kaur R, Bains TS, Bindumadhava H, Nayyar H. Responses of mungbean (*Vigna radiata* L.) genotypes to heat stress. Effects on reproductive biology, leaf function and yield traits. *Sci. Hortic*. 2015: 197, 527–541.

Kelil A, Girma Y, Hiruy M. Access and Use of Agricultural Information in Africa: Conceptual Review. *Information and Knowledge Management*, 2020. 10(7): 1-5.

KIHBS (Kenya Integrated Household Budget Survey) Kenya National Bureau of Statistics. 2015-2016. : 133Pgs. <https://statistics.knbs.or.ke/nada/index.php>

Kihoro EM. An analysis of factors influencing farmers' choice of green gram marketing channels in Mbeere south subcounty, Kenya. Msc. Thesis. University of Nairobi, Kenya. 2016:118Pgs.

Kilimo Trust. Characteristics of greengram markets in the EAC: Regional East African Community Trade in Staples (REACTS) Project. 2017.

Kirsten J, Mapila M, Okello J. De S. Managing agricultural commercialization for inclusive growth in Sub-Saharan Africa. *Global Development Network Working Paper No. 60*. 2013.

Kumara P, Pakeerathan K, Deepani LPP. Assessment of Yield Loss in Green Gram (*Vigna radiata* (L.) R. Wilczek) Cultivation and Estimation of Weed-Free Period for Eco-Friendly Weed Management. *Biology and Life Sciences Forum*. 2021; 3(1):22.

Lambrides C J, Godwin ID. Mungbean. In: Chittarajan, K., *Genome Mapping and Molecular Breeding in Plants*, 2006: 3: 69-90

Landi S, Hausman J F, Guerriero G, Esposito S. Poaceae vs. abiotic stress: focus on drought and salt stress, recent insights and perspectives. *Frontiers of Plant Sciences*. 2017: 8:1214.

MALF&C. Climate Smart Greengram (*Vigna radiata*) Resource and Training Guide. 2020.

Malik A, Fayyaz-ul-Hassan WA, Qadir G, Asghar R. Interactive Effects of Irrigation and Phosphorus on Green Gram (*Vigna radiata* L.). *Pakistan Journal of Botany*, 2006: 38, 1119-1126.

Maliva R, Missimer T. Aridity and drought Arid Lands Water Evaluation and Management, Springer, Berlin, Heidelberg. 2012: pp. 21-39

Marandu AET, Semu E, Mrema JP, Nyaki AS. Quantification of Atmospheric N₂ Fixed by Cowpea, Pigeonpea and Greengram Grown on Ferralsols in Muheza District, Tanzania. *Tanzania Journal of Agricultural Sciences*, 2010: 10 (1): 29-37.

Marshall S. The Water Crisis in Kenya: Causes, Effects and Solutions. *Global Majority E-Journal*, 2011: 2 (1):31-45

Mbow C, Brandt M, Ouedraogo I, de Leeuw J, Marshall M. What four decades of earth observation tell us about land degradation in the Sahel? *Remote Sensing* 2015: 7, 4048–4067

Mbugua I. Factors Determining Access to Credit Facilities for Farmers in Cherangany Constituency in Trans-Nzoia County. MBA thesis. University of Nairobi. 2013: 56 Pgs.

MoALF. Kenya Climate Smart Agriculture Strategy-2017-2026. 2017.

MoALF. Climate Risk Profile for Makeni. Kenya County Climate Risk Profile Series. The Kenya Ministry of Agriculture, Livestock and Fisheries (MoALF), Nairobi, Kenya. 2016.

MOALF. Situational analysis of the agriculture sector in Kenya. Final Report. Nairobi, Kenya. 2020: 154Pgs

Mogotsi KK. *Vigna radiata* (L.) R. Wilczek. In: Brink, M. & Belay, G. (Editors). PROTA 1: Cereals and pulses/Céréales et légumes secs. [CD-Rom]. PROTA, Wageningen, Netherlands. 2006.

- Mugo JW. Modelling green gram production in Kenya under the current and future climates. PhD. Thesis. University of Nairobi, Kenya. 2018: 169 Pgs.
- Muindi EM, Mrema JP, Semu E, Mtakwa PW, Gachene CK, Njogu MK. Lime-aluminium-phosphorus interactions in the Kenya Highlands. *American Journal of Experimental Agriculture*, 2015: 9 (4):1-10.
- Muindi EM, Muindi CM, Ndiso JB. The Effects of Combining Farm Yard Manure, Starter Nitrogen, Phosphorus and Zinc on Growth and Yield of Green Grams. *Journal of Agriculture and Ecology Research International*. 2019: 20(4):1-9
- Muindi EM, Wamukota AW, Mulinge JM, Okello N, Wekesa G, Ahmed H. Onion (*Allium cepa* L.) Performance as influenced by manure and fertilizer in titanium mined soils. *Sustainable Agricultural Research*, 2022: 11(3):27-37.
- Musembi EK. Demand for agricultural credit by rural smallholder farmers: a case of climate smart agriculture villages in nyando basin, Kenya. MA thesis. University of Nairobi. 2019: 77 Pgs.
- Muthoni J, Nyamongo DO. Seed Systems in Kenya and Their Relationship to On-Farm Conservation of Food Crops. *Journal of new seeds*, 2008: 9(4): 330-342.
- Nair RM, Schafleitner R, Kenyon L, Srinivasan R, Easdown W, Ebert RW, et al., Genetic improvement of mungbean. *SABRAO Journal of Breeding and Genetics*, 2012: 44: 177-190
- Ndambiri HK, Ritho C, Mbogoh SG, Ng'ang'a SJ, Muiruri EJ, Cherotwo FH. Assessment of farmers' adaptation to effects of climate change in Kenya. *Journal of economic and sustainable development*: 2012: 3(12).
- Njuguna E, Nyairo N. Formal Conditions that Affect Agricultural Credit Supply to Small-scale Farmers in Rural Kenya: Case Study for Kiambu County. *International Journal of Sciences: Basic and Applied Research*, 2015: 20 (2):59-66
- Oluwole FA, Sikhalazo D. Land degradation evaluation in a game reserve in Eastern Cape of South Africa: soil properties and vegetation cover. *Scientific Research and Essays*, 2008: 3(3), 111–119.
- Parihar AK, Basandrai AK, Sirari A, Dinakaran D, Singh D, Kannan K, et al. Assessment of mungbean genotypes for durable resistance to Yellow Mosaic Disease: Genotype x Environment interactions. *Plant Breeding*, 2017: 36, 94–100.
- Patel SA, Chaudhari PP, Desai NH. Yield and economics of green gram (*Vignaradiata* (L.) Wilczek) Cultivars as Influenced by Integrated Nutrient Management. *Crop Research*, 2016: 51:1
- Pyndji, Trutmann P. Managing angular leaf spot on common beans in Africa by supplementing farmer mixtures with resistant varieties. *Plant Disease*, 1992:76:1144-1147
- SASOL (Sahelian Solution Foundation). Green Grams Hand Book: A guide to farmers within project enhancing food security with diversified Dry land farming techniques in Kitui County. 2015: [Http://www.SASOL_foundation.co.ke/2013/wp-content/uploads/2015/02/Green-Grams-Manual.pdf](http://www.SASOL_foundation.co.ke/2013/wp-content/uploads/2015/02/Green-Grams-Manual.pdf)
- Sequeros T, Ochieng J, Schreinemachers P. et al. Mungbean in Southeast Asia and East Africa: varieties, practices and constraints. *Agriculture and Food Security* 2021: 10: 2.
- Singh JS, Singh D P. Efficient soil microorganisms: a new dimension for sustainable agriculture and environmental development. *Agriculture, ecosystems & environment*, 2011: 140(3-4): 339-353.
- Sita K, Sehgal A, Hanumantha Rao B, Nair RM, Vara Prasad PV, Kumar S, et al. Food legumes and rising temperatures: effects, adaptive functional mechanisms specific to reproductive growth stage and strategies to improve heat tolerance. *Frontiers of Plant Sciences*, 2017: 8: 1658. doi: 10.3389/fpls.2017.01658
- Sowmya T, Durga KK, Venkateshwaran K, Keshavulu K, Vidyasagar. Characterization of Green Gram Genotypes for Qualitative Traits. *Agricultural Sciences Digest*, 2019: 39(2):81-89.
- Spinoni J, Barbosa P, Bucchignani E, Cassano J, Cavazos T, Christensen JH, Dosio A. Future global meteorological drought hot spots: a study based on CORDEX data. *Journal of Climate*, 2020: 33 (9): 3635-3661.
- Suzuki N, Rivero RM, Shulaev V, Blumwald E, Mittler R. Abiotic and biotic stress combinations. *New Phytol*. 2014: 203: 32–43. doi: 10.1111/nph.12797
- Swaminathan R, Singh K, Nepalia V. Insect Pests of Green Gram *Vigna radiata* (L.) Wilczek and Their Management. *Insect Pests of Green Gram Vigna radiata* (L.) Wilczek and Their Management, Agricultural Science, 2012. Dr. Godwin Aflakpui (Ed.), ISBN: 978-953-51-0567-1, InTech,
- Swaminathan R, Singh K, Nepalia V. (2007). *Insect Pests of Grams Vigna radiata* (L.) Wilczek and Their Management, 2007.
- Udayasri S, Lavanya GR, Lal GM. Estimation of Variability and Genetic Divergence in Greengram [*Vigna radiata* (L.) Wilczek] for Yield Characters. *International Journal of Plant & Soil Science*, 2022: 34(23): 49-56.
- UNCCD. Background Document, The Economics of Desertification, Land Degradation and Drought: Methodologies and Analysis for Decision-Making. In 2nd Scientific Conference on Economic

- Assessment of Desertification, Sustainable Land Management and Resilience of Arid, Semi-Arid and Dry Sub-Humid Areas. April 9–12, 2013—Bonn, Germany.2013.
- Van Damme P. "Plant Resources of Tropical Africa 1. Cereals and Pulses," *Economic Botany*, 2007: 61(1): 108.
- Wambua MJ. Analysis of factors influencing productivity and extent of Smallholder commercialization of green grams and pigeon peas In machakos county, kenya. PhD Thesis, Egerton University, Kenya. 2021: 285 Pgs.
- Weil R, Brady NC. The nature and properties of soil. 15th edition. Pearson. 2017: 1104pp
- Wortmann C, Kirkby R, Eledu C, Allen D.. Atlas of common bean (*Phaseolusvulgaris* L.) Production in Africa. ICali.Colombia CIAT pulbication No. 297. 1998: SBN958-9439-94-2.
- Yaseen M, Xu S, Yu W, Hassan S. Farmers' Access to Agricultural Information Sources: Evidences from Rural Pakistan. *Journal of Agricultural Chemistry and Environment*, 2016: 5: 12-19.
- Yvonne M, Richard O, Solomon S, George K. Farmer Perception and Adaptation Strategies on Climate Change in Lower Eastern Kenya: A Case of Finger Millet (*Eleusine coracana* (L.) Gaertn) Production. *Journal of Agricultural Science*, 2016: 8: 33-40.
- Zandalinas SI, Sales C, Beltrán J, Gómez-Cadenas A, Arbona V. Activation of secondary metabolism in citrus plants is associated to sensitivity to combined drought and high temperatures. *Frontiers of Plant Sciences*, 2017: 7: 1954. doi: 10.3389/fpls.2016.01954
- Zinn KE, Tunc-Ozdemir M, Harper JF. Temperature stress and plant sexual reproduction: uncovering the weakest links. *Journal of Experimental Botany*, 2010: 61:1959–1968. doi: 10.1093/jxb/erq053