

# **Field and Petrographic Studies of the Guwahati metapelites, Assam, NE India: Evidence for Post-Metamorphic K-Metasomatism**

---

## **ABSTRACT**

In this study, we report mineralogical features of post-metamorphic K-metasomatic assemblage of high-grade metapelites from Guwahati, the northern extension of the central part of the Shillong Plateau, NE India. Several reaction textures of metasomatic metapelites are shown and they offer clues for relevant reactions for metasomatic modification of pre-metasomatic metapelites. Two sets of mineral reactions are recognized, which include, the alteration of high- temperature of all kinds of aluminium - silicate phases of pre-metasomatic metapelites into low-grade minerals and the replacement of coexisting metamorphic plagioclase and quartz by K-feldspar due to the involvement of a K-rich fluids at the end of all deformational events. Our structural study indicates the rocks of the area are intensely deformed with multiple folding events and post-kinematic shearing and faulting which cross-cut the dominant foliation of the rocks. The deep-seated K-rich metasomatic fluids, moved into shear zones and migrated along the foliation plane, grain boundaries of minerals transforming the parent metapelites into metasomatic rocks. As a result of metasomatic changes, all pre-existing features disappear and metapelites turn into K-feldspar -rich granitic rock.

*Key words: Cordierite, Shillong Plateau, K-metasomatism, Myrmekite.*


## **1.INTRODUCTION**

Metapelites from several localities in the high-grade terrain of the Shillong Plateau have received much attention during the past few decades. The hilly tracts of Meghalaya and some parts of Goalpara and Kamrup districts of Assam expose high-grade metapelites with profusely developed monazite, xenotime and zircon which have immense potentiality in dating the geological events. The U–Th– Pb<sub>total</sub> monazite, xenotime and zircon geochronology provide a powerful technique to elucidate the timing of thermal events for establishing their relationship with regional and global tectonic processes. For this reason, they have been subjected to extensive petrological, chronological, geochemical and tectonic scrutiny


by several workers [1,2,3,4,5,6,7,8,9,10]. However, the petrological, thermochronological history of the Guwahati metapelites [3] is somewhat cloudy and it is obscure whether these high-grade metapelites are a continuation of those situated towards the south in Meghalaya [4,5,7,8]. In this present study, observation on our structural and petrographic studies of the Guwahati metapelites demonstrates clear evidence of the involvement of post-metamorphic K-rich fluids in transforming highly deformed metapelites into metasomatic rocks.

## **2.GEOLOGICAL BACKGROUND**

The study area, Guwahati (26.1158° N, 91.7086° E) represents a northern extension of the central part of the gneissic complex of the Shillong Plateau (Fig.1a). The Shillong Plateau represents an extreme north-eastern extension of the Precambrian Indian Peninsula and is made up of the Neoproterozoic–Proterozoic to Early Palaeozoic (2.6–0.5 Ga) basement gneissic rocks [4,5,11,12], the Paleo-Mesoproterozoic intracratonic Shillong basin consisting of metasedimentary supracrustal rocks of the Shillong Group [13] and Proterozoic–Early Palaeozoic (881–479 Ma) equigranular to porphyritic granitoid plutons intruding into the basement gneisses and the Shillong Group of rocks [11,14,15,16,17,18] suggested the region as a north-eastern and eastern extension of the Eastern Ghat Belts, and the Chottanagpur Gneissic Complex respectively. The western part of the Shillong Plateau (Garo-Goalpara Hills region) is characterised by predominantly metabasic rocks with garnet corona, co-folded with quartzofeldspathic, migmatitic gneisses, orthopyroxene bearing metapelites, and post-kinematic dioritic intrusion. On the contrary, the central Shillong Plateau (e.g. Nongstoin, Kynshi, Sonapahar, Patharkhammah, Guwahati) is dominated by metapelites with coronitic cordierite and garnet-free metabasic rocks, banded iron formations, calc-silicate gneisses [19,2]. The deformational history of the Shillong Plateau is characterized by different episodes of folding ( $D_1$ ,  $D_2$ ,  $D_3$  and  $D_4$ ) and a ductile to brittle–ductile shearing episode pre-dates the last folding  $D_4$  deformation [19,20,21]. The N–S sinistral shears and mylonitization are seen to superpose the folds of earlier generations [19,20,22]. The widespread distribution of varying metamorphic grades of the characteristic NNE-trending sinistral shears and related folds throughout the Shillong Plateau indicates the early Palaeozoic southward movement of the Indian block relative to the Australo-Antarctic block in the east [23].



**Fig.1. Generalized geological map of the Shillong plateau showing the location of Guwahati, study area (GSI data)**



**Fig. 2. (a) Occurrence of metapelites as small-scale salvages of biotite within K-feldspar rich granite. (b) Thin bands of gneissic rocks in association of relics of pre-metasomatic metapelites in the intense zone of deformation. where the shear zone cross cut the foliation plane. Replacement of shear zone as well as foliation plane by K-feldspar as cross cut metasomatic veins. Note the presence of small pods of metasomatic K-rich granite at the central left part of the figure.**

The Precambrian rocks of Guwahati are co-folded bands of hornblende, biotite - bearing tonalitic gneiss, quartzofeldspathic gneiss, calc-silicate gneiss, amphibolite and cordierite - bearing metapelites, medium- to coarse-grained granite and post -kinematic grey porphyritic granitic rocks. The metapelites are scarce in the area and our collected samples occur mostly as narrow biotite salvages within pink granite (Fig. 2a) and in the zone of intense deformation, we observe occurrence of post-metamorphic, K-metasomatic metapelites of the area (Fig.2b). However, the pre-metasomatic cordierite - bearing metapelites [8] which are preserved as relics within metasomatic rocks demonstrate that the metapelites of the Guwahati area had witnessed multiple deformations and related metamorphism prior to K-metasomatism. Our structural studies show that strongly flattened  $S_2$  foliation, axial planar to tight to isoclinal earlier folds and subsequent superposed folding, and post-kinematic shearing, fracturing and minor faulting.

### 3. PETROGRAPHY

Microscopic study of relics of pre-metasomatic metapelites indicate that they are strongly deformed, flattened and characterized by a well-developed foliation defined by the alignment of sillimanite, biotite, the long axis of plagioclase and quartz that swerve around the pre- $S_2$  porphyroblasts of cordierite. The porphyroblasts of cordierite contain tiny inclusions of sillimanite, biotite, and quartz, which represent the earliest (syn- $S_1$ ) metamorphic assemblage [4,5,8]. The metasomatic metapelites, the major concern of the present study, show inadequate amount of plagioclase, cordierite, sillimanite and several textures of relevant metasomatic reactions for the consumption of these phases. Some of these, for example, are (i) symplectitic intergrowth of muscovite + chlorite within the decomposed core of cordierite (Fig.3a), (ii) pseudomorphic replacement of cordierite by K-feldspar + biotite + chlorite (Fig.3b), (iii) development of thick rim of K-feldspar + biotite around cordierite (Fig.3c), (iv) stabilization of thin rim of K-feldspar at the contact between sillimanite and quartz (Fig.3d) and (v) presence of tiny inclusions of sillimanite within K-feldspar (Fig.3e) are common. The first appearance of metasomatic K-feldspar in the present rocks is marked by penetration of K-feldspar along the fracture plane of deformed plagioclase grains ( $An_{26}$ ) in (Fig.3f). K-feldspar tends to replace adjacent plagioclase crystals along grain boundaries in some of the thin sections. The relics of plagioclase grains within K-feldspar are sub-rounded and irregular and display myrmekite with tiny quartz vermicules (Fig.3f) although myrmekites in the present rocks are not common. The only clue for the formation of large-scale metasomatic K-feldspar in the present rocks is the presence of relics of metamorphic plagioclase and quartz which are preserved within metasomatic K-feldspar (Fig.3f). This feature is common in almost all thin sections.





Fig.3 (a) Symplectitic intergrowth of muscovite + chlorite within the decomposed core of finitized cordierite. (b) Pseudomorph replacement of cordierite by K-feldspar + biotite + chlorite. Note that biotite alters into chlorite. (c) Inclusion of altered cordierite grains within the thick rim of K-feldspar + biotite indicates the replacement of cordierite due to the K-activity of the prevailing fluid phase. (d) Formation of the thin rim of K-feldspar at the equilibrium contact of sillimanite and quartz. Note the paucity of matrix sillimanite in another part of the thin section. (e) Presence of tiny inclusions of sillimanite within K-feldspar indicates a paucity of matrix sillimanite. (f) Penetration of k-feldspar along the fracture plane of deformed metamorphic plagioclase grain with albite - twin. Note the presence of inclusions of sub-

rounded and irregular altered grains of plagioclase and quartz within K-feldspar but the grain boundary between plagioclase and quartz is separated by replacing K-feldspar. In some part (lower middle), altered grain of plagioclase displays very small myrmekite with tiny quartz vermicules.

### 3.1 The sequence of metasomatic reactions

The post-metamorphic, K-metasomatic mineralogical modifications involve two sets of reactions: (i) alteration of high-temperature all kinds of aluminium- silicate phases of pre-metasomatic metapelites and (ii) stabilization of K-feldspar at the expense of plagioclase and quartz in the rocks.

Among the first set of reactions, the conversion of cordierite porphyroblasts into muscovite and chlorite or K-feldspar + biotite ± chlorite or K-feldspar + biotite (Fig.3a-c) is significant. This low-temperature assemblage may be accounted for by following metasomatic reactions.



Reaction (i) is a low-temperature alteration of biotite and pinitization of cordierite forming chlorite and muscovite through hydration by a K<sup>+</sup>-bearing fluids [24], and is most commonly observed in the present rocks. Reactions (ii) and (iii) show that cordierite becomes unstable and is replaced by K-feldspar at higher  $\mu_{\text{K}_2\text{O}}$  in fluids. The reaction of cordierite alteration into biotite and K-feldspar as shown in (ii) and (iii) has been experimentally determined which show that even relatively at a low concentration of the K -component in the fluids, it is sufficient to produce biotite after cordierite [25].

The widespread post-metamorphic metasomatic alteration of sillimanite and stabilization of K-feldspar can be represented by the following reaction (Fig.3d):



This reaction greatly reduces abundance of sillimanite in the metasomatic rocks. The coexistence of sillimanite and quartz has been observed in many high-grade rocks but at higher  $a_{\text{K}_2\text{O}}$ , in fluids, sillimanite becomes unstable and is replaced by K-feldspar [26,27]. This also accounts for the reaction texture such as tiny inclusions of sillimanite within K-feldspar (Fig.3e).

The replacement of metamorphic plagioclase by K-feldspar is a major feature of K- metasomatism which can be accomplished by several processes [28,29]. However, in the present case, replacement of coexisting metamorphic plagioclase and quartz by K-feldspar in presence of K -rich fluids is the most dominant process of K-metasomatism. The relevant reaction is as follows.



It is important to mention that the above reaction is reversed of the myrmekite- forming reaction of Becke (1908) [30]. In fact, myrmekites which are the salient feature of many metasomatic rocks are not common in the present metasomatic metapelites, although the presence of tiny quartz vermicules indicates a rare occurrence of myrmekite (Fig.3f) in a few samples. In fact, in-situ replacement of plagioclase by metasomatic K-feldspar in the present rocks is minimum.

#### **4.CONCLUSION**

Our close field observation and petrographic studies demonstrate that the metapelites of Guwahati are overprinted by an intensive episode of post-metamorphic, K-metasomatism which has destroyed and obliterated all the pre-existing metamorphic textures including foliation of the original rocks. Several reaction textures indicate modifications of high-temperature minerals of pre-metasomatic metapelites into low-grade assemblage and replacement of metamorphic plagioclase and quartz by K-feldspar in presence of K-rich fluids. The newly formed metasomatic minerals thus include K-feldspar, albite, quartz, myrmekite, muscovite, chlorite and apatite and all these newly formed minerals together with pre-existing unreplaced metamorphic plagioclase have consequently produced K-feldspar -rich granite. In the zone of intense metasomatism, the existence of post-kinematic shear zones and minor faults have formed a fracture network for the movement of fluids. The K-rich metasomatic fluids, derived from deep-seated sources probably from the mantle moved into shear zones and migrated along the foliation plane, grain boundaries and microfractures in minerals, transforming the parent metapelites into metasomatic rocks. During the progressive metasomatism, all the shear zones and fractures have been replaced by K-feldspar due to prevailing K-rich fluids (Fig. 2b). and this indicates that the event of K-metasomatism post-dated all deformational and metamorphic events.

#### **REFERENCES**

1. Gogoi K. The geology of parts of the Khasi Hills along the southwestern and southeastern extension of the Sonapahar and structure of the main Sonapahar belt, Assam; Unpublished report of Geol. Surv. Intl. Prog. 1963.
2. Lal, R.K., Ackerman, D., Seifert, F., Haldar, S.K. Chemo-graphic relationships in sapphirine-bearing rocks from Sonapahar, Assam, India. *Contrib. Miner. Petrol.* 1978; 67: 169–187.
3. Maswood, Md. and Goswami, D.N.D. The metapelites of Guwahati, Assam. *Mines, Metals and Fuels*, 1978; v.26, pp.360-363.
4. Chatterjee, N.; Mazumdar, A. C.; Bhattacharya, A.; and Saikia, R. R. Mesoproterozoic granulites of the Shillong-Meghalaya Plateau: evidence of westward continuation of the Prydz Bay Pan-African suture into Northeastern India. *Precambrian Res.* 2007; 152:1–26.
5. Chatterjee N, Bhattacharya A, Duarah B P and Mazumdar A C Late Cambrian reworking of Palaeo-Mesoproterozoic granulites in Shillong–Meghalaya Gneissic Complex (Northeast

- India): Evidence from PT pseudosection analysis and monazite chronology and implications for East Gondwana assembly; *J. Geol.* 2011; 119 311–330.
6. Bhagabaty, B., & Mazumdar, A. (1). Petrology of granulites of the Shillong Plateau from west Garo Hills district, Meghalaya, India. *Journal of Nepal Geological Society*, 2007; V. 36 (Sp. Issue) p.7
  7. Rabha, M.; and Mazumdar, A.C. Petrology of the Umpyrtha- Patharkhammah metapelites, Meghalaya in North east India. *J. Assam Sc. Soc.* 2015; 56: 61-74.
  8. Borah, P., Hazarika, P., Mazumdar, A.C. *et al.* Monazite and xenotime U–Th–Pb total ages from basement rocks of the (central) Shillong–Meghalaya Gneissic Complex, Northeast India. *J Earth Syst Sci.* 2019; 128, 68.
  9. Dwivedi S B, Theunuo K and Kumar R R Characterization and metamorphic evolution of Mesoproterozoic granulites from Sonapahar (Meghalaya), NE India, using EPMA monazite dating; *Geol.* 2020; *Mag.* 157(9) 1409–1427.
  10. Susobhan Neogi & Tapan pal, metasomatically controlled sillimanite–corundum deposit: A case study from Sonapahar, Meghalaya, Northeast India, *J Earth Syst Sci.* 2021; volume 130, Article number: 120
  11. Ghosh, S., Fallick, A.E., Paul, D.K., Potts, P.J. Geochemistry and origin of Neoproterozoic granitoids of Meghalaya, Northeast India: Implications for linkage with amalgamation of Gondwana Supercontinent. *Gond. Res.* 2005; 8, 421–432.
  12. Kumar S, Rino V, Hayasaka Y, Kimura K, Raju S, Terada K and Pathak M. Contribution of Columbia and Gondwana supercontinent assembly and growth-related magmatism in the evolution of the Meghalaya Plateau and the Mikir Hills, Northeast India: Constraints from U–Pb SHRIMP zircon geochronology and geochemistry; *Lithos* . 2017; 277 356–377.
  13. Yin, A.; Dubey, C. S.; Webb, A. A. G.; Kelty, T. K.; Grove, M.; Gehrels, G. E.; and Burgess, W. P. Geologic correlation of the Himalayan orogen and Indian craton. 1. Structural geology, U–Pb zircon geochronology, and tectonic evolution of the Shillong Plateau and its neighboring regions in NE India. *Geol. Soc. Am. Bull.* 2010; 122:336–359.
  14. Crawford, A.R., India, Ceylon and Pakistan: new age data in comparison with Australia. *Nature*. 1969; 233, 380–382.
  15. Chimote, J. S.; Pandey, B. K.; Bagchi, A. K.; Basu, A. N.; Gupta, J. N.; and Saraswat, A. C. Rb–Sr whole rock isochron age from the Myllem granite, Khasi Hills, Meghalaya. 4th National Symposium of Mass Spectrometry, Banagalore. 1988.
  16. Ghosh, S., Chakraborty, S., Bhalla, J.K., Paul, D.K., Sarkar, A., Bishui, P.K., Gupta, S.N. Geochronology and geochemistry of gran-ite plutons from East Khasi Hills, Meghalaya. *J. Geol. Soc. India.* 1991; 37, 331–342.
  17. Ghosh, S., Chakraborty, S., Paul, D.K., Bhalla, J.K., Bishui, P.K., Gupta, S.N. New Rb–Sr isotopic ages and geochemistry of granitoids from Meghalaya and their significance in middle to late Proterozoic crustal evolution. *Ind. Miner.* 1994; 48, 33–44.
  18. Crawford, A.R. Indo-Antarctica, Gondwana Land and pattern of the distortion of a granulite belt. *Tectonophysics.* 1974; 22, 141–157.

19. Mazumdar, S.K. A summary of the Precambrian Geology of Khasi Hills, Meghalaya. *Geol. Ind. Misc.* 1976; Publ. 23 (2), 311–334.
20. Bardoloi, A., Mazumdar, A.C., Chowdhury, P.K. Polyphase deformation and fold geometry study of the Precambrian gneissic complex around Nalanga Hills, Goalpara district, Assam. *Ind. J. Earth Sci.* 1994; 21, 236–246.
21. Mazumdar, A.C.; Bardoloi, A.; and Chowdhary, P.K. Structural style of the gneissic rocks in and around Umananda river island, Guwahati, Assam *Indian Min.* 1997; 51:193-198.
22. Bhattacharjee, C. C., and Baruah, N. C. The study of joints in relation to the folding episodes in the Pre-Cambrian rocks of the Hahim area. *J. Assam Sci. Soc.* 1973; 16:54–62.
23. Powell, C.McA., Pisarevsky, S.A. Late Neoproterozoic assembly of East Gondwana. *Geology* 30, 2002; 3–6.
24. Seifert, F., Schreyer, W. Lower temperature stability limit of Mg cordierite in the range 1–7 kb water pressure: a redetermination. *Contrib. Miner. Petrol.* 1970; 27, 225–238.
25. Safonov, O.G. The role of alkalis in the formation of coronitic textures in metamangerites and metaanorthosites from the Adirondack Complex, United States. *Petrology.* 1998; 6, 583e602.
26. Korzhinskii, D.S. The role of alkalinity in the formation of charnockitic gneisses. *Trudy Vostochno-Sibirskogo Instituta Akademii Nauk SSSR Series of Geology.* 1962; 5, 50e61.
27. Korikovskiy, S.P., Kislyakova, N.G. Reaction textures and phase relations in hypersthene-sillimanite crystalline schists of the Sutam Complex of the Aldan Shield. In: *Metasomatism and ore formation (Metasomatity i orudnenie)*. Nauka, Moscow.1975; pp. 314e342.
28. Kehelpannala, K.V.M. the mechanism of post-metamorphic metasomatism of orthogneisses from Ambagaspitiya, Sri Lanka, *Journal of Geological Society of Sri Lanka.* *Journal of Geological Society of Sri Lanka.* 1998; 7, 23-36.
29. Kehelpannala, K.V.M., Ratnayake, N.P. Evidence for post metamorphic metasomatism of high-grade orthogneisses from Sri-Lanka. *Gondwana Research.*1998; 2, 167-184.
30. Becke, F. *Über Myrmekite.* *Schweizerische Mine-ralogischer und petrographischer Mitteilungen.* 1908; v. 27, pp. 337-390