

Original Research Article

PHYSICO-CHEMICAL AND MICROBIOLOGICAL EVALUATION OF SOIL AND EFFLUENT CONTAMINATED WITH AZO-DYE IN ITOKU, ABEOKUTA OGUN STATE, NIGERIA

ABSTRACT

The global industrialization of chemical dyes used for household products has resulted in various dyes leaching into the soil. This study was aimed at investigating the ecological effect of azo textile dye contamination on the bacterial biomass and its selective pressure, and also to determine the associated physicochemical changes in the soil samples contaminated with textile azo dye in Itoku, Abeokuta, Nigeria. The microbiological and physicochemical changes of the soil and effluent samples collected were determined using standard procedures. The results revealed that the average count of total viable bacteria in the contaminated sampling sites ranged from $0.82 \pm 0.021 \times 10^6$ cfu/g to $1.65 \pm 0.02 \times 10^6$ cfu/g which was significantly lower than the control bacterial counts. The results showed the azo dye contaminated and the control soil samples each contained a heterogeneous population of bacteria which included but with less density in the azo dye contaminated soil, the bacteria isolated include *Paenibacillus validus*, *Bacillus licheniformis*, *Bacillus niacin*, *Serratia liquefaciens*, *Staphylococcus gallinarum*, *Bacillus subtilis*, and *Bacillus coagulans*. The physicochemical analysis of the soil revealed high levels of pH in the azo dye contaminated soil (10.4–11.1) and. The electrical conductivity was also highest in the effluent with a range of (1250 to 2943 units) compared to the control and a control with a range of (430 to 480), which is the least. Importantly, the control samples were higher in all of the other tested soil parameters including cation exchange capacity in the control is significantly higher than other samples. The control also has the highest mean value of organic matter, at 4.13%. The bulk density is almost the same across the groups with an average value ranging between 1.59–1.65 g/cm³, while the other parameters like water holding capacity, particle density, moisture content, and total porosity. Conclusion-Azo dye contamination caused decreases in bacterial density and many other soil parameters along with higher PH and electrical conductivity which suggests that have the highest value in the control sample. There is a decrease in bacteria count and the soil parameters of the azo textile dye contaminated soil is, which thus makes it unfavorable for agricultural purposes and potentially toxic to the ecosystem.

Keywords: Azo dye, Contamination, Degradation, Heterogeneous, Physicochemical

SIGNIFICANCE STATEMENT: This research is significant as pollution of our ecosystem is increasing globally due to industrialization of chemical dyes which are in turn leaching into the soil. The particular aim significance of this study was to aimed at investigating the ecological effect of azo textile dye contamination in the soil on the bacterial biomass and its selective pressure and to determine the associated physicochemical changes in the soil samples contaminated with textile azo dye in Itoku, Abeokuta. The current study found that AZO dye contaminated soil altered the PH of the soil, significantly reduced bacterial abundance in the soil, and lowered many soil parameters. This data suggests that AZO

Comment [j1]: Dear Author, this is a very interesting study and generally well written the Abstract however needed rewriting and the significance statement- but I only made small changes to the body of the paper which was well written in most spots. Feel free to use any changes that I have made that you like. You can see all my changes by going to Review in Microsoft word and clicking on all markup. Goodluck with the publication of your paper. Best wishes Lucinda Choules

Comment [j2]: Dear Author it is important to grab the readers attention with the first line by stating the importance or reason for your study so please add this line or one similar to grab the readers attention as Abstracts have to stand alone and have to interest the reader so people will be compelled to read your article.

Comment [j3]: Good Title

Comment [j4]: Possible changes "Azo Textile Dye contamination could lead to Infertile Soil"

Formatted: Font: Italic

Comment [j5]: Should "significant" be here?

Comment [j6]: This should be different then the Abstract perhaps write about how the soil becomes polluted by this dye in Nigeria and why your findings are important as they show that this dye is a problem for the environment.

dye in the soil results in adverse effects to the soil that could eventually lead to infertile soil. microbiological and physicochemical changes of soil and effluent samples collected were determined using standard procedures. The results showed the azo dye contaminated and the control soil sample contained a heterogeneous population of bacteria but with less density in the azo dye contaminated soil, the bacteria isolated include: *Paenibacillus validus*, *Bacillus licheniformis*, *Bacillus niacin*, *Serratia liquofaciens*, *Staphylococcus gallinarum*, *Bacillus subtilis*, and *Bacillus coagulans*. It was observed that the pollution of dye to the surrounding soil had deleterious effect on the physicochemical properties of the soil.

INTRODUCTION

Environmental pollution of the ecosystem is becoming worrisome due to increased industrialization of the environmentally unfriendly chemical (dyes) that are produced and consumed, affecting both living and nonliving components{1}. Synthetic dyes are of great importance in textile dyeing, paper printing, food, pharmaceuticals, cosmetics, photography, paint, petroleum products, and other industrial uses. Azo dye, Azo dye is being an artificial dye, has been used in various aspects of human endeavor with the primary aim of adding color to materials {2}. This is mostly employed in tie and dye of clothing, which is common in some parts of the country. Over the years, the use of azo dye has been increasing, which and this constantly raises concern as it has been linked to many disease conditions {3}. Dyes and pigments are designed to resist degradation, such that they remain in the environment for a long period of time. For example, the half-life of the hydrolyzed dye Reactive Blue 19 is about 46 years at pH 7 and 25°C {4}.

Azo dye is well known to cause damage to living cells via by its mutagenic and carcinogenic potential that can, by affecting the genome of living cells, it also affects the ecosystem and thus puts humanity at risk {5}. There is a continuous search for the best and most environmentally friendly method of cleaning, however, also partially degraded dyes are more toxic than the mother molecule{6,7}. This toxic potential of

the dye creates a selective pressure on the bacteria population and causes adaptation for survival{8}. The bacterial population and type present in a soil determine the ecological state and, subsequently, the fertility of the soil. Some bacteria are well adapted to live ~~an in the~~ environment impregnated with dye, ~~and while~~ some are even adapted to ~~assist with~~ degradation of the dye{9}. ~~However, it~~ has been reported that many of the azo dyes and aromatic amines produce potent toxins during their degradation and they contain heavy metals and carcinogens which may have a deleterious negative impact on bacterial cells and ecology {10}.

Conventional physical methods of treatment only transfer the pollutants from one form to another and produce secondary waste products. Examples include physical or chemical flocculation; membrane filtration; electro kinetic coagulation; electro chemical destruction; and precipitation{11}. These types of technologies are inefficient, expensive, and have low adaptability to diverse types of dye effluent. Dyes escape conventional wastewater treatment processes and persist in the environment as a result of their high stability to light, temperature, water, detergents, soap and other parameters{12}. The use of microorganisms for biodegradation of dye is convenient and versatile, with dynamic metabolisms and potential machinery of enzymes. Biodegradation is a nonhazardous, cost-efficient, environmentally friendly, and often more effective alternative to conventional methods of treatment of textile dye waste {13}.

The physico-chemical character of a soil contaminated with dye is affected. This ranges from the PH, conductivity, CEC, organic and non-organic matter, and soil structure{14}. Damage to soil property and structure can encourage leaching, thus loss of fertility. In another view, the survival of certain bacteria is dependent on the structural and chemical state of the soil, damage to ~~the~~ structural state of soil will affect the range of useful bacteria present. Dye has ~~being~~ linked to increased alkalinity and loss of fertility of soil, ~~that could result in -with-~~ long-term consequences ~~such as ef-~~ loss of substantial hectares of land, ~~especially down the stream~~ for agricultural purposes, and also damage to aquatic life. Research has shown that there is a decrease in the physicochemical properties of soil irrigated with industrial dye with a corresponding impact on crop yield{15,16}.

The search for a biological method of control of dye pollution is a necessity. This could be achieved through bioremediation or biodegradation as other methods of decontamination of dye are expensive and non-effective{17}. Bacteria that can survive in dye-contaminated soil may have the potential to degrade dye through mutation and selective pressure of the environmental toxicant{18}. The bacteria type and population are transitory as there is high turnover of mutation and development of new and resistant species{19}. There is limited literature on characterization and identification of bacteria biomass in a dye contaminated ecosystem. Thus, there is a need to isolate, characterize, and identify bacteria present in dye contaminated soil as well as the physicochemical properties of the soil which will secondarily affect the bacterial population.

EXPERIMENTAL DETAILS

Study Area

The study was carried out in Olumo rock area, Itoku, Abeokuta, Ogun state.




Figure 1:Map showing study area & sample locations

Sample Collection

Soil samples of 200g ~~were~~ collected from three different dye contaminated sites in Itoku, Abeokuta, from the top-soil layer profile of 0 - 20cm; in a sterile plastic bag and stored in ambient temperature and transported to the laboratory for analyses.

Physical parameters of soil samples and Chemical parameters of soil samples

The parameters analyzed included pH by potentiometric meter. Moisture content, bulk density and porosity. The chemical parameters included Electrical Conductivity (EC), Cation Exchange Capacity (CEC), a sample weight of 20 g was air dried and treated with 100 cm³ of 0.1 mol dm⁻³ Hydrochloric acid solution (in the ratio w: v of 1:5). The suspension ~~is~~ will be shaken in a polyethylene bottle of about 250 cm capacity on a rotary shaker (40 rpm) at room temperature for 1hour, and then filtered. The filtrates will be titrated with 0.1 mol dm⁻³ Sodium hydroxide solution in presence of phenolphthalein indicator and the sum of exchangeable alkaline cations will be calculated from the amount of sodium hydroxide solution used {20,21}.

Isolation and Characterization of Isolates Microorganisms

Isolation and enumeration of total viable bacteria will be carried out using Nutrient agar by pour plate methods. Colonies will be sub-cultured on freshly prepared Nutrient agar and the plates will be incubated at 37 °C for 24 h {22}

Biochemical identification of isolates

Phenotypic and biochemical identification of bacterial isolates will be conducted with the following tests ~~done by carrying out~~ Gram staining ~~reactions~~, catalase test, coagulase test, oxidase test, spore test, motility test, and growth on differential media, citrate utilization, Methyl Red, Voges Proskauer reactions, urease production, nitrate reduction, and sugar fermentation tests using Advance bacteria identification software 2021 version.

Statistical analysis: The data obtained in this study were evaluated using descriptive statistics through simple graphs, tables & charts and the one-way analysis of variance (ANOVA) followed by Dunnett's post hoc test, using Graphpad prism version 8.0 software (La Jolla, California, USA). The level of significance was set at $\alpha_{0.05}$.

RESULTS AND DISCUSSION

Tables 1A and 1B show the results of the bacterial isolation enumeration. The bacteria had the highest count in control soil ($4.06 \pm 0.09 \times 10^6$ cfu/g) compared to the textile dye-polluted soil ($1.23 - 1.65 \pm 0.02 \times 10^6$ cfu/g), while that of the effluent ranged from ($0.61 - 0.98 \pm 0.02 \times 10^6$ cfu/ml). These statistics were based on the average count of all viable bacteria and the standard deviation of colonies that were isolated. The low bacterial counts in the soil that was tainted with textile dye suggest that soil may contain some compounds that prevents high survival of heterotrophic bacteria. The study by {23} showed the total viable counts of bacteria from textile dye-contaminated soils were found to be comparable{23}.

Comment [J7]: Did you mean heterotrophic here?

Table 1 A: Colony count from soil Dye samples

Sample codes	TVBC ($\times 10^6$ cfu/g)
DS1	1.65 ± 0.02
DS2	0.82 ± 0.21
DS3	1.23 ± 0.11
DCS	4.06 ± 0.09

All values in mean \pm SEM, Total Viable Bacterial Counts (TVBC),

Dye Soil (DS1-3), Dye Control Soil (SC)

Table 1B: Colony count from Dye effluent samples

Sample codes	TVBC (x10 ⁶ cfu/ml)
DE1	0.98±0.05
DE2	0.63±0.04
DE3	0.61±0.05

All values in mean±SEM, Total Viable Bacterial Counts (TVBC),

Dye Effluent (DE1-3)

A total of seven bacterial isolates were found in the results using advance bacteria identification software (2021) version. Frequencies and percentages were used to identify the isolated bacteria using advanced biochemical procedures. *Bacillus niacin* had the highest frequency and percentages of bacterial species in this study, with 6 (28.57%), followed by *Bacillus coagulans* 4 (19.05%), *Paenibacillus validus* 3 (14.29%) and *Bacillus licheniformis*, *Serratia liquefaciens*, *Staphylococcus gallinarum*, and *Bacillus subtilis*, with 2 each (9.52%) It may not be surprising that there were many different species of *Bacillus* found in the polluted soil given that these organisms are native to soil environments and are known to persist there {24}. The isolated organisms from dye-contaminated soil are able to break down both the organic and inorganic dye constituents. Despite the dye's poisonous and resistant constituents, bacteria continue to exist by adapting to their surroundings. Similar research has suggested that members of the *Bacillus* genus are capable of decolorizing azo textile dye. Research has showed that *Bacillus subtilis*, and *B. cereus*, can decolorize azo dyes(25).

Table 2: Cultural and biochemical characteristics of Bacterial isolated from Soil and Effluent samples

<i>Paenibacillus validus</i>	<i>Bacillus licheniformis</i>	<i>Bacillus niacin</i>	<i>Serratia liquefaciens</i>	<i>Staphylococcus gallinarum</i>	<i>Bacillus subtilis</i>	<i>Bacillus coagulans</i>
similarity	s similarity	similarit	ns	gallinarum	similarit	similarity
96.5%	87.8%	y 90.8%	similarity	similarity	y 91.2%	95.5%

93.4% 80.2%

Gram stain	+	+	+	-	+	+	+
Spore	+	+	+	-	-	+	+
Catalase	+	+	+	+	+	+	+
Glycerol	+	+	+	+	+	+	+
H ₂ S	-	-	-	+	-	-	-
Lactose	-	-	-	-	-	-	-
Sucrose	+	+	+	+	-	-	+
Glucose	+	+	+	+	-	+	+
MR	+	+	-	+	+	+	-
V.P	-	-	-	+	+	+	-
Starch hydrolysis	+	-	+	-	-	+	+
Casein hydrolysis	-	-	-	-	-	+	-
Lipase	-	+	+	+	+	+	+
Nitrate	+	+	+	+	+	+	+
Motility	+	+	-	+	+	+	+
Indole	-	-	-	-	-	-	-
Citrate	-	+	+	+	-	-	-
Arabinose	-	+	-	+	+	+	+
Xylose	+	+	+	+	+	+	+
maltose	+	+	+	+	+	+	+

mannitol	+	+	-	+	+	+	-
Erucrose	+	+	+	+	+	+	+
Raffinose	-	+	-	+	+	-	+
sorbitol	+	+	-	+	+	+	+
galactose	+	+	+	+	+	+	+
Ribose	+	+	+	+	+	+	+
rhamnos	-	+	+	-	+	-	+
e							

UNDER PEER REVIEW

Table 3: Occurrence of bacterial isolates in the soil and effluent samples

Samples	<i>Paenibacillus validus</i>	<i>Bacillus licheniformis</i>	<i>Bacillus niacin</i>	<i>Serratia liquefaciens</i>	<i>Staphylococcus gallinarum</i>	<i>Bacillus subtilis</i>	<i>Bacillus coagulans</i>
Dye Soil 1	+	+	+	-	-	-	-
Dye Soil 2	-	-	+	+	-	-	-
Dye Soil 3	+	-	-	+	+	-	-
Control	-	-	-	+	+	-	-
Dye effluent 1	-	-	+	-	-	+	+
Dye effluent 2	-	+	+	-	-	+	-
Dye effluent 3	+	-	+	-	+	-	+

Keys: + = present - = absent




Fig. 2: Frequency and percentage occurrence of bacteria species isolated from soil contaminated with dye effluenteffluents.

Alkalinity is a measurement of a soil's buffering capacity. It is a critical indicator of how effectively soil can scavenge acids from effluent wastes. At higher alkalinity values, bicarbonates and carbonates are more common in effluents. The azo textile dye-contaminated soil during this study exhibited higher pH levels (11.1–11.5) in every sampling site, exceeding the tolerated levels {23}. The findings showed that the soil moved toward an alkaline state and that it was over the allowable limits of (6-9). The pH in the

effluent is moving toward a higher value, indicating alkalinity conditions, and this could negatively affect the permeability of the soil and the development of the soil microbial flora. Increased carbonates and bicarbonates from textile dye effluents may potentially be a contributing factor to high alkalinity. This is consistent with {23} findings, which revealed a high pH level (9.36- 9.44).

UNDER PEER REVIEW

Table 4: Physicochemical parameters of the dye contaminated soil samples

Samples	Water holding capacity (%)	Bulk density (g/cm)	Particle density (g/cm)	Moisture content (%)	Total porous (%)
DS 1	24.44±0.34	1.59±0.05	2.26±0.18	8.57±0.02	26.59±0.28
DS 2	19.26±0.33	1.65±0.03	2.21±0.01	8.39±0.04	22.19±0.22
DS 3	17.27±0.55	1.61±0.03	2.27±0.02	9.29±0.07	21.11±0.23
DCS	28.40±0.36	1.60±0.03	3.24±0.03	9.82±0.08	32.68±0.12

Dye soil (DS 1-3), Dye control soil (DCS)

Table 5A: Chemical parameters of the dye contaminated soil samples

Samples	Ph	Electrical conductivity (µS/cm)	CEC (Mol/Kg)	Organic matter (%)
DS 1	11.06±0.01	3273±0.03	0.25±0.00	1.31±0.02
DS 2	10.38±0.03	1250±0.03	0.25±0.01	2.44±0.05
DS 3	10.43±0.03	2943±0.02	0.14±0.02	1.89±0.03

Dye soil (DS 1-3), Dye control soil (DCS)

Table 5B: Physicochemical parameters of the dye contaminated effluent [samplesamples](#).

Samples	Ph	Electrical conductivity ($\mu\text{S/cm}$)	CEC (Mol/Kg)	Organic matter (%)
DE 1	12.90 \pm 0.03	9267 \pm 0.03	0.16 \pm 0.01	3.10 \pm 0.02
DE 2	13.31 \pm 0.02	8769 \pm 0.03	0.17 \pm 0.01	2.86 \pm 0.02
DE 3	13.56 \pm 0.05	9511 \pm 0.03	0.05 \pm 0.042	2.58 \pm 0,03

Dye effluent (DE 1-3), Cation exchange capacity (CEC)

The ability of the soil to retain positively charged ions known as cations is known as its cation exchange capacity (CEC). Electrostatic forces cause the negatively charged clay and organic matter particles in the soil to hold onto these cations (negative soil particles attract the positive cations). According to the results above, the control soil sample's cation level was greatest, ranging between 0.4 and 0.5, while the levels in the soil polluted with textile dye and the textile dye effluents were between 0.1 and 0.2. Because they can be readily exchanged with other cations, the cations on the CEC of soil particles are available to plants. More clay or organic matter is present in the soil the greater the CEC. This often means that clay soils with high CEC have a higher water retention capacity than soils with low CEC (sandy soils). The organic matter norm ranges from 2.00 to 6.00. The organic matter content ranges between 1-3, which is extremely low for the survival of soil organisms and the maintenance of plants {26}

Cations are positively charged ions, the capacity of the soil to hold on to these cations is called the cation exchange capacity (CEC). These cations are held by the negatively charged clay and organic

matter particles in the soil through electrostatic forces (negative soil particles attract the positive cations). From the result above result the cation from the control soil sample had the highest which ranged between 0.4-0.5 while the textile dye contaminated soil and the textile dye effluents ranged between 0.1-0.2. The cations on the CEC of the soil particles are easily exchangeable with other cations and as a result, they are plant available. The higher the CEC the more clay or organic matter present in the soil. This usually means that high CEC (clay) soils have a greater water holding capacity than low CEC (sandy) soils.

An organic matter concentration of more than 8.00 percent results in an excessive availability of nutrients as well as other issues like high pH, high soluble salts, etc. Additionally, in both uncontaminated and contaminated soil samples, soils with an organic matter content of less than 3.00 percent are stated to be less productive.

CONCLUSION

There is biodiversity in the bacterial population and a high count in the normal soil sample compared to that of the dye contaminated soil. Most of the isolated bacteria are bacillus which has the potential to biodegrade azo dye. This could have been a result of environmental selective pressure or a mutation that allows for survival. Thus, However there may be a need to genetically screen the isolated bacteria to determine if it is a new strain of bacillus. The physicochemical properties of the contaminated soil were significantly affected by the dye and thereby resulting in poor soil quality, which could lead to loss of soil fertility.

REFERENCES

- [1.] Masindi V, Muedi KL. Environmental contamination by heavy metals. Heavy Met. 2018;10:115–32.
- [2.] Fried R, Oprea I, Fleck K, Rudroff F. Biogenic colourants in the textile industry--a promising and sustainable alternative to synthetic dyes. Green Chem. 2022;24(1):13–35.
- [3.] Bakthavachalu P, Kannan SM, Qoronfleh MW. Food color and autism: A meta-analysis. Pers Food

Interv Ther Autism Spectr Disord Manag. 2020;481–504.

- [4.] Awasthi A, Datta D. Application of Amberlite XAD-7HP resin impregnated with Aliquat 336 for the removal of Reactive Blue-13 dye: Batch and fixed-bed column studies. *J Environ Chem Eng.* 2019;7(6):103502.
- [5.] Karri RR, Ravindran G, Dehghani MH. Wastewater—sources, toxicity, and their consequences to human health. *Soft Comput Tech solid waste wastewater Manag.* 2021;3–33.
- [6.] Singha K, Pandit P, Maity S, Sharma SR. Harmful environmental effects for textile chemical dyeing practice. *Green Chem Sustain Text.* 2021;153–64.
- [7.] Hassaan MA, El Nemr A. Advanced oxidation processes for textile wastewater treatment. *Int J Photochem Photobiol.* 2017;2(3):85–93.
- [8.] Paul D, Chakraborty R, Mandal SM. Biocides and health-care agents are more than just antibiotics: Inducing cross to co-resistance in microbes. *Ecotoxicol Environ Saf.* 2019;174:601–10.
- [9.] Sreedharan V, Bhaskara Rao KV. Biodegradation of textile azo dyes. *Nanosci Biotechnol Environ Appl.* 2019;115–39.
- [10.] Khan S, Malik A. Environmental and health effects of textile industry wastewater. *Environ Deterior Hum Heal.* 2014;55–71.
- [11.] Sen S, Prajapati AK, Bannatwala A, Pal D. Electrocoagulation treatment of industrial wastewater including textile dyeing effluent--a review. *Desalin Water Treat.* 2019;161(1):21–34.
- [12.] Sirisha K, Suganya B, Sivasubramanian V, Bs V, Swaminathan D, Babu AC, et al. Studies on the effect of pulsed magnetic field on the productivity of algae grown in dye industry effluent. *J Appl Biotechnol Bioeng.* 2017;3(5):409–13.
- [13.] Jabbar NM, Alardhi SM, Mohammed AK, Salih IK, Albayati TM. Challenges in the implementation of bioremediation processes in petroleum-contaminated soils: A review. *Environ Nanotechnology, Monit & Manag.* 2022;18:100694.

- [14.] Luo Z, Yao B, Yang X, Wang L, Xu Z, Yan X, et al. Novel insights into the adsorption of organic contaminants by biochar: a review. *Chemosphere*. 2022;287:132113.
- [15.] Liu M, Wang C, Liu X, Lu Y, Wang Y. Saline-alkali soil applied with vermicompost and humic acid fertilizer improved macroaggregate microstructure to enhance salt leaching and inhibit nitrogen losses. *Appl Soil Ecol*. 2020;156:103705.
- [16.] Riza M, Ehsan MN, Hoque S. Portrayal of Textile Based Pollutants and its Impact on Soil, Plants and Fisheries. *Nat Environ Pollut Technol*. 2021;20(3):1269–75.
- [17.] Barik P, Biswal T, Malik JA. Assessment of Current Soil Bioremediation Technologies. *Bioremediation Phytoremediation Technol Sustain Soil Manag*. 2022;147–84.
- [18.] Tacas ACJ, Tsai P-W, Tayo LL, Hsueh C-C, Sun S-Y, Chen B-Y. Degradation and biotoxicity of azo dyes using indigenous bacteria-acclimated microbial fuel cells (MFCs). *Process Biochem*. 2021;102:59–71.
- [19.] Blanquart F. Evolutionary epidemiology models to predict the dynamics of antibiotic resistance. *Evol Appl*. 2019;12(3):365–83.
- [20.] Keesari T, Pant D, Roy A, Sinha UK, Jaryal A, Singh M, et al. Fluoride geochemistry and exposure risk through groundwater sources in northeastern parts of Rajasthan, India. *Arch Environ Contam Toxicol*. 2021;80(1):294–307.
- [21.] Jaremko D, Kalembasa D. A comparison of methods for the determination of cation exchange capacity of soils/Porównanie metod oznaczania pojemności wymiany kationów i sumy kationów wymiennych w glebach. *Ecol Chem Eng*. 2014;21(3):487.
- [22.] Asmamaw H, Yalemtehay M. Evaluation of the antibacterial activities of leaf extracts of *Achyranthus aspera*. *African J Bacteriol Res*. 2017;9(2):9–14.
- [23.] Sirajo M, Muhammad B, Yerima SA, Ladan BB. Microbiological and Physicochemical Assessment of Soil Contaminated with Dye Effluent in Sokoto State, Nigeria. *Int J Curr Microbiol App Sci*. 2018;7(6):3800–7.

- [24.] Raffa CM, Chiampo F. Bioremediation of agricultural soils polluted with pesticides: A review. *Bioengineering*. 2021;8(7):92.
- [25.] Noor NNM, Kamaruzaman NH, Al-Gheethi A, Mohamed RMSR, Hossain MS. Degradation of antibiotics in aquaculture wastewater by bio-nanoparticles: A critical review. *Ain Shams Eng J*. 2022;101981.
- [26.] Sheoran V, Sheoran AS, Poonia P. Soil reclamation of abandoned mine land by revegetation: a review. *Int J soil, sediment water*. 2010;3(2):13.

UNDER PEER REVIEW