

Original Research Article

Trypanosomiasis in rodents from selected plague endemic foci of Tanzania; the concealed public health threat.

Abstract

Trypanosomiasis is among the zoonotic diseases that cause threat to public health, rodents are known to be reservoir of various Trypanosomes which are zoonotic. The ~~cross-sectional~~ cross-sectional studies ~~was~~ were conducted in selected areas of plague endemic foci in Tanzania, aiming at establishing the prevalence of *Trypanosoma* sp. infection in rodent species. A total of 105 rodents comprising nine species were captured in different habitats during the study ~~period~~ conducted ~~period~~ conducted between March and May 2022. The prevalence recorded was 4.8%(5/105), with individual prevalence of *Mastomys natalensis* 1.9%(2/105), *Rattus rattus* 1.9%(2/105) and *Lophuromys kilonzo* 1.0%(1/105). Prevalence of *Trypanosoma* sp. infection was not differed significantly between host species ($P>0.05$), host sex ($P>0.05$), habitat type ($P>0.05$) and wards ($P>0.05$). The ~~present study~~ present study has shed some light on the presence of *Trypanosoma* sp. infection in rodents in plague endemic foci of Tanzania, hence raising the public health concern due to their zoonotic potential to humans.

Keywords: Prevalence, *Trypanosoma* sp., Rodents, Plague, Endemic area, Tanzania.

Formatted: Left

4-Introduction.

Rodents are small mammals that belongs to the order Rodentia, the largest order of mammals comprising more than 32 families, 481 genera and 2277 species [32]. Rodent species are widely distributed occurring almost in all continents and vastly diversified [13,19]. Rodents are highly adapted to various habitats and environments [30]. Regardless of showing greater diversity in their behaviors, morphology, ecology and physiology, all rodent species have the common dentition feature. Indeed, they are characterized by the single pair of continuously ever-grown incisors, that has to be maintained in a short length by a gnawing process [1,16]. Rodent species are known to harbor many different microorganisms that are pathogenic to animals and of public health concern, various Bacteria, Viruses, Rickettsia, Fungi, Arthropods, Helminths and Protozoans have been reported [3,27].

Trypanosomiasis is a ~~disease caused~~ disease caused by protozoans of the genus ~~Trypanosoma which~~ Trypanosoma which can infect both human beings and animals [39]. These parasites are transmitted by arthropods vectors such as Tsetse flies, Tabanus, Mites, Reduviid bugs and Fleas both mechanically and biologically [31,39]. *Trypanosoma lewisi*, *Trypanosoma evansi* and *Trypanosoma lewisi* like, have been reported to infect different types of rodent species in Europe, Asia, Australia, America and Africa [21,35,37,40]. Rodents usually gets infected by *Trypanosoma* spp. by ingesting the infected fleas or through ingestion of the feces of infected fleas [12]. Several fleas have been implicated to transmit *Trypanosoma* spp. in various countries and this species includes the members under genera *Xenopsylla*, *Ctenophthalmus*, *Nosopsyllus*, *Dinopsylla* to mention a few, which are also very prevalent in the plague endemic foci of Tanzania [12,23,26,31].

Despite of being infecting rodent species, these *Trypanosoma* species are zoonotic and may cause Atypical Human Trypanosomiasis, indeed, several cases have been reported with few cases resulting into mortalities [7,5,25,38,42]. Due to close interaction between the rodents and Humans there is a high possibility of Humans to get infected with pathogenic agents from the rodents such as *Trypanosoma* spp. The closeness

interaction between the rodents and humans especially in the rural areas could facilitate the possible transmission of zoonotic agents [8]. In addition, human activities such as agricultural activities increase the abundance of the rodents, in turns the risks of zoonotic infections are also increasing.

In ~~Tanzania, to Tanzania, to~~ the best of our knowledge there are few studies that have been done to ascertain the magnitude of rodent Trypanosomiasis [14,15,16,36]. Furthermore, most of these studies have been conducted in one region only which is Morogoro region neglecting other regions. Therefore, the proposed study was sought to determine the prevalence of *Trypanosoma* species infection in rodents from selected area of plague endemic foci of Tanzania, with a view of providing a baseline information on possibility of zoonotic infection in rodent species by *Trypanosoma* species in the area and necessitating the need of advocating the plausible control and/or prevention measures to the stakeholders.

2- Materials and Methods

2.1 Study area

The study was conducted in two districts within Tanga and Morogoro Regions namely Lushoto and Mbulu districts respectively which are plague endemic foci. Lushoto ~~district location~~ district location lies between 4°32'S and 38°37'E, with majority of the part lying within Western Usambara Mountains that forms a part of Eastern Arc Mountains. Mbulu district lies between 3°51'S and 35°32'E. These districts experienced plague outbreak in ~~last thirteen~~ last thirteen years [44]. Within each ~~district~~ two districts two wards were selected and within each ward two villages were ~~selected. In~~ selected. In Lushoto district, Shume (Mavumo and Gologolo villages) and Manolo (Madala and Manolo villages) wards were selected and sampled. In Mbulu district, Nahasey (Haysali and Nahasey villages) and YaedaAmpa (Hayaseng and Arri villages) wards were selected and sampled.

Formatted: Left

2.2 Study design

A cross-sectional study design ~~were~~was employed.

2.3 Rodents trapping, identification and blood collection

Rodents were captured in different habitats by Sherman traps which was baited by a mixture of maize flour and peanut butter, from March to May 2022. The habitats selected for sampling were as follows; fallow land, crop field, bushes, plantation forests, indoor and near natural forest areas. Sixty traps were placed on the habitat overnight while five traps were placed indoor ~~per—house~~per house, the number of days for trapping in each village was three days which is equal to six days of trapping per ward. Five households were selected randomly from the list provided by local administration for indoor rodents sampling, verbal consent was sought prior to trap ~~setting-The~~setting. The traps were set during the evening (1700hrs-1800hrs) and inspected the following day in the morning (0600-0700hrs). After carefully inspection the trapped rodents were carried to the nearby area where the rodents were taken out from the traps into clothing bag, thereafter, each individual was anaesthetized by soaking the clothing bag contain the rodent into a jar that contain 98% diethyl ether one by one[16,17].Thereafter, the individuals was identified to genus or specie level by the aid of keys[19] and the blood was collected via cardiac puncture by using 2mls syringes, then the blood was placed on the well labelled 5ml Sodium Ethylene Diamine Tetraacetate (EDTA) vacutainer tube for storage and transportation to Parasitology laboratory at the Department of Veterinary microbiology, Parasitology and Biotechnology, Sokoine University of Agriculture. Various parameters were recorded from the rodents to aid their identification including tail length, head body length, pes length, weight, ear length, sex and breeding status.

2.4 Smear preparation and identification of *Trypanosomes* sp.

From the stored EDTA vacutainers containing blood sample in the refrigerator, the thin and thick smears was prepared. Thin blood smear was prepared by placing a small drop of blood on a near short end of the clean slide. By using the spreader slide holding it at an angle of 35 degree, the slide was placed in front of the drop of blood then slowly

it was pulled back until it was touching the drop of blood, then the drop of blood was allowed to spread along the juncture of the two slides. Thereafter, the spreader slide was smoothly and rapidly pushed forward to produce a smear. The slide was allowed to air dry and then stained with 10% Giemsa stain for 30 minutes. The same procedure was done in preparing the thick smear except that, after air dry the smear was fixed in methyl alcohol for 10 minutes, then stained with 10% Giemsa stain for 30 minutes. The excess stain in both smears was removed by tap water, all the procedures were performed according to [12]. The smears were then observed under compound microscope by using 100X magnification with the aid of oil immersion.

2.5 Data analysis

The raw data was double entered and cleaned on Microsoft excel 2019. Descriptive data were summarized in tables and figures. The prevalence was calculated according to [2]. The prevalence of Trypanosomiasis was compared between host species, host sex, habitat type and wards by Chi square test. All the statistical analysis was computed on Statistical Package for the Social Sciences (SPSS) version 20 IBM. In all analysis the *P*-value was set at 0.05 as significant.

2.6 Ethical clearance

Ethical clearance was obtained from the Institutional Ethical Committee of the Sokoine University of Agriculture with reference number SUA/DPRTC/186/VOL IV.

3. Results

3.1 Rodents captured

A total of 105 rodent individuals were captured during the study period, with 9 species encountered including *Arvicanthisnairobe*, *Rattus rattus*, *Grammomys* sp., *Mus* sp., *Acomys* sp., *Lophuromyskilonzoi*, *Praomysdelectorum*, *Lophuromysflavopunctatus* and *Mastomysnatalensis* (Table 1). Their abundance is also shown in Table 1. The distribution of different rodent species in various habitats is shown in Table 2. The percentage of females and males captured is shown in figure 1.

Formatted: Left

R.rattus and *M.natalensis* dominated the capture with more than 70% of the total capture. The number of rodent individuals captured within different wards are shown in Table 3.


Table 1. Number of captured individual rodents during the whole study and their abundance.

Rodents species	Number of captured individuals	Relative abundance (%)
<i>Mastomysnatalensis</i>	62	56.0
<i>Rattus rattus</i>	16	15.2
<i>Luphorumysflavopunctatus</i>	6	5.7
<i>Luphorumyskilonzoi</i>	4	3.8
<i>Acomys</i> sp.	1	1.0
<i>Grammomys</i> sp.	3	2.9
<i>Praomysdelectorum</i>	5	4.8
<i>Mus</i> sp.	2	1.9
<i>Arvicanthisnairobe</i>	6	5.7
Total	105	100

Table 2. Distribution of different species of rodents according to various habitats on selected plague endemic areas.

Rodent species	Habitats					
	Fallow land	Crop field	Indoor	Bush	Plantation forest	Near natural forest
<i>Mastomysnatalensis</i>	x	x	x	x	x	
<i>Rattus rattus</i>			x			
<i>Luphorumysflavopunctatus</i>	x	x				x
<i>Luphorumyskilonzoi</i>		x			x	
<i>Acomys</i> sp.		x				
<i>Grammomys</i> sp.				x	x	
<i>Praomysdelectorum</i>		x		x	x	
<i>Mus</i> sp.		x	x			
<i>Arvicanthisnairobe</i>	x	x				

UNDER PEER REVIEW






Figure 1. Percentage of captured rodent species according to sex.

Table 3: Number of captured individual rodents during the study with respect to their sampling wards.

Rodent species	Wards				Total
	Manolo	Nahasey	Shume	YaedaAmpa	
<i>Mastomysnatalensis</i>	14	15	21	12	62
<i>Rattus rattus</i>	0	4	7	5	16
<i>Luphorumysflavopunctatus</i>	0	2	0	4	6
<i>Luphorumyskilonzoi</i>	1	0	3	0	4
<i>Acomys</i> sp.	0	1	0	0	1
<i>Grammomys</i> sp.	1	0	2	0	3
<i>Praomysdelectorum</i>	0	0	5	0	5
<i>Mus</i> sp.	0	0	1	1	2
<i>Arvicanthisnairobe</i>	4	2	0	0	6
Total	20	24	39	22	105

3.2-Prevalence of Trypanosomiasis

Prevalence of Trypanosomiasis was found to be 4.8%, 95% CI(1.6, 10.8), with two individuals from *R. rattus* (1.9%) and *M. natalensis* (1.9%) infected while one *L. kilonzo* (1.0%) was infected. The rest of the species was found to be not infected by *Trypanosomes* sp. The Chi square test revealed that the *Trypanosoma* sp. infection prevalence was not statistically different when compared to host species ($X^2=7.198$, $df=8$, $P=0.515$), host sex ($X^2=0.008$, $df=1$, $P=0.930$), habitat type ($X^2=2.965$, $df=5$, $P=0.705$) and wards ($X^2=3.837$, $df=3$, $P=0.280$) as shown in Table 4 and Table 5.

Table 4.Prevalence of *Trypanosoma* sp. in rodents regarding to host species and host sex.

Variable	Sample size	Infected no.(%)	Chi-square	P-value
Host species				
<i>Mastomysnatalensis</i>	62	2(1.9)		
<i>Rattus rattus</i>	16	2(1.9)		
<i>Luphorumysflavopunctatus</i>	6	0(0.0)		
<i>Luphorumyskilonzoi</i>	4	1(1.0)		
<i>Acomys</i> sp.	1	0(0.0)		
<i>Grammomys</i> sp.	3	0(0.0)	7.198	0.515
<i>Praomysdelectorum</i>	5	0(0.0)		
<i>Mus</i> sp.	2	0(0.0)		
<i>Arvicanthisnairobe</i>	6	0(0.0)		
Total	105	5(4.8)		
Sex				
Males	61	3(2.9)	0.008	0.930
Females	44	2(1.9)		
Total	105	5(4.8)		

Table 5. Prevalence of *Trypanosoma* sp. in rodents regarding to habitat types and wards.

Variable	Sample size	Infected no.(%)	Chi-square	P-value
Habitat type				
Fallow land	19	0(0.0)		
Crop fields	47	2(1.9)		
Indoor	19	2(1.9)		
Plantation forest	13	1(1.0)	2.965	0.0705
Bush	4	0(0.0)		
Near Natural forest	3	0(0.0)		
Total	105	5(4.8)		
Wards				
<i>Manolo</i>	20	2(1.9)		
Shume	39	0(0.0)	3.837	0.280
Nahasey	24	2(1.9)		
YaedaAmpa	22	1(1.0)		
Total	105	5(4.8)		

4. Discussion

Formatted: Left

The present study captured a total of 105 rodent species with 9 species recorded. *M. natalensis* and *R. rattus* dominated the total capture with more than 70% of the total capture. Indeed, predation and availability of the food might influence the increase in the relative density of individuals captured. Furthermore, density of the species, nature of trap, nature of vegetation, type of habitat and number of days for trapping might affect the total number of captured individuals. Our results are concurring with the study done by [15,16,30] where by *M. natalensis* and *R. rattus* dominated the total capture. Most of relatively large number of captured individuals were from crop field followed by fallow land and indoor whereas the bushes, plantation forest and near natural forest habitats had very low number. Indeed, this may be explained due to the fact that the nature of the vegetation found in [this habitat](#) ~~this habitat~~ may influence the distribution as well as abundance of the individual species, nature of the vegetation in turns affect the food and shelter availability in respected habitats. Furthermore, the study period coincide

with harvesting period where by the crop fields had plenty crops such as maize, yams and beans.

Indubitably, the higher abundance of *M. natalensis* in various habitats in the areas might be due to the fast reproductive cycle, having large litter size, availability of various excellent quality and quantity food options and their ecological generalist behavior [9,15]. *R. rattus* were only exclusively captured indoors on almost all districts. The present findings ~~concur~~ [concur](#) with the study conducted by [18,22]. ~~This~~ [This](#) ~~invasive commensal rodent~~ [invasive commensal rodent](#) can outcompete other species, the present findings shows all the captured individuals were indoor, although *R. rattus* can occupy various large range of habitats, such as the forests, inside buildings and areas with densely profound litter cover amongst others [17,29,34]. ~~P.~~ [P.](#) ~~delectorum~~ [delectorum](#) and *Lophuromys* spp. were captured from the forest, this in fact due to their forest specialist behavior and edge dwellers. [30] reported the highest abundance of these species in forests. *A. nairobe*, *Mus* sp. *Acomys* sp. and *Grammomys* sp. were captured in relative low numbers compared to others specie, these species have been reported to occurs in relative low numbers in Tanzania [23,26].

We recorded the Trypanosomiasis infection prevalence of 4.8%, our findings are in agreement with the recently study conducted by [36] in Morogoro, Tanzania, who recorded the overall prevalence of 4.04%. In addition, our findings are also in agreement with the study conducted by [16] who reported the prevalence of 4.29% in rodents from domestic and peri-domestic areas of Morogoro Municipality, Tanzania. [15] recorded the 4.1% prevalence in the study that involved six different regions of Tanzania namely Kilimanjaro, Mtwara, Mbeya, Singida, Tanga, Dodoma and Morogoro. Our findings are in disagreement with the study conducted by [14] in Central Tanzania with high prevalence of 22.7%, with the same study reporting low prevalence of 1.33% in Swaziland. [45] recorded the high prevalence of 24.7% in captured rodents in selected areas of Egypt. Prevalence of 21.7%, 21.3% and 20% has been recorded in Brazil, Venezuela and Italy by [4,11,24].

The low prevalence of *Trypanosoma* sp. encountered in our present study could be probably due to low number of sampled animals compared to other studies. Furthermore, the decrease in vector abundance in plague endemic area due to continuous control measures against the flea vectors that are responsible for transmitting the *Yersinia pestis* could result into low transmission rate of the *Trypanosoma* sp. in the area [44]. Moreover, rodents have shown to be immune to challenge dose of *Trypanosoma* infection hence this could also attribute to the low prevalence found during our study period.

Despite of not being significant differences between host species and *Trypanosoma* sp. prevalence, *M. natalensis* and *R. rattus* were recorded to have equal percentage (1.9%) of infection. Indeed, *M. natalensis* and *R. rattus* have been reported to be infected by different *Trypanosoma* species in different parts of the world. For instance in Tanzania they have been reported to be infected by *Trypanosoma* species by [14,15,16,36]. In addition, it has been reported in Uganda, Egypt, Brazil, Mali and Australia by [24,35,37,41,45]. [43] showed a high diversity of trypanosomes in small mammals of sub-Saharan mammals. *M. natalensis* and *R. rattus* are more adapted to domestic and [peridomestic](#) [peri-domestic](#) areas, hence they are frequently [interacts](#) [interacting](#) with humans and this could increase the risk of infections to humans [36]. Also due to their behavior and their abundance in different habitats they are in high risk of getting infected by the parasite and it is known that they frequently harbor more flea vectors [26].

Although, sex was not a significant factor to the prevalence, three males were found to be infected in contrast to the two infected females. Indeed, sex-biased parasitism was shown to favor the males, indeed, males are known to be more infected by parasites due to several factors, males have large body size hence increases the probability of being infected [28]. Furthermore, males cover large territory, in turn the movements within and across territories are increased which predispose them to risk of infection, in addition, hormones such as testosterone have been shown to lower immunity hence

increase also the risk of infection [6,20,33]. [36] reported that all the infected rodents with Trypanosomes were males, none of the females were infected.

Habitat including crop field, plantation forest and indoors were shown to harbor the infected captured rodents, with indoor and crop field having relative high number of infected rodents than plantation forest although the results was not differed significantly. Crop fields during the study field was having plenty of foods such as maize, yams and beans which in fact contributed to high abundance of the rodents. Rodents that were captured indoor (*R. rattus*) two of them was found to be infected by *Trypanosoma* sp. this indeed implies the risk of humans to get infected is much more highly since these rodents occurs within the same households, indeed, they are sharing the same house with humans [16,36]

When compared the *Trypanosoma* sp. infection between the wards, Manolo ward in Lushoto district was having two rodents that were infected while Nahasey and Yaeda Ampa wards in Mbulu district had two and one rodents infected respectively, although these findings weren't statistically significant. The presence of *Trypanosoma* sp. infection in rodents captured in the two districts have clearly shown the possibility of people to get infected by these parasites, hence it alerts the public on risk that could be attributed by the rodents as well as their flea vectors in these areas.

The present study has shed some light on the presence of *Trypanosoma* sp. infection in rodents, particularly the *M. natalensis*, *R. rattus* and *L. kilonzo* captured in plague endemic foci of Mbulu and Lushoto districts in Northern Tanzania. Trypanosomiasis is still a public health threat in most of ~~Sub-Saharan~~ Sub-Saharan countries, hence the awareness should be raised coupled with continuous surveillance and need of imposing control and prevention measures.

6-References

Formatted: Left

1. Anusha B, Srinivas M, Rao NS. Survey on rodent species composition in high altitude and tribal zone (HAT) of Andhra Pradesh. Journal of entomology and zoological studies.2018;6(6): 600-602.
2. Bush AO, Lafferty KD, LotzJM,Shostak AW. Parasitology meets ecology on its own terms: Margolis et al. revisited. J Parasitol.1997;83:575–583.
3. Dahmana H, Granjon L, Diagne C, Davoust B, Fenollar F, MediannikovO.Rodents as hosts of pathogens and related zoonotic disease risk. Pathogens.2021;9(3):202.
4. De Carnieri I, Castellino S. *Trypanosomalewisi* in un allevamento lombardo di rattialbini. Parassitol. 1964 ;6:95-99.
5. DokePP, Kar A. A fatal case of *Trypanosoma lewisi* in Maharashtra, India. Ann. Trop. Med. Publ. Hlth. 2011;4:91-95.
6. Fagir DM, Horak IG, Ueckermann EA, Bennett NC, Heike H.Ectoparasite diversity in the eastern rock sengis (*Elephantulusmyurus*): the effect of seasonality and host sex. African Zoology. 2015;50(2):109-117.
7. Gao JM, Truc P, Desquesnes M, Vincendeau P, Courtois P, Zhang X, Li SJ, Jittapalapong S, Lun ZR.A preliminary serological tudy of *Trypanosoma evansi* and *Trypanosoma lewisi* in a Chinese human population. Agric Nat Resour. 2018;52(6):612–616.
8. Gholipoury M, Rezai HR, Namroodi S, Arab KF. Zoonotic and Non-zoonotic Parasites of Wild Rodents in Turkman Sahra, Northeastern Iran. Iranian journal of parasitology. 2016;11(3):350–357.
9. Haule M, Lyamuya EF, Matee MI, Kilonzo BS, Hang'ombe BN. Factors associated with flea infestation among the different rodent species in Mbulu and Karatu districts, northern Tanzania. Tanzania Journal of Health Research.2013;15 (3).

10. Hendrix CM, Robinson ED. Diagnostic parasitology for veterinary technicians.4th Edition Elsevier; 2012.
11. Herrera L, Urdaneta-Morales S.Synanthropic rodent reservoirs of *Trypanosoma (Schizotrypanum) cruzi* in the valley of Caracas,Venezuela. Rev. Inst. Med. Trop. São Paulo.1997;39:279 -282.
12. Kamaruzaman INA, TingHW, Mokhtar MAM, Yuan YK, Shah AWG, Hamid, F. F. et al. First case report on molecular detection of *Trypanosoma lewisi* in an urban rat in Kelantan, Malaysia: An accidental finding. Journal of advanced veterinary and animal research.2021;8(3):435–439.
13. Kasso M. Pest Rodent Species Composition, Level of Damage and Mechanism of control in Eastern Ethiopia. International Journal of Innovation and Applied Studies. 2013;4(3): 502-511.
14. KatakwebaAA, Mulungu LS, Eiseb SJ, Mahlaba TAA, Makundi RH, Massawe A W, et al. Prevalence of haemoparasites, leptospirae and coccobacilli with potential for human infection in the blood of rodents and shrews from selected localities in Tanzania, Namibia and Swaziland. African zoology. 2013;47(1):119-127.
15. Katakweba AAS, Kipanyula MJ, Durnez L, Mgode GF, Mhamphi G, Luziga C, et al. Rodents and Shrews as Vectors of Zoonotic Spirochetes and Trypanosomes in Tanzania. Tanzania Veterinary Journal.2013;28(1);14-19.
16. Katakweba AAS. The Prevalence of Haemoparasites in Rodents and Shrews Trapped from Domestic and Peridomestic Houses in Morogoro Municipality, Tanzania. A Hidden Public Health Threat. Tanzania Veterinary Association Proceedings.2018;36:75-82.
17. Kilonzo BS, Mbise TJ, MwalimuDC,Kindamba L. Observation on the endemicity of plague in Karatu and Ngorongoro, northern Tanzania. Tanzania Health research Bulletin. 2006;8(1).
18. Kilonzo BS, Mhina J, SabuniC,Mgode G. The role of rodents and small carnivores in plague endemicity in Tanzania. Belgian Journal of Zoology. 2005;135(Suppl.):119-125.

19. Kingdon J. The Kingdon Field Guide to African Mammals. Academic Press, San Diego;2015.
20. Krasnov BR, Shenbrot GR, Kholkhlova IS, Stanko M, Morand S, Mouillot D. Assembly rules of ectoparasite communities across scales: combining patterns of abiotic factors, host composition, geographical space, phylogeny and traits. *Ecography*.2015;38:184-197.
21. Laakkonen J, Smith A, Hildebrandt K, Niemimaa J, Henttonen H. Significant morphological but little molecular differences between *Trypanosoma* of rodents from Alaska. *J. Parasitol.* 2005;91:201–203.
22. Laudisoit A, Leirs H, Makundi RH, Van Dongen S, Davis S, Neerinckx, et al. Plague and the human flea, Tanzania. *Emerging Infectious Diseases*. 2007;13:687-693.
23. Laudisoit A, Leirs H, Makundi RH, Krasnov B. Seasonal and habitat dependence of fleas parasitic on small mammals in Tanzania. *Integrative Zoology*.2009;4:196-212.
24. Linardi PM, Botelho JR. Prevalence of *Trypanosoma lewisi* in *Rattus norvegicus* from Belo Horizonte, State of Minas Gerais, Brazil. *Mem. Inst. Oswaldo Cruz Rio De Janeiro*. 2002;97(3):411-414.
25. Lun ZR, Reid SA, Lai DH, Li FJ. Atypical human trypanosomiasis: a neglected disease or just an unlucky accident?. *Trends in parasitology*. 2009;25(3):107–108.
26. Makundi RH, Massawe AW, Borremans B, Laudisoit A, Katakweba A. We are connected: flea-host association networks in the plague outbreak focus in the Rift Valley, northern Tanzania. *Wildlife research*. 2015;42(2015):196-206.
27. Matthee S, Horak IG, Beaucournu JC, Durden LA, Ueckermann EA, McGeoch MA. Epifaunistic arthropod parasites of the four-striped mouse, *Rhabdomys pumilio*, in the Western Cape Province, South Africa. *The Journal of parasitology*. 2007;93(1):47–59.
28. Matthee S, McGeoch MA, Krasnov BR. Parasite-specific variation and the extent of male-biased parasitism; an example with a South African rodent and ectoparasitic arthropods. *Parasitology*.2010;137:651–660.

29. Michelle G, Chris R, Dickman C, Warren G. Use of habitat by the blackrat (*Rattus rattus*) at North Head, New South Wales: an observational and experimental study. *Austral Ecology*.2000;25:375–385.
30. Mulungu LS, Makundi HR, Massawe WA, MachanguRS, Mbije NE. Diversity and distribution of rodent and shrew species associated with variations in altitude on Mount Kilimanjaro, Tanzania. *Mammalia*. 2008 ;72(2008); 178-185.
31. Ortiz PA, Garcia HA, Lima L, Da Silva FM, Campaner M, Pereira CL, et al. Diagnosis and genetic analysis of the worldwide distributed *Rattus*-borne *Trypanosoma (Herpetosoma) lewisi* and its allied species in blood and fleas of rodents. *Infection, genetics and evolution*. 2017;63:380–390.
32. Phukon M, Borah KR. Species composition of field rodents in rice-vegetable cropping system at upper Brahmaputra valley zone, Assam. *Journal of entomology and zoological studies*.2019;7(1):961-969.
33. Postawa T, Nagy Z. Variation of parasitism patterns in bats during hibernation: the effect of host species, resources, health status, and hibernation period. *Parasitol Res*.2016;115:3767–3778.
34. Pryde M, Dilks P, Fraser I. The home range of ship rats (*Rattus rattus*) in beech forest in the Eglinton Valley, Fiordland, New Zealand: A pilot study. *New Zealand Journal of Zoology*.2005;32(3):139–142.
35. Salzer JS, Pinto CM, Grippi DC, Williams-Newkirk AJ, Peterhans JK, Rwego IB, et al. Impact of anthropogenic disturbance on native and invasive trypanosomes of rodents in forested Uganda. *EcoHealth*.2016;13:698–707.
36. SamijiAM, KatakwebaAS, PhiriEC. Trypanosomes Infection in Rodents and their Zoonotic Potential from Ruaha Ward in Kilosa District, Tanzania. *Proceedings of the 2nd SUA Scientific Conference held at SUA from 25th to 26th 2021*.126-133.
37. Schwan TG, Lopez JE, Safronetz D, Anderson JM, Fischer RJ, Maïga, et al. Fleas and trypanosomes of peridomestic small mammals in sub-Saharan Mali. *Parasit. Vectors* .2016;9:541.
38. Shah I, Ali US, Andankar P, Joshi RR. Trypanosomiasis in an infant from India. *J Vector Borne Dis*. 2011; 48:122–3.

39. Soulsby EJL. Helminths, Arthropods and Protozoa of Domesticated Animals. 7th edition, Lea Febiger., Philadelphia;1982.
40. TangHJ, Lan YG, Wen YZ, Zhang XC, Desquesnes M, Yang TB, et al. Detection of *Trypanosoma lewisi* from wild rats in Southern China and its genetic diversity based on the ITS1 and ITS2 sequences. *Infect. Genet. Evol.* 2012 ;12:1046–1051.
41. Thompson CK, Godfrey SS, Thompson RC. Trypanosomes of Australian mammals: a review. *Int. J. Parasitol. Parasites Wildl.* 2014;3:57–66.
42. Verma A, Manchanda S, Kumar N, Sharma A, Goel M, Banerjee PS, et al. Case report: *Trypanosoma lewisi* *T. lewisi*-like infection in a 37-day-old Indian infant. *Am J Trop Med Hyg* 2011; 85(2):221–4
43. Votýpka J, Stříbrná E, Modrý D, Bryja J, Bryjová A, Lukeš J. Unexpectedly high diversity of trypanosomes in small sub-Saharan mammals. *International journal for parasitology.* 2022;52(10):647–658.
44. Ziwa MH, Matee MI, Hang'ombe BM, LyamuyaEF, Kilonzo BS. Plague in Tanzania: An overview. *Tanzania Journal of Health Research.* 2013;15(4).
45. Dahesh AM, Mikhail WM. Surveillance of *Trypanosoma* spp. of rodents and studies in their transmission probability by fleas in some rural Egyptian areas. *Journal of the Egyptian Society of Parasitology.* 2016;46(1):157-166.