

## Original Research Article

### SEROPREVALENCE OF CYTOMEGALOVIRUS AMONG PAEDIATRIC PATIENTS WITH SEIZURES ATTENDING FEDERAL NEUROPSYCHIATRIC HOSPITAL, KADUNA , NORTH WEST NIGERIA.

#### ABSTRACT

**INTRODUCTION:** Cytomegalovirus causes a congenital infection, which poses a great public Health challenge such as hearing loss, seizures and mental disorder among children.

**METHODOLOGY:** Enzyme Linked Immunosorbent Assay (ELISA) was used to screen serum samples ( 90 ) collected from Patients diagnosed with Seizure between the ages 0-13 years for Cytomegalovirus IgG specific antibodies.

**RESULT:** The presence of CMV IgG specific antibody among Seizure patients obtained from this study recorded a prevalence of 68(75.5%). The age groups 9-13 years presented highest CMV (IgG) specific antibodies response 36(40.0%), followed by ages 5-8 years 22(24.4%) and least for ages 0-4 years 10 (11.1%),  $P>0.005$ . The CMV (IgG) specific antibodies response with respect to gender presented male with highest prevalence 50 (55.5%) than their female 18 (20.0% ) counterparts  $P<0.005$ . The study has indicated significant association between the CMV (IgG) specific antibodies response and Seizures.

**CONCLUSION:** Therefore CMV is highly prevalent within the study area and the need for necessary facilities to be available in screening pregnant women of the virus in other order to prevent the spread vertically to their fetus thereby increases the risk of Seizures.

**Key Words:** Cytomegalovirus, Seizure, IgG, ELISA

Comment [C1]: Patients presenting with

Comment [C2]: This study has not proved this association

Comment [C3]: fetuses

## INTRODUCTION

Cytomegalovirus (CMV) is a member of the Herpes family with double stranded DNA that measure approximately 200nm [1,2]. The virus is responsible for a wide range of infections in human of all ages [3]. CMV causes enlargement of cells in the body, hence the prefix Cytomegalo [4]. Infection with CMV is asymptomatic at all ages, unlike in foetus and neonates that could lead to neurological complications and death [5].

The Virus could remain latent in the body. However, severe recurrent infection may occur in immunocompromised patients [6,7]. Separate studies by other researchers have proved that Cytomegalovirus causes congenital infection and brain damage which could lead to defects in babies of infected mothers [8,9,10].

The Virus can be transmitted through close contact with body fluids of an infected person[11,12,13]. This infection poses important public health problem because of its frequency in severe congenital anomalies such as microcephaly, ~~intrauterine,~~ growth retardation, mental retardation, inguinal hernia in males, seizures, loss in hearing and neuromuscular abnormalities [14,15].

Congenital CMV infection has been reported in various countries,15-20% of sensor neural hearing loss and 7% of Cerebral palsy in United Kingdom, and rates of 0.15-2.2% of live birth from United States of America [2,16,17]. On the same hand, sero-prevalence rate of CMV IgG antibodies among pregnant women In Singapore, Thailand, Iran, Turkey and Spain have also been reported as; 87%, 100%, 93%, 98.5% and 84% respectively [19,20,21,22].

In Nigeria, Cytomegalovirus infection was also reported as 45% and 33% in breast milk,86.15% (IgG) among expected mothers, 77.1% (IgG) among primigravid women in Bida, 97.2 % (IgG) among pregnant women in Lagos, 80.5%(IgG) among children hearing

Comment [C4]: delete comma

Comment [C5]: one word: sensorineural

1 impairment and 10.5%(IgM) among pregnant women in Kafanchan Kaduna State  
2 [23,24,25,26,27].

3 Cytomegalovirus Congenital seizures disorder account for 10% of babies born with  
4 symptomatic infection and it is a very common abnormality associated with the infections. It  
5 is usually associated with children severely affected by systematic CMV infections of the  
6 nervous system. A seizure may occur at birth or within six months of life with varying  
7 severity[8]. The aim of this study therefore is to determine the seroprevalence rate of  
8 cytomegalovirus among pediatric patients clinically diagnosed of seizure at Federal  
9 Neuropsychiatric Hospital, Barnawa, Kaduna State.

Comment [C6]: presenting with seizures

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

## METHODOLOGY

### Study design/Area:

The study was a cross sectional study carried out at Federal Neuropsychiatric Hospital Kaduna a Tertiary Mental Health Hospital with 200 beds capacity. It is an institution for training Mental Health Professionals. Patients who were diagnosed clinically of seizure from ages 0-13 years at the Child and Adolescent unit ~~where were~~ used for this research.

### Sample Size:

The sample size was calculated using the descriptive studies formula [28]

$$n = \frac{z^2 pq}{d^2}$$

$$d^2$$

Where the P = Value of proportion of interest ( If no information is known about p then p= 0.5)

d= Tolerance eg within 0.05

Hence:  $n = \frac{1.96^2 \times 10/100 (1-10)}{0.05^2} = 136$  samples.

$$0.05^2$$

Comment [C7]: consider revising as suggested

### Ethical Matters/ Demographic Data

An ethical approval was obtained from Federal Neuropsychiatric Hospital Kaduna ethical committee before embarking on the work. Also consent form was designed and issued to the patients to seek their parent /guardian approval. A questionnaire was designed to collect demographic data of the entire patients.

### Inclusion Criteria

The study included paediatric patients clinically diagnosed with seizure within ages 0-13 years whose patient/guardian consented for the investigation.

1 **Exclusion Criteria**

2 The study excluded patients outside the ages 0-13 years whose ~~patient~~parent/guardian did not  
3 consent to the investigation.

4 **Sample Collection:**

5 Two milliliters of blood were collected via the ante cubical vein using sterile vacutainer into a  
6 sterile plain tube and labelled appropriately. The blood was centrifuged at 3000rpm for  
7 5minutes. The sera was harvested into clean cryo-vials and stored at -20°C until it is required  
8 for use. A total of 90 samples were collected during three months of this research.

9 **Detection of IgG Antibodies Using Enzyme Linked Immunosorbent Assay (ELISA)**

10 Samples were analyzed for the presence of IgG antibodies using the Enzyme Linked  
11 Immunosorbent Assay (ELISA) method according to the kit's manufacturer's instruction  
12 made by ASIA –LION Biotechnology Company China. Samples were incubated with mouse  
13 monoclonal antibody against human IgG bound to the solid surface for a microtitre well.  
14 Patient IgG is captured by the surface bound antibody. Unbound serum component are  
15 washed away, patient antiCMV IgG antibodies are detected and bound by an  
16 immunocomplex enzyme conjugate, consisting of CMV antigen which is conjugated to horse  
17 radish peroxidase. Unbound conjugate is removed by aspiration and washing. Substrate is  
18 then added and incubated in the presence of bound enzyme the substrate is converted to end  
19 product. The absorbance of this end product is read spectrophotometrically at 450 nm and is  
20 directly proportional to the concentration of IgM antibodies to CMV antigen present in the  
21 sample [26].

**Comment [C8]:** consider simplifying as suggested

22  
23  
24  
25  
26

## RESULT

A total of 90 samples were collected from Patients diagnosed with Seizures between the ages 0-13 years and screened for Cytomegalovirus IgG specific antibodies. The presence of CMV IgG among seizure patients obtained from this study was presented as prevalence of 68(75.5%).

Table 1. however present the seroprevalence of CMV IgG in relation to the age groups of the patients. CMV seroprevalence IgG specific antibody was highest in ages 9 - 13 years 36(40.0%) and least in ages 0 -4 years 10(11.1%).While ages 5-8 years has 22(24.4%) prevalence. The value however is statistically significant with  $X^2= 0.024$ ,  $df= 2$   $P>0.05$   $CL=95\%$ .

Table 2 shows relationship between IgG antibodies with the genders. The Males presented prevalence of 50(55.5)% IgG specific antibodies response while the females presented 18(20.0%). However, there was no statistical significant difference between the seroprevalence IgG specific antibody responses in relation to gender from this study with  $X^2=27.7$ ,  $df=1$ ,  $CL=95\%$ ,  $P<0.05$ .

Fig (i) is a pie-chart showing the frequency distribution of patient parent/guardian occupation. The data shows that 41% are civil servant, 32% businessmen, and 17% are farmers.

Fig (ii) is a pie-chart showing the frequency distribution of patients area of residence. The chart reveals 57% living in Urban settlement, 14% semi-urban and 19% in rural area.

Fig (iii) is a pie-chart demonstrating the patients educational status. The chart shows that 54% are in school, 8% have been withdrawn from school while 28% were not in school

Comment [C9]: presenting with seizures

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

Table 1: Prevalence of CMV (IgG) Among Patients in Relation to Age Group

Age(years)	Samples Screened	Sex		CMV (IgG)	
		Males	Female	Positive	Negative
0 – 4	16	16	0	10(11.1%)	6(6.6%)
5 – 8	32	24	8	22(24.4%)	10(11.1%)
9 – 13	42	30	12	36(40%)	6(6.6%)
<b>TOTAL</b>	<b>90</b>	<b>70</b>	<b>20</b>	<b>68(75.5%)</b>	<b>22(24.4%)</b>

$X^2 = 0.024$  df= 2 p>0.05 CL = 95%


1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

TABLE 2: Prevalence of CMV (IgG) in Relation to Gender

Gender	Samples Screened	CMV ( Positive)	CMV(Neg)
Male	70	50(55.5%)	20(22.2%)
Females	20	18(20.0%)	02(2.2%)
<b>TOTAL</b>	<b>90</b>	<b>68(75.5%)</b>	<b>22(24.4%)</b>

$\chi^2 = 27.7$  df = 1 P<0.05 CL= 95%

UNDER PEER REVIEW



1  
2  
3  
4




Fig.1: CMV (IgG) Positive Parents/Guardian Occupation

5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

1  
2  
3  
4  
5




Fig.2: Frequency Distribution of CMV (IgG) Positive Patients with Respect to Residence

1  
2  
3  
4  
5  
6  
7  
8  
9




Fig.3:Frequency of Distribution of CMV (IgG) Positive Patients with Respect to Educational Status

44  
45  
46  
47  
48  
49

## DISCUSSION

Cytomegalovirus infection is commonly spread in developing and low social economic countries. CMV infection was also earlier reported to be the viral cause of birth defects in industrialized countries [29]. This study however has recorded a prevalence of 75.5% of CMV (IgG) in Seizure patients.

In other separate previous studies, it revealed 80% prevalence of CMV infection in children who attend daycares, particularly children younger than 2 years was reported [26,30,31,32].

The 75.5% prevalence obtained in this study however is similar to the work of Redwan, *et al.*, [30] who also reported prevalence of 80.7%. Similarly a separate study, Adisa, *et al.*, [31] reported higher prevalence of 98.9% IgG specific antibodies to CMV in samples of 94 mothers while their corresponding newborns had 86.2% sero prevalence rate. More so, Yeroh *et al.* [33] recorded 94.8% of CMV IgG among pregnant women in Kaduna state. Although infection of the mother during pregnancy could be without consequence, however, there may be a serious effect on the fetus [2]. Other reports from other countries was also documented; In Singapore, Thailand, Iran, Turkey and Spain the seroprevalence rate of CMV IgG antibodies among pregnant women was reported as; 87%, 100%, 93%, 98.5%, and 84% respectively [19,20,21,22].

In Nigeria, a study conducted in 2008, reported a prevalence of 45.0% and 33% IgM antibodies among breastfeeding mothers and their infants [23]. Similarly, Okwori *et al.* [24] in a study among expectant mothers in Bida, Nigeria, reported IgG antibody prevalence of 86.1% among multigravid women and 77.1% among primigravid women. Therefore, there is justification on high rate CMV in Children since it can be transmitted from mother to child.

From this study it was revealed that ages 9-13yrs had the highest prevalence 68(40%), followed by ages 5-8yrs 22(24.4%) and least for ages 0-4yrs 10(11.1%). This trend had

Comment [C10]: patients presenting with seizures

1 earlier been explained by Redwan, *et al.*, [30] who explained that CMV Sero prevalence  
2 increases gradually with age. This findings however shows statistically significant with  $X^2=$   
3 0.024,  $df=2$ ,  $CL=95\%$ ,  $P>0.05$ .

4 In this studies males 50(55.5%) have a higher frequency than female 18(20.0%) as shown in  
5 table 2. This is not in agreement with the work of Redwan, *et al.*, [30], where females have a  
6 86.8 % and males 75%. This shows a gradual increase in the prevalence between female and  
7 male. However, there is no statistical significant difference between the two genders towards  
8 contracting the infection with  $X^2= 27.7$ ,  $df=1$ ,  $CL=95\%$ ,  $P< 0.05$ .

9 This study further reveals, that 41% of the patient's parent/guardian of CMV positive (IgG)  
10 patients were civil servants, while 32% were businessmen and 17% farmers. This trend with  
11 high value from civil servants was explained earlier by other authors that civil servants and  
12 business class keep their children to a daycare during their working hours and this could  
13 predispose the children to the virus since most of the care givers are either secondary school  
14 level graduate or not even knowledgeable of the virus since is not common like the HIV  
15 [26,31,32].

16 The study further reveals that 54% of the patients are in school, 8% were withdrawn from  
17 school by parent and 28% are not in school. This suggests that Seizures ~~has~~ have a greater  
18 negative effect on children, especially on their educational and social status. Some parent will  
19 prefer their children at home due to the stigmatization and trauma associated to seizures in  
20 any episode that occur to the patient. Seizure is attributed to believes by most Africans that is  
21 associated with Evil spirit, and most times such patients are taken to prayer houses or native  
22 doctors especially those in the rural area before resolving to the orthodox.

Comment [C11]: please delete

Comment [C12]: percentages should add up to 100%

Comment [C13]: with

23

24

1       **Conclusion**

2       Cytomegalovirus is highly associated with Seizure, therefore the need for public awareness  
3       with the purpose of educating the populace on CMV and its public health challenge  
4       associated with it, in order to eradicate stigmatization in relation to the patient's condition,  
5       traumatization that comes with seizure and its social economic challenge . There is a need for  
6       training for community and primary Health care workers so that they can be involve in  
7       managing of such patients. The need of awareness to day care and crèche worker and  
8       pregnant women on how to prevent the spread of the disease through hygienic way of live.

9       **Recommendation**

10      Based on our findings from this study it is recommended that:

- 11      -Aggressive awareness is highly needed to the populace on the consequences of the infection.  
12      -Routine screening of sexually active women is needed to avoid transmission from mother to  
13      foetus of the infection  
14      - Hand wash should encourage especially after changing diapers  
15      - Personal hygiene should encourage especially to those handling children and pregnant  
16      women.  
17      - There is a need to develop Vaccine or any prophylaxis against CMV infection

18      **COMPETING INTERESTS DISCLAIMER:**

19  
20      Authors have declared that no competing interests exist. The products used for this research  
21      are commonly and predominantly use products in our area of research and country. There is  
22      absolutely no conflict of interest between the authors and producers of the products because  
23      we do not intend to use these products as an avenue for any litigation but for the  
24      advancement of knowledge. Also, the research was not funded by the producing company  
25      rather it was funded by personal efforts of the authors.  
26

1  
2 **REFERENCE**

- 3 [1]. Boppana, S. B.; Rivera, L. B. ; Fowler, K. B.; Mach, M. and Britt, W. J. (2001):  
4 Intrauterine Transmission of Cytomegalovirus to Infants of Women with  
5 Preconceptional Immunity. *New England Journal of Medicine*. 3334:1366 – 1371(s).
- 6 [2]. Nathalie, J. S. and Richard ,W. E. (2005):Statement in: Diagnostic Procedures for  
7 Viral, Rickettsia and Chlamydial Infections 6th Edition pp. 321 –329
- 8 [3]. Suzuki Y, Taribe Y, Mogami Y, Yanagihara K, Nishikawa M, (2008) Epilepsy in  
9 patients with congenital Cytomegalovirus infection .*Journal of Brain Dev*. 2008,  
10 *Jun*;30(6);420-4.doi:1010161 j. braindev
- 11 [4]. Prescott, M. L.; Harley, J. P.; and Klein, D. A; (2005): Microbiology. 6<sup>th</sup> edition pp  
12 862.
- 13 [5]. Griffiths, P.D. and Walter, S. (2005): Cytomegalovirus. *Current opinion in infectious*  
14 *Disease* 18(3):242-245
- 15 [6]. Timbury , C. M. (1986) : Herpes Disease: Notes on Medical Virology 8th edition: pp  
16 88- 89.
- 17 [7]. Bello, C. and Whittle, H. (1991): Cytomegalovirus infection in Gambian Mothers  
18 and their Babies : *Journal of Clinical Pathology* 1991; 44: 366-369
- 19 [8]. Spring 1998: cytomegalovirus Registry; [www.bcm.tmc.edu/pedi/infect/cmv](http://www.bcm.tmc.edu/pedi/infect/cmv)
- 20 [9] Fowler, K. B. and Pass, R. F. (2006): Risk factors for congenital cytomegalovirus  
21 infection in the offspring of young women: Exposure to young children and recent  
22 onset of sexual activity. *Paediatric*; 118: 286-292
- 23 [10]. Dworsky, M. E.; Welch, K.; Cassady, G. and Stagno, S. (1983). Occupational risk for  
24 primary cytomegalovirus infection among paediatric health-care workers. *New*  
25 *England Journal of Medicine*, 309(16), 950–953.
- 26 [11]. Gabrielli, L.; Lazzaroho, T.; Foschini, M. P.; Lanari, M.; Guerra, B.; Eusebi, V. *et al.*,  
27 (2003): Horizontal in utero acquisition of cytomegalovirus infection in a twin  
28 pregnancy. *Journal of clinical microbiology*: 14(3) 1329 – 1331(s).
- 29 [12]. Stagno, S.; Pass, R. F.; Cloud G. Brith; W. J.; Henderson, R. F.; Walton, P. O. *et al.*,  
30 ( 1986): Primary Cytomegalovirus infection in pregnancy incidence,  
31 transmission of foetus and clinical outcome.; *Journal of the American Medical*  
32 *Association* 256: 1904-1908
- 33 [13]. HO, M. (1990) Epidemiology of Cytomegalovirus infection. *Reviews of Infectious*  
34 *Diseases*; 12(7) :5701-5710
- 35 [14]. Jawetz, M. and Adelberg (2007): Text Book Medical Microbiology 24th Edition.  
36 Pp. 441 – 445.

- 1 [15]. Morton, C. C. and Nance, W. E. (2006): New-born hearing screening-a silent  
2 revolution. *New England Journal of Medicine*, 354:2151-2164.
- 3 [16]. Negro, G. (2005): Passive immunization during pregnancy for congenital  
4 Cytomegalovirus infection. *New England Journal of Medicine*; 353 (13): 1350- 1362.
- 5 [17]. Grosse, S. D.; Dollard, S. C.; Ross, D. S.; (2008) Congenital cytomegalovirus (CMV)  
6 infections a cause of permanent bilateral hearing loss: A quantitative assessment.  
7 *Journal of Clinical Virology. Epub ahead of print.*
- 8 [18]. Lazzarotto, T.; Varani, S.; Guerra, B.; Nicolosi, A.; Lanrari, M.; Landini, M. P.;  
9 (2001): Prenatal Indicators of Cytomegalovirus infection, *Journal of. Pediatr*;  
10 (13791): 90 – 95.
- 11 [19]. Wang, A.; Tank, H.; Tee, C. S.(2000): Seroprevalence of Cytomegalovirus,  
12 Toxoplasmosis, and Parvovirus in pregnancy. *Singapore Med J. ;41(4):151–155.*
- 13 [20]. Arabpou, M.; Kariyane, K.; Jankhah, A.; (2007). Human cytomegalovirus in women  
14 of child bearing age throughout Fars province-Iran: A population based cohort study.  
15 *Malaysian Journal of Microbiology.;3(2):23–28.*
- 16 [21]. Ali, S.; Askin, G.; Hakan, O; (2007): CMV seroconversion in pregnancy and the  
17 incidence of congenital CMV infection. *Turk J Peadiatr. ; 49:30–36.*
- 18 [22]. Estripeant, D.; Moreno, Y.; Ahumada, R. (2007): Seroprevalence of Cytomegalovirus  
19 infection in peuperal women and its impact on their newborns. *Ann Pediatr. ;66:135–*  
20 *139.*
- 21 [22]. Kassim O. O.;Afolabi, O.; Ako-Nai, K. A.;Torimiro, S. E. A.; Littleton, G. K.; Oke,  
22 O. O. *et al.*, (1987): Cytomegalovirus antibodies in breast milk and sera of  
23 mother- infant pairs. *J Trop Pediatr*; 33(2) : 75-77.
- 24 [24]. Okwori, A.; Olabode, A.; Emmuwen, E.; Lugos, M. ;Okpe, E.; Okopi, J.*et al.*, (2008):  
25 Sero- Epidemiological Survey of human cytomegalovirus infection among expectant  
26 mothers in Bida Nigeria. *The Internet Journal of infectious Diseases;6 (2)*
- 27 [25]. Akinsegun, A. A.; Kabiru, A. R.; Adeniyi, A. A.; Kikelomo, O. W. (2011) Sero  
28 prevalence of Cytomegalovirus antibodies amongst normal pregnant women in  
29 Nigeria. *Int. J Women Health 2011; 3 : 423-428.*
- 30 [26]. Edward, D. S., Edward, I. U., Nwankiti, O., Shallangwa, I. B., Abdullahi,  
31 M.M.,(2015): The Sero-Prevalence of Cytomegalovirus (IgM) Antibodies among  
32 Pregnant Women Attending Ante-natal Clinic at the General Hospital Kafanchan,  
33 Kaduna State Nigeria. *British Microbiology Research Journal;2015,Vol*  
34 *9(5)pp1-6 , 2015. Article no. BMRJ. 16863 ISSN : 2231-0886. DOI:*  
35 *10.9734/BMRJ/2015/16863*
- 36 [27]. Edward I. U., Elayo S, Ogboi J. S., Musa A. M.1,Edward D. S, Shallangwa B. I,  
37 Meshubi F.1, Isaac A. (2017):Seroprevalence of Cytomegalovirus Among Paediatric  
38 Patients with Hearing Loss Attending National Ear Care Centre Kaduna  
39 Northwest Nigeria. *Pathology and Laboratory Medicine 2017; 1(2): 48-53*  
40 <http://www.sciencepublishinggroup.com/plm> doi: 10.11648/j.plm.20170102.14

- 1 [28]. Daryl, S. Paulson (2008) Biostatistic and Microbiology A Survival Manual Pp 4:73-  
2 81 ISBN:978-0-387-77281 doi:10.1007/978-0-387-77282-60.
- 3 [29]. Staras SA, Dollard SC, Rayford KW, Flanders WD, Pass RF, Cannon MJ. (2006)  
4 Seroprevalence of Cytomegalovirus infection in the United State (1988-1994).  
5 *Clinical Infectious Diseases.* ;43(9):1143- 1151.
- 6 [30]. Redwan, A. N. Ahmed, M.M. AlAwfi, M.S. (2011) Prevalence Study of  
7 Cytomegalovirus infection among foreign man power in Jeddah Saudi Arabia; *African*  
8 *Journal Of Microbiology Research Vol 5(17) pp 2539-2549*
- 9 [31]. Adisa, T. Bukbuk, D. Harry, T.(2010): Maternofoetal transfer of Cytomegalovirus  
10 IgG antibodies in Maiduguri, North Eastern Nigeria. *The Internet Journal of*  
11 *Microbiology. Volume 9 Number 1. DOI: 10.5580/c53*
- 12 [32]. Brooks, G. F.; Butel, J. S. and Morse, S. A. (2007) : Herpes Viruses In: Jawetz,  
13 Melnick and Adelberg's (Ed) Medical Microbiology. 24th Edition, McGraw-Hill.  
14 New York 441-445
- 15 [33]. Yeroh, M. Aminu, M. Musa, B. O. (2014): Seroprevalence of Cytomegalovirus  
16 infection among Pregnant Women in Kaduna Nigeria . *African Journal of Clinical*  
17 *and Experimental Microbiology Vol 16(1) 37-44*

18  
19  
20  
21  
22  
23  
24  
25  
26  
27