

Assessment of Renal Functions among Sudanese Patients with Acute Lymphoblastic Leukemia's Receiving Chemotherapy in Radiation and Isotopes Center of Khartoum (RICK), Sudan

Abstract

Background: A wide array of disturbances in electrolyte equilibrium is commonly seen in patients with acute leukemia (AL). These abnormalities present a potential hazard in these patients, as that of enhancing the cardio-toxic effects of certain chemotherapeutic regimens.

Material and Methods: This study was designed as interventional study, which includes 50 samples, the data was collected from hospital archives, include both male and female (the males is 24 with 48% while the rests 26 is females with 52%), from different area in Sudan (east is 2%, west is 64%, north is 10%, south is 12%, and center is 12%), have different classification of Acute Lymphoblastic Leukemia's (L1 is 17 with 34%, L2 is 27 with 54% and L3 is 6 with 12%). Carried out in Radiation and Isotopes Center of Khartoum in Khartoum state, during the period from February 2020 to January 2021, and the obtained data were analyzed by SPSS.

Results: The result of this study showed that there was insignificant difference ($p > 0.05$) in the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, and magnesium in the study groups before and after treatment, and significant increase in the serum level of potassium in the study groups after treatment p. value of 0.007, the mean of serum level of potassium before and after treatment is 2.688 and 3.702 respectively. And insignificant difference ($p > 0.05$) in the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium in the study groups before and after treatment according to gender and FAB classification.

Conclusion: The study conclude that the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, and magnesium was not affected by treatment, but the potassium is increased after treatment. And the gender and classification of Acute Lymphoblastic Leukemia's have insignificant effect on the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium.

Key words: Acute Lymphoblastic Leukemia's, urea, creatinine, uric acid, sodium, potassium, calcium, magnesium, and phosphorous, Sudanese.

Introduction:

“The kidneys are vital organs that perform a variety of important functions, homeostasis (maintenance of equilibrium) of the body’s water, electrolyte and acid base status, and participation in hormonal regulation”⁽¹⁾.

“Acute lymphoblastic leukemia (ALL) is a cancer of the lymphoid line of blood cells characterized by the development of large numbers of immature lymphocytes. Symptoms may include feeling tired, pale skin color, fever, easy bleeding or bruising, enlarged lymph nodes, or bone pain. As acute leukemia, ALL progresses rapidly and is typically fatal within weeks or months if left untreated, and it is the most common type of leukemia in young children”⁽²⁾. “Acute myeloid leukemia (AML): which affect myeloid cells and grows quickly, hepatic involvement in acute leukemia’s is usually mild and silent at the time of diagnosis”⁽³⁾. “Some study showed liver infiltration in 95% of ALL and 75% of AML patients”⁽⁴⁾. “In ALL, infiltration was confined to the portal tract, whereas in AML, infiltration was observed in both portal tract and sinusoids, massive leukemic cells infiltration of the liver may present as a fulminant hepatic failure”⁽⁵⁾. “The aminotransferases are normally present in the serum in low concentration; these enzymes are released into the blood in greater amounts when there is damage to the liver cell membrane resulting in increased permeability”⁽⁶⁾. “The activation of ALP and GGT are elevated in hepatic infiltration by leukemic cells”⁽⁷⁾.

“In acute lymphocytic leukemia, renal complications occur due to several factors, including leukemic infiltration of the kidneys, therapy-related side effects such as tumor, nephrotoxic drugs, and septicemias”⁽⁸⁾. “Hyperuricemia, as a manifestation of tumor lysis syndrome, is a well-recognized complication and in most cases, it occurs after the initiation of chemotherapy. The most common symptoms in children are easy bruising, pale skin, fever, and an enlarged spleen or liver”^(9, 10 and 11).

Materials and Methods:

This is a facility based, interventional study, which includes fifty samples, the data was collected from hospital archives, include both male and female (the males is 24 with 48% while the rests 26 is females with 52%), from different area in Sudan (east is 2%, west is 64%, north is 10%, south is 12%, and center is 12%), have different classification of Acute Lymphoblastic Leukemia's (L1 is 17 with 34%, L2 is 27 with 54% and L3 is 6 with 12%). Carried out in Radiation and Isotopes Center of Khartoum in Khartoum state, during the period from February 2020 to January 2021, and the obtained data were analyzed by SPSS.

Study Area:

This study was conducted in Khartoum state at Radiation and Isotopes Center of Khartoum (RICK), is the largest hospital in Sudan and infective patients from all the century.

Study Population:

Participants involved in this study were Sudanese patients with Acute Lymphoblastic Leukemia's for different classification of Acute Lymphoblastic Leukemia's, and including both males and females from different area in Sudan, attending the Radiation and Isotopes Center of Khartoum (RICK).

Data Collection:

Data was collected form hospital archives.

Sampling Technique:

Simple random sampling technique was used to enrolled participants in this study.

Statistical Analysis:

Collected data was computed and analyzed using the SPSS (statistical package for social sciences) software program; version 21. Independent 'T. test' was used for comparison (p-value of ≤ 0.05) was consider significant.

Results:

This study was designed as interventional study, conducted in Khartoum state in Radiation and Isotopes Center of Khartoum (RICK), to assess the renal functions among Sudanese patients with Acute Lymphoblastic Leukemia's Receiving Chemotherapy. The study include 50 samples, the participants included from different area in Sudan; east is 2%, west is 64%, north is 10%, south is 12%, and center is 12%, that illustrated in table (1).

The patients is divided into two groups before and after treatment, then the levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium is compared between them, and statistically showed insignificant difference between the levels of urea, creatinine, uric acid, sodium, calcium, phosphorous and magnesium (p. value less than 0.05), and the potassium is significantly increased after the treatment, that illustrated in table (2).

The patients have different classification of Acute Lymphoblastic Leukemia's (L1 is 17 with 34%, L2 is 27 with 54% and L3 is 6 with 12%), and classified into two groups (before and after treatment) for each classification of ALL, and the study showed statistically insignificant difference between the parameters for each classification of ALL (p. value less than 0.05), that illustrated in tables (3), (4) and (5).

The study include both males and females, the males were 24 with 48% while the rests 26 were females with 52%, and the levels of parameter is compared between them, and showed statistically insignificant difference (p. value less than 0.05), that illustrated in table (6).

Table (1): Show the geographical distribution of Acute Lymphoblastic Leukemia's

| Area | Frequency | Percent |
|--------|-----------|---------|
| East | 1 | 2 |
| West | 32 | 64 |
| North | 5 | 10 |
| South | 6 | 12 |
| Center | 6 | 12 |
| Total | 50 | 100 |

Table (2): Comparison of renal parameters among study population before and after treatment

| Parameters | Before (Mean \pm SD) | After (Mean \pm SD) | p. value |
|------------|------------------------|-----------------------|----------|
| Urea | (49.54 \pm 28.423) | (50.28 \pm 36.913) | 0.538 |
| Creatinine | (1.730 \pm 0.964) | (2.388 \pm 2.120) | 0.717 |
| Uric Acid | (3.160 \pm 1.152) | (5.744 \pm 2.594) | 0.078 |
| Calcium | (8.645 \pm 0.906) | (10.803 \pm 12.280) | 0.182 |
| Phosphorus | (4.860 \pm 1.088) | (4.378 \pm 0.905) | 0.675 |
| Magnesium | (2.042 \pm 0.367) | (1.869 \pm 0.377) | 0.834 |
| Sodium | (124.500 \pm 9.729) | (134.340 \pm 7.058) | 0.069 |
| Potassium | (2.688 \pm 0.916) | (3.702 \pm 0.643) | 0.007 |

*Significance of p.value ≤ 0.05

Table (3): Comparison of renal parameters according to FAB classification (L1) among study population before and after treatment

| Parameters | L1 | | p. value |
|------------|---------------|---------------|----------|
| | (Mean ± SD) | | |
| | Before | After | |
| Urea | 20.896±25.666 | 20.195±32.915 | 0.273 |
| Creatinine | 2.434±0.970 | 2.604±1.372 | 0.296 |
| Uric Acid | 2.873±0.988 | 3.809±2.345 | 0.345 |
| Calcium | 5.072±3.141 | 4.962±3.030 | 0.247 |
| Phosphorus | 5.276±1.172 | 3.397±1.215 | 0.256 |
| Magnesium | 2.388±0.772 | 2.302±0.835 | 0.393 |
| Sodium | 123.17±10.038 | 54.381±64.883 | 0.209 |
| Potassium | 2.627±0.883 | 3.770±0.698 | 0.237 |

*Significance of p.value ≤ 0.05

Table (4): Comparison of renal parameters according to FAB classification (L2) among study population before and after treatment

| Parameters | L2 | | p. value |
|------------|---------------|--------------|----------|
| | (Mean ± SD) | | |
| | Before | After | |
| Urea | 18.846±25.660 | 25.008±34.19 | 0.248 |
| Creatinine | 2.106±1.089 | 2.252±1.22 | 0.386 |
| Uric Acid | 2.757±0.953 | 3.935±2.248 | 0.234 |
| Calcium | 5.276±3.121 | 5.495±3.230 | 0.267 |
| Phosphorus | 3.463±1.243 | 3.854±1.322 | 0.378 |
| Magnesium | 2.393±0.756 | 2.921±0.875 | 0.390 |
| Sodium | 55.269±60.884 | 53.269±61.34 | 0.345 |
| Potassium | 2.614±0.857 | 2.845±0.925 | 0.234 |

*Significance of p.value ≤ 0.05

Table (5): Comparison of renal parameters according to FAB classification (L3) among study population before and after treatment

| Parameters | L3 | | p. value |
|------------|-------------|------------|----------|
| | (Mean ± SD) | | |
| | Before | After | |
| Urea | 66.8±46.11 | 48.3±36.94 | 0.224 |
| Creatinine | 1.56±0.886 | 2.32±1.24 | 0.324 |
| Uric Acid | 3.45±1.21 | 6.18±2.75 | 0.237 |
| Calcium | 8.78±0.97 | 8.17±0.90 | 0.376 |
| Phosphorus | 5.73±1.05 | 3.90±1.16 | 0.073 |
| Magnesium | 2.13±0.40 | 2.02±0.57 | 0.083 |
| Sodium | 119.8±11.2 | 137.5±4.68 | 0.715 |
| Potassium | 2.62±0.85 | 3.88±.519 | 0.291 |

*Significance of p.value ≤ 0.05

Table (6): Comparison of renal parameters among study population before and after treatment according

| Parameters | Male | | Female | | p. value |
|------------|-------------|----|---------------|----|----------|
| | Mean ± SD | No | Mean ± SD | No | |
| Urea | 53.85±30.92 | 24 | 47.461±27.598 | 26 | 0.563 |
| Creatinine | 1.490±0.941 | 24 | 1.900±0.973 | 26 | 0.716 |

| | | | | | |
|-------------|---------------|----|----------------|----|-------|
| Ureic Acid | 3.140±1.351 | 24 | 3.134±1.020 | 26 | 0.262 |
| Calcium | 8.340±0.755 | 24 | 8.124±0.679 | 26 | 0.676 |
| Phosphorous | 4.700±1.257 | 24 | 5.038±0.915 | 26 | 0.297 |
| Magnesium | 1.987±0.409 | 24 | 2.069±318 | 26 | 0.978 |
| Sodium | 125.347±8.386 | 24 | 123.615±11.031 | 26 | 0.055 |
| Potassium | 2.656±1.021 | 24 | 2.699±0.852 | 26 | 0.421 |

*Significance of p.value \leq 0.05

Discussion:

The present study was carried out to estimate the serum levels of urea, creatinine, uric acid, sodium, potassium, calcium, phosphorous, and magnesium among patients with Acute Lymphoblastic Leukemia's in Radiation and Isotopes Center of Khartoum (RICK), during the period from February 2020 to January 2021; 50 samples were included, the participants involved including both males and females, the males were 24 with 48% while the rests 26 were females with 52% from different area in Sudan (east is 2%, west is 64%, north is 10%, south is 12%, and center is 12%), have different classification of Acute Lymphoblastic Leukemia's (L1 is 17 with 34%, L2 is 27 with 54% and L3 is 6 with 12%).

The present study showed statistically insignificant difference between the mean of the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, and magnesium in the study groups before and after treatment (P-value $>$ 0.05), and showed statistically significant difference between the mean of the serum level of potassium in study groups before and after treatment (P-value $<$ 0.05), the serum levels of potassium is low in the study group before treatment when compared with study group after treatment. That illustrated in table 1. The result of potassium, phosphorous and magnesium is agree with study conducted by (Alea F. Salman, 2013) ⁽¹²⁾, and disagree in the result of sodium and calcium, and the results of urea and creatinine are disagree with the study conducted by (Aisha Alhejazy Abdalla, 2018) ⁽¹³⁾ which concluded is increased.

The results of this study showed statistically insignificant difference between the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium in the study groups before and after treatment according to gender and FAB classification (P-value $>$ 0.05). That illustrated in tables 2, 3, 4 and 5.

The treatment of leukemia (imatinib) may cause hypocalcaemia and hypophosphatemia ⁽¹⁴⁾. "Hyponatremia has been estimated to occur in approximately 10% of patients with acute hematologic malignancies" ⁽¹⁵⁾, "patients with acute leukemia was significantly lower in potassium when compared with healthy controls" ⁽¹⁶⁾, "hypermagnesuria occurs in approximately 15% in acute leukemia patients" ⁽¹⁷⁾, "hypophosphatemia may be multifactorial in patients with a hematologic malignancy" ⁽¹⁸⁾, "in leukemia, hypercalcemia can occur in the acute and chronic phases" ⁽¹⁹⁾.

Conclusion:

This study concluded that the serum level of potassium is increased after the treatment, and the treatment of ALL not affect the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, and magnesium.

The gender of patients not affect the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium before and after treatment, also the FAB classification of ALL have no effect of the serum levels of urea, creatinine, uric acid, sodium, calcium, phosphorous, potassium and magnesium before and after treatment.

Recommendations:

The serum level of potassium should be estimated during the treatment. Further studies should be carried out with larger sample size and including the patient diet.

Ethical Approval and consent :

Ethical approval was obtained from ethical committee of Al Yarmouk College and Informed consent was taken from administrative.

References:

1. Fraumeni, E. and Demers, L.M., 1986. Beta2-microglobulin. Chicago, Ill.: ASCP Check Sample Continuing Education Program. Clinical Chemistry.
2. Ondreyco, S.M., Kjeldsberg, C.R., Fineman, R.M., Vaninetti, S. and Kushner, J.P., 1981. Monoblastic transformation in chronic myelogenous leukemia: presentation with massive hepatic involvement. *Cancer*, 48(4), pp.957-963.
3. Bruguera, M. and Miquel, R., 2007. The Effect of Haematological and Lymphatic Diseases on the Liver. *Textbook of Hepatology: From Basic Science to Clinical Practice*, pp.1662-1670.
4. Abdalla, A.A. and Akasha, R., 2018. Liver Enzymes, Urea and Creatinine among Acute Lymphocytic Leukemia in Sudanese Patients. *Journal of Medical and Biological Science Research*, 4(2), pp.72-75.
5. Litten, J.B., Rodríguez, M.M. and Maniaci, V., 2006. Acute lymphoblastic leukemia presenting in fulminant hepatic failure. *Pediatric blood & cancer*, 47(6), pp.842-845.
6. Braunwald, E., Fauci, A.S., Kasper, D.L., Hauser, S.L., Longo, D.L. and Jameson, J.L., 2001. *Harrison's principles of internal medicine*. McGraw Hill.
7. Shimizu, T., Tajiri, T., Akimaru, K., Arima, Y., Yokomuro, S., Yoshida, H., Mamada, Y., Taniai, N., Mizuguchi, Y., Kawahigashi, Y. and Naito, Z., 2006. Cholecystitis caused by infiltration of immature myeloid cells: a case report. *Journal of Nippon Medical School*, 73(2), pp.97-100.
8. Munker, R., Hill, U., Jehn, U., Kolb, H.J. and Schalhorn, A., 1998. Renal complications in acute leukemias. *Haematologica*, 83(5), pp.416-421.
9. Lommatzsch, S.E., Bellizzi, A.M., Cathro, H.P. and Rosner, M.H., 2006. Acute renal failure caused by renal infiltration by hematolymphoid malignancy. *Annals of diagnostic pathology*, 10(4), pp.230-234.
10. American Cancer Society (2010). "How is Leukemia Diagnosed?". Detailed Guide: Leukemia-Adult Chronic. American Cancer Society. Archived from the original on 5 April 2010. Retrieved 4 May 2010.
11. Ross JA, Kasum CM, Davies SM, Jacobs DR, Folsom AR, Potter JD (2002). "Diet and risk of leukemia in the Iowa Women's Health Study". *Cancer Epidemiol. Biomarkers Prev.* 11 (8): 777-81. PMID 12163333. Archived from the original on 10 September 2017.
12. Salman, A.F., Ali, K.F. and Alwan, A.F., 2013. Evaluation of Electrolytes in Adult Patients with Acute Leukemia before and after Chemotherapy. *Baghdad Science Journal*, 10(2), pp.362-367.
13. Abdalla, A.A. and Akasha, R., 2018. Liver Enzymes, Urea and Creatinine among Acute Lymphocytic Leukemia in Sudanese Patients. *Journal of Medical and Biological Science Research*, 4(2), pp.72-75.
14. Bassam Francis Matti, Shahla'a Fadhil Sabir, Maysaa Ali Abdul Khaleq, Marwah Hasan Al kaabi, (2017). Serum Calcium and Phosphate levels in Patients with Chronic Myeloid Leukemia Taking Different Dose of Tyrosine Kinase Inhibitors. *AJPS*, 2017, Vol. 17, No.1; 12-6.
15. Filippatos TD, Milionis HJ, Elisaf MS. Alterations in electrolyte equilibrium in patients with acute leukemia. *Eur J Haematol.* 2005;75(6):449-460.
16. Lantz B, Carlmark B, Reizenstein P. Electrolytes and whole body potassium in acute leukemia. *Acta Med Scand.* 1979;206(1-2):45-50.
17. Milionis HJ, Bourantas CL, Siamopoulos KC, Elisaf MS. Acid-base and electrolyte abnormalities in patients with acute leukemia. *Am J Hematol.* 1999;62(4):201-207.
18. Zamkoff KW, Kirshner JJ. Marked hypophosphatemia associated with acute myelomonocytic leukemia. Indirect evidence of phosphorus uptake by leukemic cells. *Arch Intern Med.* 1980;140(11): 1523-1524.
19. Noguchi M, Oshimi K. Extensive bone marrow necrosis and symptomatic hypercalcemia in B cell blastic transformation of chronic myeloid leukemia: report of a case and review of the literature. *Acta Haematol.* 2007;118(2):111-116.