

Original Research Article

Study of Phytoplankton Abundance and Species Diversity in Shahjangi Pond of Bhagalpur District, Bihar (India)

Comment [IB1]: Remove the parenthesis

Abstract

In the present study, species diversity and phytoplankton abundance were examined in Shahjangi Pond in the Bhagalpur region of Bihar. The present study was carried out from April 2021 to March 2022. Phytoplankton is considered a type of microscopic plankton capable of a photosynthesis process that is found as an essential element of the aquatic ecosystem. Phytoplankton can range in size and shape and they are photosynthesizing autotrophic organism. The primary goal of this study is to learn more about the variety of phytoplankton in this pond. A blotting silk net was used to capture the sample. The planktons were collected in wide, thick bottles with a 200ml capacity. The vial containing the filtrate was then brought to the lab. In the laboratory, 5ml of 4% formalin solution was added to the collected filtrate. Under a compound microscope, sample analysis was carried out at 10x and 40x magnification.

Comment [IB2]: Rephrase this statement.

Comment [IB3]: Full stop is missing

Comment [IB4]: Rephrase this statement. "... Is to determine/examine..."

Comment [IB5]: Write in full. Laboratory,

Comment [IB6]: Comma is missing

Comment [IB7]: To the

Comment [IB8]: Rephrase this statement and briefly highlight key findings in the abstract.

Keywords: *Chlorophyceae, Myxophyceae, Phytoplankton, Pond, Species*

Introduction

Phytoplankton is a very important character of an aquatic ecosystem and plays a crucial role in the energy transfer among the aquatic food chains or food webs. Freshwater phytoplankton plays a vital role in aquatic ecosystem like ponds, lakes and reservoirs (Sant Manickam et al., 2014). Phytoplankton is the chief primary producer in almost all aquatic systems and is an imperative food source for other organisms (Sukumaran et al, 2008). Zooplankton feeds on phytoplankton. They are accountable for eating millions of little algae and other phytoplankton. Phytoplankton not only serves as nutrition or food or diet for the

aquatic animal but also plays a significant role in keeping the biological equilibrium and quality of water (Pandey et al., 1998).

Material & Method

Sample Collection

For collection of water sample, 3 stations were cast in the pond. Water samples for the Phytoplankton study were collected from three different stations. Monthly collections were performed between 8:00 AM to 9:00 AM at the three stations when weather conditions allowed sampling from the Pond. There was surface sampling at stations 1, 2 and 3. At the station, samples were collected at mid-depth (9 m) and the bottom depth (19 m). All the collected samples were transported to the laboratory. Plankton sampling was carried out by using the plankton net of mesh size 55 µm by hauling horizontally for five meters according to the method.

Comment [IB9]: Remove this statement

Filtered water samples were stored in the sample bottle and then preserved with Lugol's iodine solution of 10%. Identification of the phytoplankton species was according to Botes (2003) Emi and Andy (2007).

Comment [IB10]: Buttress this point by highlighting key steps in a clear description and reproducible manner

H' - Shannon-Wiener diversity index

S - Species richness (number of species), p_i

Comment [IB11]: Explain the rationale behind this

$$H' = - \sum_{i=1}^k p_i \log p_i$$

Result & Discussion

Four families (Bacillariophyceae, Chlorophyceae, Chrysophyceae, and Myxophyceae) and 18 phytoplankton species were found after the six-month survey. Flagellaria recorded the highest with a percentage abundance of 28.49% followed by Ankistrodesmus with 22.08%, Aphanocapsa with 21.27% while Eudorina with 0.24% is the least abundant (Table 1). Tables 2 and 3 of the study revealed the Shannon-Wiener Diversity Index (H') and the Spatial Variation in the Diversity indices of the study.

Comment [IB12]: Italicize class names

Comment [IB13]: Italicize this name

Comment [IB14]: Same as above

Comment [IB15]: Same as above

Comment [IB16]: Same as above

Species	Total Abundance	% Abundance
<i>Flagilaria</i>	113217	28.49
<i>Tabellaria</i> sp.	13535	3.40
<i>Naviculales</i>	4121	1.03
<i>Nituschia</i>	2452	0.61
<i>yclotella</i>	13349	3.36
<i>Ankistrodesmus</i>	87791	22.08
<i>Chlorella</i>	19723	4.96
<i>Ulothrix</i>	4523	1.13
<i>Enteromorpha</i>	1009	0.25
<i>Closterium</i>	5145	1.29
<i>Eudorina</i>	1030	0.24
<i>Oocystis</i>	11458	2.88
<i>Zugrema</i>	1523	0.38
<i>Microspora</i>	5121	1.28
<i>Aphanocapsa</i>	84521	21.27
<i>Anabaena</i>	12456	3.13
<i>Oscillatoria</i>	1321	0.33
<i>Aphanizomenon</i>	8452	2.12
<i>Mallomonas</i>	6521	1.64
Total	= 397268	

Table 1. Species composition and abundance of phytoplankton in Shahjangi Pond (Bhagalpur)


Comment [IB17]: The title should be on top of the table, please

	Site		
	A	B	C
Shannon-Wiener Diversity Index (H')	3.12	2.57	3.09
Evenness Index (E)	0.75	0.73	0.76
Simpson's Density Index (D)	0.16	0.19	0.20

Table 2. Spatial variation in diversity indices of phytoplankton population across the study sites

Comment [IB18]: The title of the table should be placed on top of the table

fig.1 Graphical Representation of abundance of phytoplankton in Shahjangi Pond (Bhagalpur)



Comment [IB19]: The title of the figure should be below the figure

	Site		
	A	B	C
Shannon-Wiener Diversity Index (H')	3.12	2.57	3.09
Evenness Index (E)	0.75	0.73	0.76
Simpson's Density Index (D)	0.16	0.19	0.20

Table 3. Spatial variation in diversity indices of phytoplankton population across the study

sites

Comment [IB20]: The title should be on top of the table

fig.2 Graphical representation of Spatial variation in diversity indices of phytoplankton population across the study sites



Comment [IB21]: The title of the figure should be below the figure

Table 1 and 2 of the study revealed the Shannon-Wiener Diversity Index (H') and the Spatial Variation in the Diversity indices of the study. The Shannon-Wiener Diversity Index (H') ranged between 2.57 – 3.12 across the three study sites. Pielou Evenness Index (E) ranged between 0.73 - 0.76; Simpson's Diversity Index (D) ranged between 0.16 - 0.20. The study supports Azma's results that a Simpson Index value of 0.83 to 0.93 suggests that the communities are mature and stable since the dominance is shared by a sizable number of individuals. According to Azma and Frutos et al., pointed out that if the Pielou Index values are less than 0.5, it could be an indicator of the presence of ecological stress.

Comment [IB22]: Coma and year missing

Comment [IB23]: year

It is generally accepted that the dynamics of water circulation, nutrient concentrations, rainfall patterns, location, and the physical environment—which varies primarily in accordance with the dry and wet seasons in tropical waters—are the primary factors driving seasonal variation of phytoplankton community structure (Fonge et al., 2012 & Jeje et al., 1986). During the six-month research period, more *Ankistrodesmus* (22.08%) and *Flagilaria* (28.49%) cases were noted. Nineteen (19) phytoplankton species in total were discovered during the study period. In general, the three sites' plankton species compositions were comparable. This is comparable to the findings of Mohammed et al., 2009 & Anago et al., 2013, who reported phytoplankton and zooplankton abundance in a study of phytoplankton diversity from Koil Coastal waters in India and Awba Reservoir Ibadan Nigeria respectively. Furthermore, a study conducted by Fonge et al. on Phytoplankton diversity and abundance in the Ndop wetland plain, Cameroon showed a high diversity of phytoplankton in study areas. Anthropogenic activities at Station A may be to blame for the high quantity of phytoplankton there. The results support those of Fonge et al., who noted that the ongoing input of nutrients, particularly through the usage of nitrate and phosphate fertilizers adjacent to the station, may also be responsible for the high abundance of phytoplankton species. All sites showed an increase in species richness, variety, and evenness. This might be explained by more microhabitats emerging as a result of increasing living space.

Comment [IB24]: Year and space

Comment [IB25]: year

Conclusion

In conclusion, it can be said that the phytoplanktonic communities of Shajangi pond are mature and stable with great diversity. The study has been made throughout the year so season and geography both have an impact on

phytoplankton diversity and abundance, but despite this, the diversity of phytoplankton has remained stable.

Additional research should be done to assess the physicochemical and phytoplankton abundance and composition in various water bodies throughout the year.

References

Comment [IB26]: Correct Reference format

- ❖ Anago IJ, Esenowo IK, Ugwumba AAA. The physico-chemistry of Awba reservoir university of Ibadan, Nigeria. *Research Journal of Environmental and Earth Sciences*. 2013;5(11):638-644.
- ❖ Azma HI, Siti AZ. A comparative study of zooplankton diversity and abundance from three different types of water body. 2nd International Conference on Agriculture, Environment and Biological Sciences (ICAEB'S'15), Bali (Indonesia); 2015.
- ❖ Botes L. Phytoplankton identification catalogue – Saldanha Bay, South Africa. *Glo Ballast Monograph Series No. 7*. IMO London. Boyd, C.E. and Tucker CS. (1998). *Pond Aquaculture Water Quality Management* Lower Academic Publishers. London. 2003;15-20.
- ❖ Emi Y, Andy G. Phytoplankton identification guide. University of Georgia Marine Education Centre and Aquarium; 2007.
- ❖ Fonge BA, Tening AS, Egbe EA, Yinda GS, Fongod AN, Achu RM. Phytoplankton diversity and abundance in Ndop wetland plain, Cameroon. *African Journal of Environmental Science and Technology*, 2012;6(6):247-257.
- ❖ Fonge BA, Tening AS, Egbe EA, Yinda GS, Fongod AN, Achu RM. Phytoplankton diversity and abundance in Ndop wetland plain, Cameroon. *African Journal of Environmental Science and Technology*, 2012;6(6):247-257.

- ❖ Frutos SM, Poi de Neiff AS, Neiff G. Zooplankton abundance and species diversity in two lakes with different trophic states (Corrientes, Argentina). *Acta Limnol. Bras.* 2009; 21(3): 367–375.
- ❖ Jeje CY, Fernando R. A practical guide to the identification of Nigerian zooplankton (Cladocera, Copepoda and Rotifera). Kainji Lake Research Institute, Niger. *Journal of Limnology.* 1986;50(3):142-149.
- ❖ Mohamed AS, Thurumaran G, Arumugam R, Ragupathi RK, Anantharaman P. Studies of plankton diversity from koi coastal waters, India. *Global Journal of Environmental Research.* 2009;3(2):118-125.
- ❖ Pandey. Ushapandey Tyagi, H.R. and Rai, N., Algal flora and physico-chemical environment of Fateh saker lake. *Phykos,* 37(187): 29-30 (1998).
- ❖ SantManickam N, Saravana Bhavan P, Santhanam hanam P, Muralisankar T, Srinivasan V, P, Muralisankar T, Srinivasan V, S, Vijayadevan K, Chitrrasu P Radhakrishnan and Jawahar Ali A (2014). Seasonal Variations and Jawahar Ali A (2014). Seasonal Variations of Zooplankton Diversity in a Perennial of Zooplankton Diversity in Reservoir at Thoppaiyar, Dharmapuri District, South India. *Austin Journal of Aquaculture and Marine Biology;* (1): 1- 7.
- ❖ Sukumaran, M. Brintha, M. Mathavan Pillai, M., Species composition and diversity of phytoplankton of Pechparai dam, India. *J of Theor. and Expl. Bio.* 4(4): 157-161 (2008).

Comment [IB27]: Correct the format