

Original Research Article

PERFORMANCE OF BROILER FED DIETS CONTAINING GRADED LEVEL OF CASSAVA (*Manihot esculentus*) FLOUR MEAL AT STARTER PHASE

ABSTRACT

The experiment was conducted in the Federal University of Agriculture Zuru Kebbi State in the Department of Animal Science to determine the performance of broiler starter fed diet contain graded level of cassava (*Manihot esculentus*) flour meal (CFM 0%, 20% and 30%), diet tagged T1, T2 and T3 were compounded respectively. In a four weeks experiment, using one hundred and fifty (150) Ross 308 day old chicks were used in the experiment group, replicated five times each treatment had 10 chicks per treatment group. Feed intake and body weight gain of the bird were determine and compared. Result of final body weight, feed intake g/b/d, body weight gain and feed conversion ratio were significantly different ($P < 0.05$) for all the treatment while mortality and feed intake g/b showed that there was no significant difference ($P > 0.05$). It is concluded that broilers fed 20% cassava flour meal performed better in the study area.

Key words:- Performance, Broiler starter, Diets, Cassava,

INTRODUCTION

A serious competition exists between the feed industry and other channels in the food chain (especially man) over convention feed ingredients such as maize and Soybeans. This has resulted in the high cost and scarcity of this conventional feedstuff. Poultry feed producers are thus faced with the task of finding alternatives feed stuff that will not compromise quality. The search for such alternatives has exercised Animal Nutritionists in Nigeria for over a decade Oke *et al.*, 2002).

Cassava has been used as an alternative energy source and its inclusion in diet for poultry has been extensively studied (Tewe and Egbunike, 2000 and Adegbola, 2001). Nigerian cassava production is by far the largest in the world, a third more than production in Brazil and almost double the production in Indonesia and Thailand. The Food and Agriculture Organization of the United Nations estimated 2002 Cassava production in Nigeria to be approximately 34 million tonnes (FAO, 2004). The trade of cassava production reported by the Central Bank of Nigeria put the highest estimate of production at 37 million tons in 2000 by the Federal Ministry of Agriculture and Natural Resources (FMARD, 1997). The third series provided by (PCU, 2003) had the most conservative estimate of production at 28 million tons in 2002. Nigeria's production was targeted at 40 million tonnes by 2005 and 60 million tonnes by 2020. International Trade Administration (ITA, 2002) Nigeria; being the world's largest producer of cassava should be able to utilize its vast potential to provide enough of the tuber, not only for human consumption, but also for animal feed, it can be used as flour after being peeled or chipped with the peels and ground before use. Studies have shown that levels as high as 40% could be used in broiler diet (Erurbetina and Afolami, 1992).

The greatest limitation to the use of cassava for livestock feed is its content of cyanogenic-glucosides, linamarin and lotaustralin. Toxicity of cassava is caused by hydrocyanic acid (HCN) which is liberated when glucoside is hydrolysed by the action of linamarase enzyme. The degree of toxicity depends upon the variety, ecological condition for growth of the plant. The form of the product being fed and its processing technology (Coursey, 1999). The normal range of hydrocyanic acid in fresh cassava root is 15-400 ppm. It has long been established that the peel contains 5-10 times prussic acid of the content of the pulp. Where the hydrocyanic acid is below

100pm, as in cassava flour or chips cassava can be safely incorporated into rations as is allowed in the EEC (Delange and Ahuwalia, 2002).

The aim of this is to determine the performance of broiler fed diets containing cassava as energy supplement

MATERIALS AND METHODS

Study Area

This research work was conducted in the Department of Animal Science, Federal University of Agriculture Zuru, the area is geographically located in the guinea savanna on the latitude $11^{\circ}405N$ and longitude $5^{\circ}2.39E$ of the equator (Baba *et al.*, 2013).

Experimental Birds

The bird used for this experiment was sources from Zarm Farm in Oyo State in Nigeria. The strains used are Ross 308 broiler and was purchased at day old. One hundred and fifty (150) day old broiler chicks were used for this research.

Experimental Diets

The experimental diets used for the research was formulated using the available feed ingredients such as maize, cassava flour meal, soya bean, groundnut cake, bone meal, vitamin-mineral premix, methionine, lysine and salt. The diet was fed to the birds throughout the experiment the gross and chemical compositions of the diets are shown in table 1 below.

Table 1: Gross and chemical composition of the experimental diets.

Ingredients (%)	T ₁ (0%)	T ₂ (20%)	T ₃ (30%)
Maize	60.00	40.00	30.00
Cassava Flour Meal	-	20.00	30.00
Palm Oil	-	0.30	0.40
Soybeans Meal	2.70	5.00	3.30

Groundnut Cake	25.50	26.00	27.50
Wheat Offal	3.60	0.50	0.20
Fish Meal	0.20	0.40	0.50
Blood Meal	4.00	4.10	5.00
Bone Meal	2.00	1.7	1.70
Limestone	1.00	1.00	0.50
Lysine	0.25	0.25	0.25
Methionine	0.25	0.25	0.25
Premix	0.25	0.25	0.25
Common Salt	0.25	0.25	0.25
Total	100.00	100.00	100.00
Calculated chemical composition of the experimental diet `			
Energy ME(kcal/kg)	2993.00	2993.00	2993.00
Crude Protein (%)	22.60	22.60	22.60
Lysine (%)	1.00	1.00	1.00
Methionine (%)	0.50	0.50	0.50
Calcium Ca (%)	1.00	1.10	1.00
Phosphorus (%)	0.5	0.50	0.50

Experimental Design

One hundred and fifty (150) day old broiler chicks were used for the experiment. They were randomly divided into three treatment groups of 50 chicks each. Each treatment was further being divided into 5 replicates, with ten chicks per replicate. In a completely randomize design. Each treatment was fed one of the experimental diets for four weeks.

Data Collection

Feed intake was recorded daily by subtracting the amount offered from the left over of the previous day. Body weight was monitored weekly and mortality was recorded as it occurs.

Statistical Analysis

The data collected was subjected to analysis of variance (ANOVA) using the Stat View Statistical Package (SAS, 2003).

RESULTS AND DISCUSSION

At the end of the experiment, broiler chicks feed diet with the highest level of cassava flour meal supplement (T₃) consumed less feed (673.82g/b) compared to those on the other treatment (800.00 and 717.04g/b) for T₁ and T₂ respectively (P>0.05) (Table 2). Similarly, birds on the highest level of cassava flour meal (T₃) recorded the lowest body weight gain (183.14g/b) compared to the value recorded for T₁ and T₂ (232.21 and 250.77g/b) respectively. The final body weight (g/b) differ significantly between the treatment (P<0.05) even though the highest value was recorded for T₁ (290.71) as against 288.00 (g/b) and 234.00 (g/b) for T₂ and T₃ respectively. Feed conversion ratio (FCR) differ significantly between the treatment (P<0.05) even though the best value was recorded for T₂ (2.89) as against 3.58 and 3.49 for T₁ and T₃ respectively. Incidence of mortality did not vary between the treatment (P>0.05) even though the highest value was recorded for T₁ (3.00) as against 1.60 and 2.60 for T₂ and T₃ respectively (Table 2).

Table 2: Performance of broiler fed diets containing cassava flour meal at starter phase

Parameter	T1	T2	T3	SEM
Initial body weight	58.50	59.76	58.85	1.48
Final body weight (g/b)	332.71 ^c	288.00 ^b	234.00 ^a	13.27
Feed intake (g/b)	2561.32	717.04	673.81	546.67
Feed intake (g/b/d)	34.90 ^b	25.60 ^a	24.06 ^a	1.67

Body weight gain (g/b)	274.21 ^b	250.76 ^b	183.14 ^a	10.78
Body weight gain g/b/d)	9.79 ^b	8.79 ^b	6.52 ^a	0.40
Mortality (%)	3.00	1.60	2.60	1.21
Feed conversion ratio	3.58 ^b	2.89 ^a	3.49 ^b	0.54

^{abc} Means in the same row followed by different superscripts are significant difference ($P < 0.05$)

From table 2.0 Results on body weight gain for chick on the control group was significantly ($P < 0.05$) better compared to chicks on others treatment diets. However, feed intake for chick on all treatment diets differ significant ($P < 0.05$) during the starter phase this is in tandem with the work of Erurbetine and Afolani (2001) that showed a progressive decline in body weight gain of birds increasing concentration of cassava meal. It also agreed with earlier report of Akinfala (2000), who recommended inclusion level of up to 25% cassava root in poultry rations. Stevenson and Jackson (1983) reported that level rate of up to 30 percent of cassava in the diet improved growth performance of poultry, whereas (Erurbetine and Afolami, 1992). observed decrease in the weight of poultry resulting from the increase in the quantity of cassava included in the diets. This research showed that birds on 0% CFM inclusion level had the highest body weight gain than the rest of the treatments. This supports the findings of (Adegbola, 2001) who reported a significant difference in weight gain of starter broilers fed cassava meal.

The non-significant effect ($P > 0.05$) of cassava flour meal levels in feed intake of chicks conformed to the work of (Erurbetine and Afolami, 1992). who reported similar non-significant effects of this parameter. Stevenson and Jackson (2003), reported similar findings when they fed up to 30% cassava flour meal without any significant effect on the consumption index of

broilers. However, (Tewe and Egbunike, 2000) reported that feed consumption was affected when cassava level of 30% was fed to young chicks.

The mortality rate for this work falls within the accepted range of 0-5% for good management practices and safety of diets as advocated by ISA (1996).

The result of the feed conversion ratio of chicks is significantly affected ($P < 0.05$). This is similar with the report of (Erurbetine and Afolami, 1992). who reported significant effect on feed conversion ratio among birds, fed dietary cassava meal. (Tewe and Egbunike, 2000 and Adegbola, 2001) reported that the feed conversion ratio of broilers fed sundry cassava flour meal was poor when compared to those on maize meal. It is also similar with the finding of (Adegbola, 2001) who reported a significant difference on feed conversion efficiency when cassava meal was added to the diets of starter broilers at graded levels.

CONCLUSION AND RECOMMENDATIONS

It is concluded that broilers fed 20% cassava flour meal performed better in most parameters

It is also recommended that further investigation should be carried but to determine the hematological parameters and proximate analysis of shea butter cake. Feeding trials should be repeated to ascertain the passive impact of CFM on performance characteristics. Lower levels of CFM should be tried.

REFERENCES

Adegbola, A.A. (2001). Methionine as an additive to cassava based diets. In:Nestle, B. and Graham, M. (eds). Cassava as an animal feed. Proceedings of a workshop held at the University of Guelph, 18-20 April 2000. Ottawa International Development Research Center, IDRC

095C. pp. 9-17.

Baba M.D., J.M Yelwa, J.S Dabai and A.M. Sakaba., (2013), published paper on the

economics of goat production in Zuru, Kebbi State Nigeria. College of Agriculture, Zuru
Pp 543-546.

Oke, O.L. (2002). Problems in the use of cassava as animal feed. *Animal Feed Science and Technology* 3:345-380.

Tewe, O.O., and Egbunike I. (2000). Utilisation of Cassava in Non-Ruminant Livestock Feeds in: Cassava as Livestock feed in Africa proceedings of IITA/ILCA/University of Ibadan Workshop on the Potential Utilisation of cassava as livestock Feed in Africa (Eds S K Hahn, L Reynolds and G N Egbunike), Pp 28-38.

Eruvbetine, D. and Afolami, C.A. (1992). Economic evaluation of Cassava (*Manihot esculenta*) as a feed ingredient for broilers. Proc. Of The sixth world's poultry congress, Netherlands. pp. 532-535.

F.A.O. (2004) Food and Agricultural Organization of the United Nations *African agricultural, the next 25 years*, Rome, Italy. pp.22-29.

FMARD, (1997). Department of Planning, Research and Statistics. Nigerian
Agricultural Statistics. Federal Ministry of Agriculture and Rural Development, Abuja.

PCU, (2003). Crop Area Yield Survey. Abuja Nigeria: Federal Ministry of
Agriculture and Rural Development: Project Coordinating Unit. Pp. 10 - 11

ISA (Institut de sélection Animale) (1996). Guide d'Élevage des poulets de chair service production et d'Assistance Technique, Quintin, France. Pp. 3-10.

IITA, (2002). Nigerian's Cassava Industry: Statistical Handbook.

Coursey, D.G. (1999). Cassava as Food. Toxicity and technology. In:
chronic cassava toxicity. Nestel, B. and P. Maclyntre (Eds).
Proceedings of an Inter disciplinary Workshop. London, pp. 29-30

Delange, E., and Ahluwalia R., (2002). Cassava toxicity and thyroid:

Research and public health Issues. Proceedings, International Workshop on Cassava Toxicity, 31 May -2 June 1982, Ottawa, Canada. IDRC-207e: Ottawa.

Akinfala, E. O. (2000). Chemical and biological evaluation of whole cassava plant-based diets in life-cycle feeding of pigs in South-Western Nigeria Ph. D thesis, University of Ibadan, Nigeria, Pp. 236-239.

Stat view, statistical package (SAS 2003). Incorporated New York.

Stevenson, M.H., and N. Jackson. (2003). The Nutritional value of dried Cassava root meal in broiler diets. *Journal of the Science of Food and Agriculture* 34: 1361-1367.

Eruvbetine, D. and Afolami, C.A.(2001). Economic evaluation of Cassava (*Manihot esculenta*) as a feed ingredient for broilers.Proc. Of The sixth world's poultry congress, Netherlands. pp. 532-535.