

Original Research Article

Polymorphism and Haplotype Structure of TLR2 Gene Associated with Endometritis Infection in River Buffalo

ABSTRACT

Background and Aim: Toll-like receptors (TLRs) are important for the recognition of pathogen-associated molecular patterns. Single nucleotide polymorphisms (SNPs) within TLRs have a potential impact on the alteration of susceptibility or resistance to inflammatory diseases. This work focused on a case-control study for the distribution of SNPs in TLR2 gene to test their role for endometritis occurrence in river buffalo.

Materials and Methods: Egyptian buffaloes from the slaughterhouse were tested, where forty samples divided into apparently healthy uteri (20n) and clinically infected ones (20n) were used to detect the genetic association between TLR2 gene SNPs and endometritis disease.

Results: Nineteen novel polymorphic sites were identified. Three SNPs were found to be statistically significant. AA genotype at 5'UTR and CC genotype at the coding region had significant association with susceptibility to endometritis, while GG genotype at 5'UTR had significant association with resistance to endometritis. Polyphen 2 analysis revealed three amino acids substitutions in TLR2 protein having potential functional significance. Haplotype reconstruction revealed the statistical significance of haplotype frequencies between cases and controls indicating its association with the occurrence of endometritis.

Conclusions: It is to be concluded that the innate immune activation response may be interfered by the effect of polymorphisms and mutations of TLRs due to the reduction of the protein ability to recognize PAMPs.

Keywords: *Bubalus bubalis*, TLR2, endometritis, SNPs

1. INTRODUCTION

Water Buffalo has been for long playing a pivotal role in livestock and agriculture economy of many developing countries across the globe. Domesticated species of buffalo are river and

swamp buffalo which are typically found in tropical and subtropical forests, wet grasslands, marshes and swamps regions of the world. The current water buffalo population is 195 million head including both river and swamp buffaloes, 77% of which are river buffalo. Egyptian buffaloes are of the river type with a population of 4 million head, it ranks the third after Indian (over 115 million) and Pakistani buffalo (31.7 million)[1] and it is a representative of the Mediterranean buffalo.

Uterine diseases related to microbial infection represent the main cause of reduced fertility in bovine; the economic losses related to these diseases are double to those caused by ovarian disorders with a great impact on the economy [2]. The incidence of uterine infections in buffaloes is higher than cows which might be related to poor hygiene, vaginal stimulation for milk let down and possibly wallowing. Buffaloes are culled and sent to slaughterhouses either because they are uneconomic to maintain or because they have some disease problems. Hence, abattoirs are a good source for studying pathological conditions of buffalo reproductive organs that are severe enough to cause infertility and even sterility [3].

Endometritis is the most common uterine disease observed in buffaloes slaughtered at abattoirs and is one of the main causes of infertility in both cattle and buffalo [4]. Endometritis disease is highly prevalent, and lead to economic losses due to decreased milk yield, medication as well as prolonged calving intervals [5]. Numerous bacteria in a variety of combinations have been isolated from infected uteri of buffaloes, where *Arcanobacterium pyogenes*, *E.coli*, *streptococcus spp.*, and *Staphylococcus spp.* are responsible for toxic metritis and endometritis in buffaloes [6]. *E.coli* are usually associated with uterine infection with other bacteria in buffaloes [7]. Furthermore, *Escherichia coli* and *Arcanobacterium pyogenes* are the most prevalent bacteria isolated from the uterine lumen of bovine with uterine disease [8].

Toll-like receptors (TLRs) are part of pattern recognition receptors (PRRs) that play an essential role in recognizing molecules that are produced by pathogens, inflammatory initiation, and stimulation of adaptive immune responses. PAMPs include various components of pathogens such as lipopolysaccharides (LPS), peptidoglycans, flagellin, bacterial DNA and viral double stranded RNA [9]. Polymorphisms within TLR genes are associated with the occurrence of wide range of diseases in human and mice. Furthermore, polymorphism in TLRs genes is also

associated with variations in disease resistance traits in livestock. TLR2 and TLR4 are the most two studied TLRs due to their essential role in the recognition of bacterial pathogens. Several studies have focused on the functional consequences of TLR2 and TLR4 SNPs where it was found that polymorphisms in TLR2 and TLR4 genes have potential effect on the binding of these TLRs to its ligands and disease resistance [10]. TLR2 gene in buffalo consists of two exons with 2355 bp open reading frame encoding 784 amino acids. The gene plays an essential role in the innate immune response to a variety of pathogens, making TLR2 polymorphism to have a potential impact on disease resistance in livestock [11].

2. MATERIALS AND METHODS

2.1 Sampling and study design:

Egyptian buffaloes at the slaughterhouse were tested, where forty samples divided into apparently healthy uteri (20n) and clinically infected ones (20n) were used to detect the genetic association between TLR2 gene SNPs and endometritis disease. Therefore, microbiological processing was performed on all uterine samples to identify bacterial pathogens in clinically infected samples.

2.2 TLR2 sequencing:

Overlapped primers were designed to amplify two exons of TLR2 gene in healthy and diseased uterine samples. DNA extraction was performed for single nucleotide polymorphisms (SNPs) analysis of TLR2, using DNeasy Blood & Tissue kit (Qiagen) according to Manufacturer's instructions. In this investigation, primers for TLR2 gene were designed using known DNA sequences of *Bubalus bubalis* published in database with accession No. HM756161 and the web interface primer3 as shown in Table (1). The TLR2 coding sequence was amplified using five overlapped primers according to the following thermal profile: initial denaturation at 95°C for 1-3 min followed by 40 cycles of 30s at 95°C, 30s at TMann(C) which is the annealing temperature specific for each primer in and 1 min at 72°C, with a final extension at 72°C for 1 min. PCR reactions were performed with PCR Master Mix (2X) kit (Thermo Fisher Scientific) which included 50ng of genomic DNA and 10 µM of each primer (forward and reverse) in a final volume of 50 µl. Amplicons were purified (QIAquick PCR Purification Kit, Qiagen) and bi-directionally sequenced by Laragen company (USA).

2.3 Genotyping and SNP selection:

CLUSTAL - W program [12] was used to analyze the forty river buffalo Egyptian breed sequences of TLR2 gene with CDs of publicly available GenBank: HM756161. A consensus sequence for each sample was generated from alignment of each forward and reverse sequence of a single DNA sample to identify polymorphisms among the samples.

2-4 Genetic association and protein analysis:

Genotype frequencies were tested for deviation from Hardy-Weinberg equilibrium (HWE) by using Fisher's Exact test, where the allelic and genotypic frequencies were carried out for each SNP independently.

Univariate logistic regression model was performed for the SNPs that were significantly associated with the disease. Odds Ratio test (OR) was calculated with 95% confidence interval (CI)[13]. All statistical analyses were performed using R statistical program and P-value was corrected using Bonferroni method [14]. Haplotype reconstruction was performed using PHASE software, version 2.1[15]. Moreover, the potential impact on protein structure and function due to amino acid substitutions was performed by PolyPhen-2 software (Polymorphism Phenotyping V2) with default parameters [16] and the colored figures are available at <http://genetics.bwh.harvard.edu/cgi-bin/ggi/ggi2.cgi>. Protein domain has been predicted from the CD sequence by using the Conserved Domain Database available at: <http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi> [17].

3. RESULTS AND DISCUSSION

3.1 Bacterial results:

Microbial infections of the genital tract cause infertility by disrupting uterine and ovarian function [18]. Uterine lumen contamination can be categorized depending on the bacterial pathogenicity [19] which is also correlated in part with the severity of postpartum endometritis; although, the establishment and persistence of uterine infection is also influenced by the genetic factors, uterine environment and the animal's innate and acquired immunity. Uterine pathogens including *Escherichia coli*, *Proteus* species, *Arcanobacterium pyogenes* and *Prevotellamelaninogenicus* are associated with greater endometrial inflammation and more severe clinical uterine disease [20].

Examination of the collected uterine swab samples was performed involving colony characters, cellular morphology and the purity of the culture and the suspected colonies were identified. The most highly isolated bacterium from diseased samples was found to be *Escherichia coli*. Other potential bacterial pathogens were also detected with different percentages including *Streptococcus pyogenes* and *S. aureus*. Opportunist contaminants like *Klebsiella spp.*, *Micrococcus species*, *Proteus spp.* and *S. epidermidis* were also identified.

Several studies demonstrated that *Escherichia coli* bacteria is one of the most common isolates in endometritis cases. Furthermore, it has been widely reported that *Escherichia coli* presence was correlated with increased endometrial inflammation and more severe clinical diseases [20 and 4]. In this respect, *E. Coli* was isolated with high incidence from Egyptian buffalo with endometritis [21]. Moreover, *E. coli* (36.66%), *Klebsiella spp* (30%), *Proteus* (13.33%) were isolated from endometritis cases of bovine [22].

3.2 SNPs identification:

Buffalo TLRs can be used as positional markers in the genome and polymorphism reported in their genes may be related to differences in resistance and susceptibility to various bacterial infections [23]. Single nucleotide polymorphisms (SNPs) within TLRs genes have a potential impact on the response of animals to TLR ligands which lead to the alteration of susceptibility or resistance to infectious and inflammatory diseases [24]. On the other hand, polymorphisms, and mutations of TLRs genes were reported to interfere with the innate immune activation due to the reduction of the protein ability to recognize PAMPs [25].

Sequence analysis of TLR2 exons in healthy samples yielded twenty-six SNPs which were bi-allelic (two in 5' UTR, 22 in coding region and two in 3' UTR), in addition to the identification of a dinucleotide SNP at genomic positions 482 and 483 bp of coding region. Sequence analysis of TLR2 exons in diseased samples yielded nineteen SNPs which were bi-allelic (two in 5' UTR, 15 in coding region and two in 3' UTR), in addition to the identification of a dinucleotide SNP at genomic positions 482 and 483 bp of coding region.

This study focused on the mutual SNPs found in healthy and diseased samples to identify SNPs of TLR2 gene associated with endometritis occurrence in Egyptian buffalo. Nineteen novel polymorphic sites were found in both samples (two in 5' UTR, 15 in coding region and two in 3'

UTR). Fourteen SNPs (73.7 %) showed transitions while five SNPs (26.3 %) showed transversions. Seven mutual SNPs, located at the coding region were non-synonymous (nsSNPs) with five showing transitions and two transversions, while eight mutual SNPs were synonymous (sSNPs) with seven transitions and one transversions. There was a mutational bias between transitions and transversions with nsSNPs and sSNPs ($p=0.017$) by using Fisher's two tailed exact test. The *TLR2* CD sequences generated from healthy and diseased samples were submitted to GenBank with the accession numbers KU984438 and KU984439, respectively.

3.3 Allele and Genotype frequencies of TLR2 mutual SNPs:

Genotype and allele frequency of each mutual SNP of *TLR2* gene were calculated in healthy and diseased samples as the first step of genetic association study using Fisher's exact test [26]. All *TLR2* mutual SNPs conformed to Hardy-Weinberg equilibrium ($P>0.05$) [27]. Three mutual SNPs were found to be statistically significant in allele and genotype frequency after applying Bonferroni correction to reduce the false positive results, one of those SNPs was located at coding region (374bp), while the other two SNPs were located at 5' UTR (90&124bp). On the other hand, there was no relationship in allele and genotype frequencies of 14,1034,1375 mutual SNPs ($P=1$). Moreover, there was no genetic association between other mutual SNPs (Table 2).

3.4 Odds Ratio test analysis of statistically significant TLR2 mutual SNPs:

Only three SNPs (90bp, 124bp located at 5' UTR and 374bp at CD) showed significant P values ($p<0.01$) in allele and genotype frequency (Table 2) were subjected for Odds Ratio test analysis after logistic regression model performance to determine the association between these SNPs and endometritis occurrence in river buffalo Egyptian breed. Three mutual SNPs in *TLR2* gene, associated with endometritis disease, were found to be located at 5' UTR and coding region, where the dominant homozygous genotypes AA and CC of 90bp and 374bp SNPs had a significant P -value (0.0015) with OR value more than one (2.88), indicating a significant association with susceptibility to endometritis. Whereas OR value of recessive homozygous genotypes of those SNPs (CC & TT) were less than one (0.297) with significant P -value (0.011) revealed association of those genotypes with resistance to endometritis. On the other hand, SNP at 124 bp showed OR value less than one (0.320) for dominant homozygous genotype (GG) with P -value (0.0002) indicating a significant association with resistance to endometritis while showed OR value more than one (3.833) with P -value (0.002) for recessive

homozygous genotype (AA) indicated the significant association with susceptibility to endometritis (Table 3).

Despite the relatively small number of samples used, the results give a rather good idea of the association between TLR2 genotypes and resistance / susceptibility to endometritis disease. Those results are in good agreement with previous results reporting that TLRs polymorphisms are associated with resistance and susceptibility to diseases in many species [28; 29 and 24].

Previous studies have demonstrated the influence of polymorphisms on susceptibility to several bacterial diseases, where *TLR2* polymorphisms were found to increase the risk of infections like tuberculosis [30], urinary tract infections [31], Mastitis [32] and other disease conditions [33]. On the other hand, the association between genetic polymorphisms and susceptibility to COVID-19 disease and severity was revealed [34]. The relation between endometritis occurrence and genetic polymorphism of TLRs in human and cattle was previously reported [35]. It is also to be mentioned that the polymorphism in the 5'UTR of TLRs genes and its role in diseases susceptibility was previously detected in bovine [36]. In this respect, several polymorphisms have been found in the *TLR2* gene, with a study demonstrating the association between *TLR-2* and the increased type 2 diabetes (T2DM) risk, since carrying the D allele of the *TLR-2* del -196-174 variant may be related as a risk factor for T2DM [37]. Another study performed a meta-analysis suggesting that TLR2 Arg753Gln polymorphism is significantly associated with high TB risk, where in the subgroup analysis based on ethnicity it was found that TLR2 Arg753Gln polymorphism elevates the risk of TB in Asian and Caucasian populations, but not in African or mixed populations [38].

3.5 Protein analysis of TLR2 CD of healthy and diseased samples:

Translation of nucleotide sequences (2352 bp) of TLR2 coding regions of healthy and diseased samples was carried out by Open Reading Frame software (ORF). Eight non-synonymous (nsSNPs) were detected in coding region of healthy samples which, leads to seven amino acids substitution, while Seven non-synonymous (nsSNPs) were detected in coding region of TLR2 of diseased samples, which leads to six amino acids substitution (Table 4).

For protein analysis, PolyPhen-2 software was used to predict the potential impact on the protein structure and function due to amino acid substitutions in TLR2 coding region of healthy and

diseased samples. The majority of amino acid substitutions were detected to be benign. Three substitutions (L 5 W, T 18 M and N 345 S), which were found within *TLR2* coding region, were found to be possibly damaging and may have a potential impact on the alteration of protein structure and function [39]. It is to be noted that the change of polarity from polar Threonine to non-polar Methionine (T 18 M) might also affect the protein structure and function [40]. This effect of change in protein function related to that of amino acid polarity was also reported in cattle *TLR* SNPs [41]. It was also previously reported that non-synonymous SNPs that modify amino acids polarity in TLRs may affect ligand binding and recognition [42]. Furthermore, amino acid substitutions that alter the amino acid polarity may also have a potential impact on host immune responses and resistance to diseases [43]. On the other hand there was no effect on protein function due to change the amino acid (583) from serine to threonine in coding region of healthy samples (Table 4).

Tables 5A & 5B show all the SNPs in diseased and healthy samples, respectively, their positions in *TLR2* CDs, the protein domain and PolyPhen-2 analysis of amino acid substitution effects of these SNPs. Four non-synonymous SNPs were detected in Leucine-rich repeats (LRR) domains of *TLR2* of diseased samples in addition to one other in healthy samples and might have a role in altering its ability to identify extracellular pathogens [44]. These results are in accordance with other studies, which reported that LRR domains are rich in non-synonymous SNPs in canine, human, porcine and bovine TLRs [41], [45] and [46]. Furthermore, polymorphisms that occur in LRR domains may cause changes in responsiveness towards pathogenic microorganisms [47].

3.6 Haplotype prediction:

The determination of haplotype reconstruction for several SNPs in one gene is a powerful tool to provide more information about genotype-phenotype associations than individual SNPs [48]. PHASE v2.1 was used to detect haplotypes for *TLR2* coding region SNPs of healthy and diseased samples. Unphased diploid data was used. The best pairs for each sample were determined as well as frequencies of each determined haplotype.

It was found that 5% of healthy samples had the potential to transmit GTTACCAGCCGCACT haplotype, while 8% of diseased samples had the potential to transmit TTTACCAGCGACGTT haplotype. Moreover, there was a statistical significance ($P=0.03$) of haplotype frequencies of

healthy and diseased samples by performing a permutation test, included in PHASE v2.1, which indicated the presence of a significant association between haplotype frequency of TLR2 gene in healthy and diseased samples with endometritis occurrence in Egyptian buffalo. This result was previously found to apply also for TLR4 gene [49]. Haplotype reconstruction of TLRs and its association with disease occurrence has been reported in human, cattle [50] and buffalo [24].

4. CONCLUSION

The innate immune activation response may be interfered by the effect of polymorphisms and mutations of TLRs due to the reduction of the protein ability to recognize PAMPs. The present study describes novel polymorphic sites in the river buffalo Egyptian breed *TLR2* gene and their association with endometritis disease occurrence. We can improve our ability to understand complex diseases by studying the genetic factors involved, which may help identifying their underlying physio-pathological pathways and therefore improving our ability to understand the disease in its entirety and to determine the risk of developing it. This will pave the way to both improvement of disease resistance in herds by selective breeding and also to identification and synthesis of innovative drugs.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- [1] Lewandowski CM. The effects of brief mindfulness intervention on acute pain experience: An examination of individual difference. 2015, Dissertation. Southern Illinois University Carbondale.
- [2] Sheldon IM, Roberts MH. Toll-like receptor 4 mediates the response of epithelial and stromal cells to lipopolysaccharide in the endometrium. *PLoS One*. 2015; 5(9): e12906.
- [3] Azawi OI, Ali AJ, Lazim EH. Pathological and anatomical abnormalities affecting buffalo cows reproductive tracts in Mosul. *Iraqi J. Vet. Sci*. 2008; 22(2): 59-67.

- [4] Ajevar G, Muthu S, Sarkar M, Kumar H, Das G K, Krishnaswamy N. Transcriptional profile of endometrial TLR4 and 5 genes during the estrous cycle and uterine infection in the buffalo (*Bubalus bubalis*). *Vet. Res. Commun.* 2014; 38(2): 171-176.
- [5] Goshen T, Shpigel NY. Evaluation of intrauterine antibiotic treatment of clinical metritis and retained fetal membranes in dairy cows. *Theriogenol.* 2006; 66(9): 2210-2218.
- [6] Azawi OI. Pathogenesis of postpartum metritis in buffaloes: a review. *Buffalo Bull.* 2013; 32(1).
- [7] Azawi OI, Omran SN, Hadad JJ. Clinical, bacteriological, and histopathological study of toxic puerperal metritis in Iraqi buffalo. *J. Dairy Sci.* 2007; 90(10): 4654-4660.
- [8] Herath S, Lilly ST, Santos NR, Gilbert RO, Goetze L, Bryant CE, White JO, Cronin J, Sheldon IM. Expression of genes associated with immunity in the endometrium of cattle with disparate postpartum uterine disease and fertility. *Reprod. Biol. Endocrinol.* 2009; 7(1): 55.
- [9] Jang JH, Shin HW, Lee JM, Lee HW, Kim EC, Park SH. An Overview of Pathogen Recognition Receptors for Innate Immunity in Dental Pulp. *Med. Inflamm.* 2015; 794143. <http://dx.doi.org/10.1155/2015/794143>
- [10] Zhu L, Yuan H, Jiang T, Wang R, Ma H, Zhang S. Association of TLR2 and TLR4 polymorphisms with risk of cancer: a meta-analysis. *PloS one.* 2013; 8(12): e82858.
- [11] Tschirren B, Andersson M, Scherman K, Westerdahl H, Mittl PR, Råberg L. Polymorphisms at the innate immune receptor TLR2 are associated with *Borrelia* infection in a wild rodent population. *Proc. Royal Society of London B: Biol. Sci.* 2013; 280(1759): 20130364.
- [12] Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 1994; 22 (11): 4673- 4680.
- [13] McHugh ML. The odds ratio: calculation, usage, and interpretation. *Bioch. Medica.* 2009; 19(2): 120-6.
- [14] Bland JM, Altman DG. Calculating correlation coefficients with repeated observations: Part 2-Correlation between subjects. *BMJ.* 1995;310(6980): 633.
- [15] Stephens JC, Schneider JA, Debra A, Tanguay DA, Julie Choi J, Tara Acharya T. Haplotype variation and linkage disequilibrium in 313 Human Genes. *Sci.* 2001; 293(5529): 489-493.
- [16] Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, Kondrashov AS and Sunyaev SR. A method and server for predicting damaging missense mutations. *Nat Methods.* 2010;7(4): 248-249.

- [17] Marchler-Bauer A, Anderson JB, Chitsaz F, Derbyshire MK, DeWeese-Scott C, Fong JH, Geer LY, Geer RC, Gonzales NR, Gwadz M, He S, Hurwitz DI, Jackson JD, Ke Z, Lanczycki CJ, Liebert CA, Liu C, Lu F, Lu S, Marchler GH, Mullokandov M, Song JS, Tasneem A, Thanki N, Yamashita RA, Zhang D, Zhang N, Bryant SH. CDD: specific functional annotation with the Conserved Domain Database. *Nucleic Acids Res.* 2009; 37: 205-210.
- [18] Galvao KN. Uterine diseases in dairy cows: understanding the causes and seeking solutions. *Anim. Reprod.* 2013; 10(3): 228-238.
- [19] Williams EJ, Fischer DP, Pfeiffer DU, England GC, Noakes DE, Dobson H, Sheldon IM. Clinical evaluation of postpartum vaginal mucus reflects uterine bacterial infection and the immune response in cattle. *Theriogenol.* 2005; 63(1): 102-117.
- [20] Bonnett BN, Martin SW, Gannon VP, Miller RB, Etherington WG. Endometrial biopsy in Holstein-Friesian dairy cows. III. Bacteriological analysis and correlations with histological findings. *Canadian J. Vet. Res.* 1991; 55(2): 168.
- [21] Hanafi EM, Ahmed WM, Abd El Moez SI, El Khadrawy HH, Abd El Hameed AR. Effect of clinical endometritis on ovarian activity and oxidative stress status in Egyptian buffalo cows. *American-Eurasian J. Agri. Environ. Sci.* 2008; 4: 530-536.
- [22] Udhayavel S, Malmarugan S, Palanisamy K, Rajeswar J. Antibioqram pattern of bacteria causing endometritis in cows. *Vet. World.* 2013; 6 (1): 100.
- [23] Banerjee P, Gahlawat SK, Joshi J, Sharma U, Tantia MS, Vijh RK. Sequencing, Characterization and Phylogenetic analysis of TLR genes of *Bubalus bubalis*. *DHR-IJBLs.* 2012; 3:137-158.
- [24] Alfano F, Peletto S, Lucibelli MG, Borriello G, Urciuolo G, Maniaci MG, Desiato R, Tarantino M., Barone A, Pasquali P, Acutis PL. Identification of single nucleotide polymorphisms in Toll-like receptor candidate genes associated with tuberculosis infection in water buffalo (*Bubalus bubalis*). *BMC genetics*, 2014; 15(1): 139.
- [25] Mariotti M, Williams JL, Dunner S, Valentini A, Pariset L. Polymorphisms within the toll-like receptor (TLR)-2,-4, and-6 genes in cattle. *Diversity.* 2009; 1(1): 7-18.
- [26] McDonald JH. Handbook of biological statistics. 2009. Sparky House Publishing.
- [27] Salanti G, Amountza G, Ntzani EE, Ioannidis JPA. Hardy–Weinberg equilibrium in genetic association studies: an empirical evaluation of reporting, deviations, and power. *Eur. J. Human Genet.* 2005; 13: 840-848.
- [28] Plantinga TS, Johnson MD, Scott WK, van de Vosse E, Edwards DRV, Smith PB, Alexander BD, Yang JC, Kremer D, Laird GM, Oosting M. Toll-like receptor 1 polymorphisms increase susceptibility to candidemia. *J. Infect. Dis.* 2012; 205(6): 934-943.

- [29] Sun Q, Zhang Q, Xiao HP, Bai C. Toll-like receptor polymorphisms and tuberculosis susceptibility: A comprehensive meta-analysis. *J. Huazhong Uni. Sci. Techno. [Med. Sci.]*. 2015; 35: 157-168.
- [30] Hawn TR, Scholes D, Li SS, Wang H, Yang Y, Roberts PL, Stapleton AE, Janer M, Aderem A, Stamm WE, Zhao LP. Toll-like receptor polymorphisms and susceptibility to urinary tract infections in adult women. *PloS one*, 2009; 4(6): e5990.
- [31] Kannaki TR, Shanmugam M, Verma PC. Toll-like receptors and their role in animal reproduction. *Anim. Reprod. Sci.* 2011; 125 (1): 1-12.
- [32] Schnetzke U, Spies-Weisshart B, Yomade O, Fischer M, Rachow T, Schrenk K, Glaser A, von Lilienfeld-Toal M, Hochhaus A, Scholl S. Polymorphisms of Toll-like receptors (TLR2 and TLR4) are associated with the risk of infectious complications in acute myeloid leukemia. *Genes and Immunity*. 2015; 16 (1): 83-88.
- [33] Dieter C, Brondani LA, Leitao CB, Gerchman F, Lemos NE, Crispim D. Genetic polymorphisms associated with susceptibility to COVID-19 disease and severity: A systematic review and meta-analysis. *PLoS One* 2022; 17(7): e0270627.
- [34] Pinedo PJ, Galvao KN, Seabury CM. Innate immune gene variation and differential susceptibility to uterine diseases in Holstein cows. *Theriogenol.* 2013; 80(4): 384-390.
- [35] Dubey PK, Goyal S, Kathiravan P, Mishra BP, Gahlawat SK, Kataria RS. Sequence characterization of river buffalo Toll-like receptor genes 1–10 reveals distinct relationship with cattle and sheep. *Inter. J. immunogenet.* 2013; 40(2): 140-148.
- [36] Karaali ZE, Candan G, Aktuğlu MB, Mustafa Velet M, Ergen A. Toll-Like Receptor 2 (*TLR-2*) Gene Polymorphisms in Type 2 Diabetes Mellitus. *Cell J.* 2019; 20 (4): 559–563.
- [37] Hu L, Tao H, Tao X, Xiaolong Tang X, Xu C. TLR2 Arg753Gln Gene Polymorphism Associated with Tuberculosis Susceptibility: An Updated Meta-Analysis. *BioMed. Res. Inter.* 2019: 1- 9.
- [38] Adzhubei I, Jordan DM, Sunyaev SR. Predicting functional effect of human missense mutations using PolyPhen2. *Current Protocols in human Genet.* 2013; 7-20.
- [39] Allen SJ, Schwartz AM, Bush MF. Effects of polarity on the structures and charge states of native-like proteins and protein complexes in the gas phase. *Anal. Chem.* 2013; 85(24): 12055-12066.
- [40] White SN, Taylor KH, Abbey CA, Gill CA, Womack JE. Haplotype variation in bovine Toll-like receptor 4 and computational prediction of a positively selected ligand-binding domain. *Proc. Nat. Acad. Sci.* 2003; 100(18): 10364-10369.

- [41] Zhang XY, Lei M, Xie L, Zhang CX, Zheng J, Yang C, Deng XD, Li JL, Huang DP, Xie XH Detection of polymorphisms and protein domain architectures in rabbit toll-like receptor 2. *World Rabbit Sci.* 2014; 22(1): 83-90.
- [42] Shinkai H, Tanaka M, Morozumi T, Eguchi-Ogawa T, Okumura N, Muneta Y, Awata T, Uenishi H. Biased distribution of single nucleotide polymorphisms (SNPs) in porcine Toll-like receptor 1 (TLR1), TLR2, TLR4, TLR5, and TLR6 genes. *Immunogenet.* 2006; 58(4): 324-330.
- [43] Seabury CM, Cargill EJ and Womack JE. Sequence variability and protein domain architectures for bovine Toll-like receptors 1, 5, and 10. *Genomics*, 2007; 90(4): 502-515.
- [44] Tapping RI, Omueti KO, Johnson CM. Genetic polymorphisms within the human Toll-like receptor 2 subfamily. *Bioch. Society Transact.* 2007; 35(6): 1445-1448.
- [45] Cuscó A, Sánchez A, Altet L, Ferrer L, Francino O. Non-synonymous genetic variation in exonic regions of canine Toll-like receptors. *Canine Genet. Epidemiol.* 2014; 1(1): 11.
- [46] Matsushima N, Tanaka T, Enkhbayar P, Mikami T, Taga M, Yamada K, Kuroki Y. Comparative sequence analysis of leucine-rich repeats (LRRs) within vertebrate toll-like receptors. *BMC genomics.* 2007; 8(1): 124.
- [47] Ciampolini R, Ciani E, Roux M, Cecchi F, Mazanti E, Tancredi M, Castellana E, Presciuttini S, Amarger V. Haplotype reconstruction from unphased genotype data at the bovine PRKAG3 gene. *Annali della Facoltà di Med. Vet.* 2007; 59: 81-86.
- [48] Osman NM, Abou Mossallam AA, El Seedy FR, Mahfouz ER. Single Nucleotide Polymorphisms in TLR4 Gene and Endometritis Resistance in River Buffalo (*Bubalus bubalis*). *Jordan J. Biol. Sci.* 2018; 11 (5): 577- 583.
- [49] Bochud PY, Chien JW, Marr KA, Leisenring WM, Upton A, Janer M, Rodrigues SD, Li S, Hansen JA, Zhao LP, Aderem A. Toll-like receptor 4 polymorphisms and aspergillosis in stem-cell transplantation. *New England J. Med.* 2008;359(17): 1766-1777.
- [50] Jann OC, King A, CorralesN, Anderson SI, Jensen K, Ait-Ali T, Tang H, Wu C, Cockett NE, Archibald AL, Glass EJ. Comparative genomics of Toll-like receptor signalling in five species. *BMC genomics*, 2009; 10 (1): 1.

Table (1): PCR primers , Annealing temperature and amplicon length of amplified bubaline TLR2

Region of TLR2 gene	Sequence 5`-3`	Annealing Temp. TM _{ann} (C)	Size (bp)
5`UTR(Exon1)	F: GGAAATGCTTAGGTTCAAATCG R:TTACCCTGAAACACTCGGTCT	54 °C	382

CD (Exon2)	F: TGAATTAAGCCATGATGTCAA R:TTCTGTCCAAACTCAGTGCTC	54 °C	980
CD(Exon2)	F:TCAATCCAGAACATTAGCCATCT R:GGAAATATAAAGTTCTTTGAGTTGTGG	53 °C	870
CD(Exon2)	F:GGCAGAATCGTTTGAAATCAC R:AATGGGAGAAGTCCAGCTCA	54 °C	981
CD(Exon2)	F:AGGGACATCTGCTACGATGC R:CGTCGCTAAATTCTAACTCTTGC	55 °C	812

Table (2): Allele and genotype frequencies of TLR2 mutual SNPs

SNPs position	P value of Allelic comparison	P value of Genotypic comparison
5` UTR 90M(A/C)	0.043*	0.005*
5` UTR 124R(G/A)	0.044*	0.003*

CD 14K(T/G)	1	1
CD 53 Y(T/C)	0.36	0.88
CD 108 Y(T/C)	0.36	0.95
CD 153 R(A/G)	0.68	0.99
CD 156 Y(C/T)	0.08	0.43
CD 374 Y(C/T)	0.043*	0.005*
CD 381 R(A/G)	0.08	0.43
CD 482 S(G/C)	0.08	0.43
CD 483 Y(C/T)	0.08	0.43
CD 519 S(G/C)	0.68	0.98
CD 1034 S(G/C)	1	1
CD 1375 Y(C/T)	1	1
CD 1678 R(G/A)	0.68	0.97
CD 1707 Y(T/C)	0.68	0.96
CD 2064 Y(C/T)	0.078	0.19
3' UTR 158 S(G/C)	0.07	0.27
3' UTR 231 Y(C/T)	0.68	0.95

-*Statistically significant values (P value <0.01).

-Fisher Exact test was used because of the small number of samples.

-Bonferroni was used for p value correction.

Table (3): Odds Ratio test results of statistically significant TLR2 mutual SNPs.

SNPs	Genotype	P-value	OR	95% CI
90M(A/C)	AA	0.0015*	2.88	1.58-5.25
	AC	0.771	0.724	0.415-1.265

	CC	0.011*	0.297	0.131-0.674
124R(G/A)	GG	0.0002*	0.320	0.176-0.583
	GA	0.202	1.439	0.823-2.517
	AA	0.002*	3.833	1.634-8.992
374 Y(C/T)	CC	0.0015*	2.88	1.58-5.25
	CT	0.771	0.724	0.415-1.265
	TT	0.011*	0.297	0.131-0.674

-*Statistically significant p values ($p < 0.05$) and ORs.

-CL: confidence level.

Table (4): Non-synonymous SNPs and amino acids substitution in TLR2 CDs of healthy and diseased uterine samples

DNA sequence variants	Amino Acid position	Predicted Amino acid
14 K(T/G)	5	Leucine(L)/Tryptophan (W)







53Y(T/C)	18	Methionine(M)/Threonine(T)
374 Y(C/T)	125	Alanine(A)/Valine(V)
482 S (G/C)	161	Serine(S) /Threonine(T)
483 Y(C/T)	161	Serine(S) /Threonine(T)
1034 R(A/G)	345	Asparagine(N)/Serine(S)
1678 R(G/A)	560	Alanine(A)/Threonine(T)
1747 W (T/A)	583	Serine(S) /Threonine(T)

- Bold SNPs indicate mutual SNPs in healthy and diseased samples.

- Italic SNPs indicate SNPs in healthy samples only.


Table (5A): Detected SNPs and their effect on protein function in TLR2 CDs of diseased samples

CDs Exon	SNPs positions	Amino acid substitution	Protein domain	Amino acid substitution effect
-----------------	-----------------------	--------------------------------	-----------------------	---------------------------------------

Exon(2) CDs	14 K (T/G)	Leu (Non Polar) 5 Trp (Non Polar)	-*	This mutation is predicted to be PROBABLY DAMAGING with a score of 0.999 (sensitivity: 0.14; specificity: 0.99) 
Exon(2) CDs	53Y (T/C)	Thr (Polar) 18 Met (Non Polar)	-*	This mutation is predicted to be POSSIBLY DAMAGING with a score of 0.954 (sensitivity: 0.79; specificity: 0.95) 
Exon(2) CDs	374 Y (C/T)	Ala (Non Polar) 125 Val(Non Polar)	LRR_8 (Vs)	This mutation is predicted to be BENIGN with a score of 0.001 (sensitivity: 0.99; specificity: 0.15) 
Exon(2) CDs	482 S (G/C) 483 Y (C/T)	Ser (Polar) 161 Thr (Polar)	LRR_8 (Vs)	This mutation is predicted to be BENIGN with a score of 0.002 (sensitivity: 0.99; specificity: 0.30) 
Exon(2) CDs	1034 R (A/G)	Asn (Polar) 345 Ser (Polar)	LRR (Vs)	This mutation is predicted to be POSSIBLY DAMAGING with a score of 0.491 (sensitivity: 0.89; specificity: 0.90) 
Exon(2) CDs	1678 R (G/A)	Ala (Non Polar) 560 Thr (Polar)	LRR_CT	This mutation is predicted to be BENIGN with a score of 0.000 (sensitivity: 1.00; specificity: 0.00) 

Table(5B): Polyphen-2 analysis of amino acid substitution effects in TLR2 CD of healthy samples only

CDs Exon	SNPs	Amino acid	Protein	Amino acid substitution
----------	------	------------	---------	-------------------------

	positions	substitution	domain	effect
Exon(2) CDs	1747 W (T/A)	Ser (Polar) 583 Thr(Polar)	LRR_CT	<p>This mutation is predicted to be BENIGN with a score of 0.000 (sensitivity: 1.00; specificity: 0.00)</p>  <p>0.00 0.20 0.40 0.60 0.80 1.00</p>

-SNPs positions were calculated by taking the ATG start codon as position 1 on the GenBank sequence: JN786600

- Amino acids substitutions are considered according to the ATG start codon

-* Protein position without known function

UNDER PEER REVIEW