

BIO-EVALUATION OF THE ECOLOGICAL HEALTH STATUS OF PETROLEUM EFFLUENT RECEIVING WATER OF ELEME RIVER, NIGERIA

ABSTRACT

Ecological Health Status of Eleme River receiving treated petroleum effluent wastes from Eleme Petroleum Refinery Company was evaluated using a combined approach of biological methods; petroleum effluent bioindicators, functional diversity and community assemblage metrics, and physico-chemical methods; TPH, nutrients, turbidity, DO, pH, Heavy metals, and total dissolved solids analysis. This was carried out, knowing that effluent discharges changes the biodiversity of aquatic macroinvertebrates particularly insects, reduce their functional roles, and changes their physico-chemical requirements. The River was divided into 3 stations; upstream (receiving point), midstream and downstream, with each station further divided into 3 sampling points. Collection of macroinvertebrates was done, using handnets and Eckman's grab from floating vegetations, logs and sediments. The samples were placed in a 4-L bucket containing 70% ethanol and 5% glycerin, and transported to the Entomology Research Laboratory, Department of Animal and Environmental Biology, University of Port Harcourt for sorting and identification. Composite water samples were taken for analyses of physico-chemical properties using standard devices and methods. Results obtained indicated that out of 18 species encountered, 12 were monitor indicators, 5 facultative tolerant indicators and none were sensitive indicators. Five species (taxa) occurred at both up-and mid-streams and seven occurred at the downstream. *Daphnia barbata* and *macrobranchium sp.*, occurred commonly in the three stations. The insect-indicators encountered were seven, all of the downstream, and was dominated by insects belonging to the order: Hemiptera. Three functional groups; Detritivorous, collector-herbivorous and predators were encountered with detritivorous being dominant. The TPH, nitrate, phosphate, copper and nickel concentrations in all sampling stations were above the intervention limits of World Health Organization (WHO). Concentrations of physico-chemical properties except DO and pH were higher in downstream. Shannon-weiner index showed that absence of species belonging to Ephemeroptera, plecoptera and Trichoptera in all stations indicated that Eleme River have a poor ecological health status.

Keywords: Ecological Health, bioindicators, functional diversity, petroleum effluents, physico-chemicals.

INTRODUCTION

Eleme River is a fresh water body and a branch of Imo River; situated at Aleto and Agbonchia communities, in Rivers State, Nigeria. Eleme is an oil producing area, hosting two of Nigeria's four petroleum refineries whose treated effluent wastes are usually discharged into the river. Petrochemical wastes discharged into rivers is highly toxic (Abduli et al., 2006) and usually modify the physical and chemical components of the receiving water bodies.

Some of the waste effluents are persistent in the water bodies and lead to instability, disorder, harm or discomfort to both physical systems and living organisms inhabiting the ecosystem (Dawodu et al., 2015). Kuehn et al (1995), pointed out a positive correlational relationship between fish health and the aromatic hydrocarbons from petrochemical waste effluents, while Sangodoyin (1991) stated that such waste effluents affect the biodiversity of aquatic flora and fauna.

The impact of petrochemical waste effluents on the physicochemical and heavy metals of receiving water bodies has been described. Kanu and Achi (2011) reported an increase in physicochemical properties of Eleme river while Gbarakoro et al. (2020) stated that though variations in such parameters occur but they were within World Health Organization (WHO), (1990) and Nigeria Federal Ministry of Environment (1991) permissible limits. Parameters such as Electrical conductivity, turbidity, TSS, TDS, calcium hardness, COD, BOD were pointed out as above permissible limits with pH values showing alkaline upstream and slightly acidic downstream (Gbarakoro, et al., 2020). Gheorghe, et al., (2017) reported that petrochemical effluents wastes increased heavy metal levels in water bodies which caused its bioaccumulation in aquatic organisms.

The discharge of petrochemical effluent wastes and its attendant changes on the stability of the water body constitute disturbance of the ecosystem. Pickett, and White (1985) defined disturbance as any relatively discrete event in time that disrupts ecosystem, community or population structure and changes resources, substrata availability or the physical environment.

The discrete event of petrochemical effluent waste discharge into Eleme river can be assessed using indicators principally physico-chemical and biological indicators in rivers (Kambwiri et al., 2014). The use of biological indicators for evaluation of the ecological health of rivers and their landscapes is the best method because it provides complete spatial-temporal integration of knowledge of fresh water ecosystems and their landscapes (Uherek et al., 2014). Abel (2002) stated that the method can reveal the occurrence of ecologically-significant environmental changes and provide needed data for further investigation. Tagliapietra and Sigovini, (2010), stated that biological indicators particularly benthic communities are suitable for use because they provide information on environmental conditions either due to two factors; i. sensitivity of single species to environmental changes (indicator species)

ii. possession of features that enable them to integrate environmental signals over a long period of time. They listed out the features as exposure to chemical contaminants that accumulate in the sediment; and to low dissolved oxygen levels that often occur near the bottom surface; and limited mobility that restricts their ability to avoid adverse conditions. Physicochemical approach is vested with limitations such as restriction of data to the moment of sampling without taking cognizance of the patchy distributions of chemicals in the environment at long term (Resende et al., 2010). It is time consuming and expensive for most developing countries (Deborde et al., 2016), and often limited to suspect compounds, priority pollutants, or major component of a mixture.

Biological indicators such as benthic macro invertebrates which have high efficacy for bio indication of ecosystem health have been used for evaluation (Zerniawska-Kusza, 2005). Okiwelu (1996) stated that insects are effective bioindicators for monitoring landscape changes because of their abundance, species richness, ubiquitous occurrence and importance in the function of natural ecosystems. Uwaigbae et al (2020) observed that insects provide information on the habitat and aquatic ecosystem integrity due to their sensitivity to pollution. The determination of the ecological health status of fresh water bodies in the tropics, particularly Nigeria and Eleme to be specific has been restricted to physicochemical approach. Available literature on the use of biological approach is limited, therefore, the present study, used both approaches with an objective to ascertain the ecological health status of Eleme river. The use of combined approach to obtain information on the causes of degradation and the consequent effect on aquatic ecosystem has been suggested (Carmen, 1988; James and Daniel, 1981). We applied biological evaluation methods as contained in Elios and Chilima, 2017, and functional diversity index.

MATERIALS & METHODS

DESCRIPTION OF STUDY AREA

The area was divided into three sampling stations; upstream station located on $04^{\circ} 48' 46.2''$ N and $07^{\circ} 06' 45.3''$ E. It is the receiving point of the treated petrochemical effluent waste discharge from petrochemical company. This station is also stressed by mechanical in-stream sand mining, and there is virtually no vegetation cover. The midstream station is located at 1.5km from the upstream on $04^{\circ} 48' 29.8''$ N and $07^{\circ} 06' 38.6''$ E. Vegetation cover is absent, and manual in-stream sand mining operations is occurring in this station. Both stations are contained in the Aleto Eleme section of the river, and have deeper depths.

Station 3 is the downstream located at Agbonchia section of the river where there is vegetation cover, and no in-stream sand mining operations. The depth is shallow and the station which lies on a coordinate of $04^{\circ} 05' 20.2''$ N and $06^{\circ} 53' 54.4''$ E is about 2km from station 2.

Each station was divided into three sampling points which serve as replicates in a completely Randomized Design.

COLLECTION OF BIOLOGICAL MATERIALS

Biological materials used for analysis of ecological health status were collected from the three sampling stations, using Eckman's grab and hand net devices. The devices were used to sample randomly from the sediments, logs, and vegetation cover. The Eckman's grab which measure 225cm^3 was used in a wooden canoe to collect biological materials from about 15 to 28cm depth, and sieved through a 0.5mm mesh size net and immersed into the water to wash off unwanted particles. The biological materials were collected from upstream and mid-stream and preserved in a 4-litre bucket with 70% ethanol and 5% glycerin.

Floating vegetation and logs at the river banks and channels were searched and dislodged, and hand nets placed underneath for collection of biological materials. Unwanted materials

collected along were sieved and removed, and the biological materials placed in a 4-L bucket containing 70% ethanol and 5% glycerin. All the biological materials were stained with 0.1% Eosin red stain and transported to the Entomology Research Laboratory, Department of Animal and Environmental Biology, University of Port Harcourt.

SORTING AND IDENTIFICATION

The sorting technique used enhanced easy sorting; which consists of picking up from the sieve material all the animals that were alive at the moment of the sampling (Tagliapietra & Sigovini, 2010). Sorting and identification were carried out under a binocular dissecting microscope.

Identification was conducted using standard keys, illustrations and typed specimens (Dudgeon, 1999), in which small samples were spread onto petri dishes and carefully examined to identify the species. Biological materials (macro invertebrates) were picked and placed in different containers according to main taxonomic groups; oligochaetes, crustaceans and insects. The containers were labeled according to the sampling stations with identified species counted and grouped according to their functions, indicator ability, pollution tolerance, species richness and abundance. This was used to determine pollution indicators, community assemblage metrics and functional diversity.

DETERMINATION OF PETROLEUM EFFLUENT INDICATORS

Three categories of effluent indicators were determined by counting the taxa or species that were encountered in the three respective sampling stations of the river sections. Species counted for each section were recorded as tolerant to its habitat specific concentration of the stressor. Those species which were somewhat tolerant as they occur at one sampling station but absent at another station, were counted and called Facultative indicators. Those taxa that usually occur at fresh water bodies but are absent in the studied sites are observed to be sensitive to the stressor and called sensitive indicators.

DETERMINATION OF COMMUNITY ASSEMBLAGE METRICS

Total number of individuals in a taxa (abundance) that belong to three orders of insects; Ephemeroptera, plecoptera and Trichoptera and those that belong to the family Chironomidae were counted and analysed using EPT Metrics. Individual macroinvertebrates belonging to three subcommunities; oligochaetes, Insects and crustaceans were counted and recorded according to their respective habitat-types; upstream, midstream and downstreams to determine their abundances. The abundances obtained were subjected to assemblage or biodiversity analyses using metrics such as Simpson's, Shannon-Weiner and Margalef richness indices to determine the significant differences between the habitat-types on one land, and between subcommunities on the other land.

FUNCTIONAL DIVERSITY INDEX

Organisms that perform the same or similar functions in the water body encountered were assembled together and used to assess the level of functionality performance of the water body.

PHYSICO- CHEMICAL EVALUATION OF ECOLOGICAL HEALTH STATUS OF ELEME RIVER, NIGERIA.

Physico-chemical materials used for evaluation of ecological health status of Eleme river were collected from the sampling stations, using standard devices and methods. Composite water samples from stations were collected and analysed. For pH, temperature and conductivity, calibrated handheld pH electronic meter (D1-4337), digital thermometer and electronic conductivity meter model (H₁-4103) respectively were used.

Turbidity was determined by Nephelometric turbidity meter (NTU), while Total Dissolved Solids (TDS) and total suspended solids (TSS) were determined by Calibrated handheld Electronic TDS meter (D4-7103) and gravimetric method, respectively.

Titrimetric method with AgNO₃ was used to determine chloride concentration, while calibrated handheld electronic TDS/Salinity meter (D4-7103) for salinity, ASTM D3875 for alkalinity. Dissolved oxygen (DO) was determined by calibrated handheld HACH Electronic DO meter, Total Petroleum Hydrocarbon (TPH) by Flame Ionization Detector (FID) - Gas Chromatograph with Agilent 7890N. Heavy metals (Lead, copper and nickel) by calibrated Atomic Absorption spectrophotometer (AAS) and non-metals (Nitrates & Phosphates) by calibrated HACH 3900DR spectrometer.

The values obtained from the analyses were subjected for Hierarchical Cluster Analysis by Jaccard Similarity Index to determine the similarity between the three habitat-types, and Principle Component Analysis (PCA) to determine the relationship between the parameters and insect distribution.

RESULTS

Petroleum Effluent Sediment Indicators

Three key performance indicators of taxa were encountered in the study; indicators or taxa that were tolerant of the effluents; indicators that were somewhat tolerant and tolerated certain level of effluents which are referred to as facultative tolerant indicators; and indicators that were sensitive to the effluents and were not encountered but usually occurred in unpolluted ecosystems.

Petroleum Effluent Tolerant Indicators

Five taxa of macro invertebrates; *Paranais sp.*, *Nais sp.* (oligochaetes), *Daphnia barbata*; *Macrobrachium sp.*, and *Sudanonautus africanus africanus* (crustaceans) were encountered. They tolerated the effluent wastes received by the Aleto section of the Eleme River at the sediments (Table I).

Seven insect taxa; *Lethocerus indicus*, *Naucoris sp.*, *Leptonea sp.*, *Aphelocheirus grik*, *Dytiscus sp.*, *Chironomus sp.*, and *Dragonfly* nymphs, and three crustacean taxa; *Daphnia sp.*, *Macrobrachium sp.*, and *Apus sp.*, tolerated the effluents at the vegetation cover and water column and were encountered during the study (Table 1). Out of the twelve (12) petroleum effluent indicators encountered in the study, five taxa occurred at both the up-and mid-streams, and seven occurred at the downstream (Table 2). Two taxa; *Daphnia barbata* and *Macrobrachium* (Crustacean group) occurred commonly in the samples collected from the three stations.

Seven species of insect indicators encountered were collected from the downstream, and they were dominated by insects that belong to Hemiptera (bugs); *Leptonea sp*; *A.grik*, *L.indicus*, *Dytiscus sp*; and *Naucoris sp*. The other two taxa (species) collected were *Chironomus sp* (Diptera: Chironomidae), and *Dragonfly* nymphs (Odonata: Lebillulidae). (Table 2). Insect-Indicators were all collected from the vegetation cover following a loosening, and oligochaetes were collected from the sediment samples.

Facultative Effluent Tolerant Indicators

In the sediment samples, five taxa or species; *Diplonychus rusticus*, *Ophidonais sp.*, *Chaetogaster sp.*, *Aeolosoma hemprichi* and *Elseniella tetrahedral* were encountered. Four taxa apart from the insecta; *D.rusticus* were absent in the sediment samples collected from the upstream but occurred at the midstream, indicating their somewhat tolerant characteristics (Table 3).

Pollution Sensitive Indicators

None of the species belonging to the EPT orders were encountered at either the Aletto receiving point of the effluents or the Agbonchia downstream end of the river (Table 1). Out of 18 species encountered, 12 tolerated the petroleum effluent waste, and only 5 species were somewhat tolerant. Among the insects, hemipterans (bugs) dominated with 5 species, while one species each was recorded in Coleoptera, Diptera and Odonata.

Functional Diversity

Detritivorous trophic group dominated the sediment samples among the oligochaetes and crustaceans encountered while herbivorous trophic group, particularly collectors dominated among insects. The predator trophic group represented by *Chaetogaster* (Oligochaeta) and *L.indicus* (Insecta) were also encountered.

Community Assemblage Metrics

The result of statistical analyses of Simpson index, Shannon-weiner diversity index, and Mergalef richness index showed that diversity was more the unmined and low at the Mechanical habitat-types (Table 5).

Table 1: Physicochemical Parameter Values of Petroleum Effluent Receiving Eleme River.

Values of Range, Mean, Standard Error of Physicochemical Parameters of Eleme River

Parameters				WHO
				Standard Limit 2016 Edition
	Upstream	Midstream	Downstream	
Temperature (⁰ c)	30	28.0-28.4	27.5-28.0	27.5-29.1
		28.13±0.13	27.77±0.15	28.13±0.49
Salinity (mg/L)	600	20.0-22.1	21.7-23.4	25.0-28.1
		21.1±0.61	22.47±0.50	26.7±0.91
pH	6.5-8.5	6.63-6.68	6.71-6.79	6.34-6.40
		6.65±0.01	6.75±0.02	6.37±0.02
Dissolved oxygen (mg/L)	7.5	5.4-5.8	5.0-6.1	4.02-4.51
		5.63±0.12	5.63±0.33	4.35±0.17
Total suspended solids (TSS) mg/L	30	8.8-20.3	17.9-18.6	20.8-22.4
		19.53±0.43	18.20±0.21	21.57±0.46
Turbidity (NTU)	15	13.8-15.5	11.5-15.8	14.3-16.7
		14.7±0.49	14.1±1.32	15.7±0.72
Nitrate (NO ³⁻) mg/L	10	10.5-12.4	11.3-14.8	15.6-17.5
		11.27±0.58	12.67±1.08	6.33±0.59
Phosphate (PO ₄ ³⁻) mg/L	0.1	0.259-0.271	0.212-0.288	0.37-0.383
		0.266±0.004	0.260±0.024	0.377±0.004
Copper (Cu) mg/L	1.5	0.25-0.31	0.24-0.29	0.31-0.35
		0.29±0.02	0.27±0.01	0.33±0.01
Lead (Pb) mg/L	0.05	0.021-0.025	0.023-0.031	0.029-0.034
		0.023±0.001	0.027±0.004	0.031±0.002
Nickel (Ni) mg/L	0.03	0.114-0.19	0.115-0.122	0.131-0.142
		0.116±0.001	0.119±0.002	0.137±0.003
Total Petroleum Hydrocarbon (TPH) mg/L	0.01	5.78-7.37	4.6-7.08	9.47-13.51
		6.66±0.47	5.84±0.72	11.34±1.18
Total Dissolved Solids (TDS) mg/L	2000	44.30-46.00	47.00-48.00	48.3-52.0
		45.37±0.54	47.40±0.80	49.60±1.02

Table 2: Categories of Petroleum Effluent Indicators Taxa (species) Encountered at Eleme Branch of Imo River During the Period of Study.

Taxa (species)	Group	Tolerant Indicators (TI)	Facultative Moderately Tolerant Indicators (MTI)	Sensitive Indicators (SI)
<i>Nais sp.</i>		E		
<i>Paranais sp.</i>	olig	E		
<i>Ophidonais</i>	och		E	
<i>Chaetogaster sp.</i>	aet		E	
<i>Aeolosoma hemprichi</i>	es		E	
<i>Elseniella tetrahedra</i>			E	
<i>Apus sp.</i>	cr		E	
<i>Daphnia barbata</i>	us	E		
<i>Macrobrachium sp.</i>	ta	E		
<i>Sudanonautus</i>	ce			
<i>Africanus africanus</i>	a	E		
<i>Lethocerus indicus</i>		E		
<i>Naucoris sp.</i>	in	E		
<i>Leptonea sp.</i>	se	E		
<i>Aphelocheirus grik</i>	ct	E		
<i>Dytiscus sp.</i>	s	E		
<i>Chironomus sp.</i>		E		
<i>Dragonfly nymphs</i>		E		
EPT species				
NE				

Legend: TI = Tolerant Indicators
 MTI = Moderately Indicators (Facultative)
 SI = Sensitive Indicators
 E = Encountered
 NE = Not Encountered

Table 3: Petroleum Effluent Indicators Encountered at the three study stations of Eleme River During the Period of Study.

Taxa	Upstream	Midstream	Downstream
<i>Nais sp</i>	E	E	NE
<i>Paranais sp.</i>	E	E	NE
<i>Daphnia barbata</i>	E	E	E
<i>Macrobrachium sp.</i>	E	E	E
<i>Sudanonautus africanus africanus</i>	E	E	NE

<i>Leptonea</i> sp.	NE	NE	E
<i>A.grik</i>	NE	NE	E
<i>L.indicus</i>	NE	NE	E
<i>Dytiscus</i> sp.	NE	NE	E
<i>Chironomus</i> sp.	NE	NE	E
<i>Dragonfly nymphs</i>	NE	NE	E
<i>Naucoris</i> sp.	NE	NE	E

Table 4: Petroleum Effluent Facultative Indicators Encountered at the Stations During the Period of Study.

Vegetation Taxa	Sediments		
	Upstream	Midstream	Downstream
<i>Diplonychus rusticus</i>	E	NE	NE
<i>Ophidonais</i> sp.	NE	E	NE
<i>Chaetogaster</i> sp.	NE	E	NE
<i>Aeolosoma hemprichi</i>	NE	E	NE
<i>E.tetrahedra</i>	NE	NE	E
<i>Apus</i> sp.	NE	NE	E

Table 5: Impact of Sand mining on Species Richness of Macroinvertebrates at Eleme River, Rivers State, Nigeria

Phylum or class	Diversity group	Species (Taxa)	Order	Family	Common name	Up-stream	Mid-Stream	Down Stream
Annelida	OLIGOCHAET (MEIOBENTHIC)	1. <i>Ophidonais</i> sp.	Naididae			-	+	-
		2. <i>Nais</i> sp	Naididae			+	+	-
		3. <i>Paranais</i>	Naididae			+	+	-
		4. <i>Chaetogaster</i>	Naididae			-	+	-
		5. <i>Aeolosoma</i> sp	Naididae			-	+	-
		6. <i>Eiseniella tetrahedra</i>		Lumbricidae	Square tailed worm	-	+	-
Sub-Total						2	6	8
Arthropoda	INSECTA (AQUATIC ENTOMOFAUNAL)	1. <i>Diplonychus rusticus</i>	Hemiptera	Belostomatidae		+	-	-
		2. <i>Lethocerus indicus or americanus</i>	Hemiptera	Belostomatidae	Giant water bug	-	-	+
		3. <i>Naucoris</i> sp	Hemiptera	Naucoridae	creeping water bug	-	-	+

		4. <i>Leptonea sp</i>	Hemiptera	Hydropsychidae		-	-	+
		5. <i>Aphelocheirus grick</i>	Hemiptera			-	-	+
		6. <i>Dystiscus sp.</i>	Coleoptera	Dytiscidae	Predacious diving beetle	-	-	+
		7. <i>Chironomus sp.</i>	Diptera	Chironomidae	Midges	-	-	+
		8. <i>Dragonfly nymphs</i>	Odonata	Libellulidae		-	-	+
Sub-Total						1	-	7
Crustacea	CRUSTACEA	1. <i>Daphnia barbata</i>		Brachiopoda	Water flea	+	+	+
		2. <i>Apus sp.</i>		Brachiopoda	Triops or Tadpole	-	-	+
		3. <i>Macrobrachim sp</i>		Malacostraca	Shrimp Crayfish	+	+	+
		4. <i>Sudanonautus africanus</i>		Malacostraca	Freshwater crab	+	+	+
Sub-Total						3	3	4
Grand Total						6	9	11

Table 6: Diversity indices of the Habitat-types

Diversity indices	Habitat-types		
	Mechanical	Manual	Unmined
Taxa_S	6	9	11
Individuals	11	121	46
Dominance_O	0.273	0.166	0.124
Simpson_1-D	0.727	0.835	0.876
Shannon-H	1.509	1.904	2.214
Evenness_e^H/S	0.754	0.746	0.832
Brillouin	1.160	1.782	1.903
Menhinick	1.500	0.818	1.622
Margalef	1.803	1.668	2.612
Equitability_J	0.842	0.866	0.923
Fisher_alpha	3.487	2.248	4.580
Berger-Parker	0.438	0.223	0.196
Chao-1	9	9	11.5

Table 7: Percentage abundance (%) of invertebrate groups collected from the study sites

Phylum	1	% abundance	2	% abundance	3	% abundance	Total	% abundance
Annelida	10	12.5	115	38.08	0	0	125	26.60
Arthropoda	69	86.25	186	61.59	88	100	343	72.98
Mollusca	1	1.25	1	0.33	0	0	2	0.43
Total	80	100	302	100	88	100	470	100

Table 8: Percentage abundance of arthropod classes Encountered during the study

Phylum	1	% abundance	2	% abundance	3	% abundance	Total	% abundance
Annelida	4	5.80	0	0	42	47.73	46	13.41
Arthropoda	65	94.20	186	100	41	46.59	292	85.13
Mollusca	0	0	0	0	5	5.68	5	1.46
Total	69	100	186	100	88	100	343	100

The TPH, Nitrate, Phosphate, copper, Nickel concentrations in all the sampling stations were above the intervention limits of World Health Organization (WHO) (Table 4). The salinity and nitrate of the up-stream that received the effluent waste discharge was less in concentration with mean values of 21.1 0.61mg/L, and 11.27±0.58mg/L, respectively. The station also contained the least concentration of Lead (0.023±0.001mg/L) and Nickel (0.116±0.00mg/L) (Table 4). The least concentration of TPH was recorded at mid-stream with a mean of 5.84±0.72mg/L.

The highest concentrations of physicochemical properties were recorded at the downstream: salinity (26.7±0.91mg/L), turbidity (15.7±0.72ntu), Nitrate (16.33±0.59mg/L), Phosphate (0.377±0.004mg/L), Copper (0.33±0.01mg/L), Lead (0.031±0.002mg/L), Nickel (0.137±0.003mg/L) and TPH (11.34±1.18mg/L) (Table 3). Downstream recorded the least concentrations of Dissolved Oxygen (4.35±0.17mg/L) and pH (6.37±0.02). Both suspended and dissolved solids were high in the station with mean values; 21.57±0.46mg/L (TSS) and 49.60±1.02mg/L (TDS).

DISCUSSION

The absence of EPT species in all stations of the Eleme river investigated is an indication of Ecological poor health status. This is because, species of EPT orders occur in good healthy water bodies and are usually sensitive to pollution and cannot inhabit polluted water ecosystem. This collaborate reports of Barman (2014) and Elias and Chilima (2017) that organisms belonging to the Ephemeroptera (Mayflies), Plecoptera (stoneflies), and Trichoptera (caddisflies) (EPT) are sensitive to pollution and typically occur in good quality water.

The number of taxa (species richness) and its individuals (species abundance) occurring in an ecosystem indicates the ecological health status of the ecosystem. Ecosystem richer in species and have high abundance is adjudged to be healthier, while otherwise is unhealthy and data

(Burman, 2014) collected on the reveal the effects of pollutants or changes in the ecosystem and are used to assess anthropogenic impacts on the ecosystem. In this present study, most of the species encountered are tolerant to the petroleum effluent waste discharged into the water body. The collection of a high level of tolerant taxa is an indication that the Eleme river is ecologically poor in health. Infact, out of 18species encountered, 12 (66.67%) are in the tolerant category.

The high number of taxa of oligochaetes and insect particularly Hemipterans (bugs) that dominated the sediments and vegetation, respectively, showed that these group of species have high tolerance for petroleum effluent wastes. The high number of oligochaetes collected from the mid-stream over the upstream implies that though both streams are poor in ecological health, that of the upstream is poorer (Table 2). Moreso, the occurrence of *E.tetrahedra* which usually occur in good freshwater in the mid-stream showed that the mid-stream is a little bit healthier. This agrees with the report that some genera of Lumbriculidae associate with water body that have good water quality (Fend, and Lenat, 2007).

Cheimonopoulou, et al. (2010) stated that the main river reaches of Greek river possess moderate water quality, while downstream mainly contained poor water quality, due to its dominance by pollution tolerant macro invertebrate taxa. This was also observed in our study, as the absence of insect species at both up-and mid-streams, and its dominance by pollution tolerant species at downstream is an indication of poor-quality water. The non-availability of vegetation habitat at both up-and mid-streams is responsible for the absence of petroleum effluent waste-tolerant insect species. Payakka and Prommi (2014) argued that the presence of riparian vegetation and suitable substrate increase insect taxa and abundance in aquatic habitats because vegetation may provide protection from predators, and food for the insects, sites for oviposition and source of respiratory oxygen. This situation may have prevailed at a very reduced rate at the downstream and completely absent at both up-and mid-streams of our study sites.

The high level of TPH recorded at the downstream is an indication that the downstream is more impacted by the petroleum effluent waste. The reduction in the TPH level at up-stream (point of effluent entry) is due partly to increase in the volume of water and water flow (current) caused by sand mining. This led to flow of the pollutant to the downstream where there is less mixing of water. The presence of insect taxa that belong to three orders (Hemiptera, Odonata and Diptera) of tolerant species at the downstream, indicates that the downstream is more impacted, and contain ecologically unhealthy water (poor quality water). These species; Dragonfly (Odonata), Bugs (Hemiptera), and *Chironomus* sp; (Diptera) has been described as pollution tolerant species that occur in polluted water (Barman, 2014). Our study agrees with the description.

The natural stream water contain dissolved oxygen range of 5-13mg/L, and less than 1.0mg/L for nitrates (Barman, and Gupta, 2015). However, in this present study, DO ranges is 4.35-5.63mg/L. The low DO, and pH, high nitrates and phosphates recorded in our study; particularly in the downstream is an indication of poor water quality. This agrees with the report that such situation can cause a reduction in water quality (Payakka, and Prommi,

2014), especially as it does not favour assemblage of Trichoptera which is a pollution sensitive group whose assemblage is favoured by high values of DO and low levels of nitrate (Barman, and Gupta, 2015).

The high level of TPH triggered by the petroleum effluent wastes caused a variation in the levels of some physicochemical properties which consequently cause a poor ecological health status of the Eleme water body of Aletto and Agbonchia.

Living organisms contribute immensely through their functions to their communities and ecosystems. In our study three groups of functional diversities were encountered; predatory, detritivorous and collectors. The abundance of detritivores; oligochaetes and crustaceans, including *Daphnia* sp indicates that decomposition or degradation is on-going in the ecosystem at the sediment levels in the midstream. The reduced number of detritivores at the upstream is an indication that decomposition is slow and inefficient when compared with the midstream.

In the downstream, collector-functional group is abundant showing that filterer-detritivorous activities are also on-going.

The inducement of the dominance of decomposers comprising detritivorous and collectors functional groups over predatory functional group by petroleum effluent wastes is in good faith for the sediments and water at midstream and downstream, respectively. It will naturally, though at a longer time degrade the pollutant. This responsibility is lacking in the upstream, but at the midstream it is performed by Oligochaetes and Crustaceans, while the insects carry it out in the downstream.

This agrees with the report that the abundance of a functional group in an ecosystem indicates different quality of the water body (Barman, 2014) and provides information on the state of ecosystem functionality (Paraschiv *et al.*, 2013). Ecosystem dominated by predator-insects are usually not good while those dominated by collector-insect functional group are better ecosystem (Barman, 2014) as it will help in natural recovery.

CONCLUSION

The absence of insect species belonging to the orders; Ephemeroptera, plecopteran and Trichoptera indicate that the Eleme river is polluted, however, the dominance of collector-functional groups is an indication that the ecosystem is undergoing natural recovery. The study also show that Eleme river contain petroleum effluent tolerant indicators.

REFERENCES

- Abduli, M.A., Samieifard R, and Jalili Ghazi Zade (2008). Rural Solid Waste Management. *Int. J. Environ Res.* 2(4): 425-430.
- Abel, P.D. (2002). *Water Pollution Biology* 2nd Edition. Taylor and Francis. Sunderland, UK. 287p.
- Barman, B. (2014). The importance of Aquatic Insects as Biomonitors of Freshwater Ecosystem. *International Journal of Environmental and Natural Sciences*, Vol.1, 82-85.

- Barman, B., and Gupta, S. (2015). Aquatic Insects as Bioindicator of water quality. A study on Bakuamari stream, Chakras hila Wildlife Sanctuary, Assam, North East India. *Journal of Entomology and Zoology Studies*, 3(3):178-186.
- Cheimonopoulou, M.T., Bobori, D.C., Theocharopoulos, I., Lazaridou, M. (2011). Assessing Ecological water quality with Macro invertebrates and Fish: a case study from a small mediterranean River. *Environmental Management* 47:279-290.
- Czerniawska-Kusza, I. (2005). Comparing modified biological monitoring working party score system and several biological indices based on macroinvertebrates for water quality assessment. *Limnologica*. Vol.35(3):169-176.
- Dawodu, B., Zaid A., Olelekan, E.I. Falilie, A., (2015). Consequences of Anthropogenic Activities on Fish and the Aquatic Environment. *Poultry, Fisheries and Wildlife Sciences* 3:138.
- Deborde, D.D., Hernandez, M.B.M. and Magbanua, F.S. (2016). Benthic Macroinvertebrate Community as an Indicator of Stream Health: The Effects of Land use on Stream Benthic Macroinvertebrates: *Science Dilima*.28(2), 5-26.
- Elias, R.C. and Chilima, L. (2017). Use of Benthic Macroinvertebrate Indices as Bioindicators of Ecosystem Health for the Detection of Degraded Landscapes in Malawi. *International Journal of Agriculture, Forestry and Fisheries*. 5(6): 128-134.
- Fend, S.V, and Lenat, A.R. (2007). Two new genera of Lumbriculidae (Annelida, Clitellata) from North Carolina, USA *Zootaxa* 1666:1-22.
- Gbarakoro, S.L., Gbarakoro, T.N., and Eebu, W.L. (2020). Impact of Industrial Effluent Discharge on the Physicochemical Properties of Aletto stream, Eleme, Rivers State, Nigeria. *Annual Research and Review in Biology*, 35(1):79-89.
- Georghe S, Stoica, C, Vasite, G.G., Nita Lazar, M. Stanescu, E. and Lucaciu, I.E. (2017). Metal Toxic Effects in Aquatic Ecosystems; Modulators of Water Quality (eds. Hlanganani Tutu), 4: 65-89. Available: <https://www.intechopen.com/chapters/5263>.
- Georghe, S., Stoica, C., Vasite, G.G., Nita Lazar, M., Stanescu, E., and Lucaciu I.E. (2017). Meal Toxic Effects in Aquatic Ecosystems: Modulators of Water Quality (ed: by Hlanga-nani Tutu), 4:65-89. Available: <https://www.intechopen.com/chapters/52639>.
- Kambwiri, A.M., Changadeya, W., Chimphamba, J, and Tandwe, T.(2014). Land use impacts on water quality of Rivers draining from Mulanje Mountain: A case of Ruo River in the Southern Malawi. *Malawi Journal of Science and Technology*, Vol.10(1), 2014
- Kanu, I., and Achi,O.K. (2011). Industrial Effluents and their impacts on water quality of receiving Rivers in Nigeria - Incomplete
- Kuehn, R.h., Berlin, K.D., Hawkins, W.E., and Ostrander, G.K.J. (1995). Toxicology and Environmental Health, 46:101-106 - Incomplete
- Okiwelu, S.N. (1996). Entomology, Industry and Environment-Reflections. A symposium paper presented at the 27th Annual Conference of the Entomological Society of Nigeria, at the Rivers State University of Science and Technology, Port Harcourt, 25-28 November, 1996.
- Paraschiv, G, Bucur, M., Ehlinger, T., Negreanu-Pirjol, T., Vasile, D., Hennessy, J, and Tofan, L. (2013). Benthic invertebrates species as Indicators of Benthic Habitats Health from Siutghiol lake, Romania. *Carpathian Journal of Earth and Environmental Sciences*. Vol.8, (4):185-194.

- Payakka, A., and Pormmi, T. (2014). Aquatic Insects Biodiversity and Water Quality Parameters of Receiving Water body. *Current World Environment*. Vol.9(1),53-58.
- Pickett, S.T.A. and White, P.S. (1985). *The Ecology of Natural Disturbances and Patch Dynamics*, Academic Press, Science 472 pages, Orlando.
- Resende, P.C., Resende, P., Pardal, M., Almeida, S. and Azeiteiro, U., (2010). Use of Biological Indicators to Assess Water Quality of the UI River (Portugal). *Environ. Monit. Assess.* 170:535-544.
- Sangodoyin, A.Y. (1991). Groundwater and Surface Water Pollution by Open Refuse Dump in Ibadan, Nigeria. *Journal of Discovery and Innovation*, 3(1):24-31
- Tagliapietra, D., and Sigovini, M (2010). Benthic Fauna: Collection and Identification of Macroinvertebrates. *NEAR Curriculum in Natural Environmental Science, Terre et Environment*, Vol.88:253-261.
- Uherek, C.B., and Gouvela, F.B.P (2014). Biological Monitoring Using Macroinvertebrates of Water Quality of Maroaga stream in the Maroaga cave system, Presidente Figueiredo, Amazon, Brazil. *International Journal of Ecology*, Vol.2014, Article ID 308149, 7pages. Hindawi Publishing Corporation.
- Uwaigbae, M.A.E, Abdul-Rhaman, D., Gbarakoro, T.N., El-Surtasi, E.L., Agwu, J.E., Martins, S.O., Rotimi, J.A. Impact of Disturbances on the Biodiversity of Ijala-Ikeren Wetland Ecosystem in Niger Delta. In: L.Hufnal (ed.) *Changing Ecosystems and their services* (1st Edition)pp1-19.