

Temperature variation in Kerala:Present and future

ABSTRACT

Temperature is changing worldwide and a warming trend is expected in most parts. This study investigates the long-term (1983-2020) trend analysis of the maximum temperature in Kerala and the future trend in maximum temperature. The future temperature has been downloaded from GFDL-CM3, which is most suited for Kerala conditions was used to assess the future climate change under different representative concentration pathways (RCP) 4.5 and 8.5 scenarios. The Mann-Kendall test was also used for trends analysis for the downloaded data for each RCPs. The baseline (1980 – 2020) temperature was compared with future (2021-2050 and 2051-2080) simulations for the 14 districts of Kerala under different RCPs. An increasing trend of maximum temperature was observed in all districts of Kerala during the period of 1980 -2020. The increasing temperature trend was confirmed by Mann- Kendall test at a 0.01 significant level. The warming is expected to increase in future simulations under RCP 4.5 and 8.5. In the high-range zone, a decrease in temperature is expected during the beginning of the mid-century, but it continues to show an increasing trend in the projected climate. Under RCP 4.5 in most districts, an increase in temperature by 0.03 C per year is expected in the future. While under RCP 8.5 the temperature is expected to increase by 0.05 C per year and 0.06 C per year by mid and end century respectively.

Keywords: Temperature, GFDL-CM3, Warming, Mann-Kendall, Representative Concentration Pathways (RCP)

1. INTRODUCTION

Temperature is a vital climate quantity that directly influences human and natural systems. The annual mean temperature over India is showing a warming trend (Kothawale, 2005). According to AR5, IPCC, 2013 the global mean surface temperature is a significant indicator of climate change because it increases quasi-linearly with increasing greenhouse gas emissions. Recent IPCC reported last decade *i.e.* 2011-2020 as the warmest one. The temperature of earth has increased by 1.1 °C from late 1800. The last decade between 2011-2020 was the warmest one on record (Kikstra, 2022). Mondal et al. (2015) conducted a study on the long-term trend in temperature from 1901 to 2007 over India. He observed an increase in the mean temperature of 0.003 °C/year. Nayak et al. (2022) reported a possible warming of 3–5 °C over the entire Indian region in the period 2051–2110 with reference to the period 1951–2010. Raj et al., (2022) studied the long-term (1950-2018) temperature variation over Pattambi Palakkad district in Kerala and observed an increasing trend in maximum temperature. The future climate has been downloaded using Global climate models. Global climate models aid in obtaining information on climate change based on different greenhouse gas emission scenarios. Impacts of climate change on the area of interest are obtained from regional climate models used for downscaling global climate simulations to smaller areas (Chou, 2014). Different paths are taken by the models to reach four different radiative forcings that correspond to different concentration paths of


32 the greenhouse gases, the so-called Representative concentration pathway (RCPs)
33 (Zhai, 2019). The pathways describe different climate futures, all of which are
34 considered possible depending on the volume of greenhouse gases (GHG) emitted in
35 the years to come. Under RCP 4.5 a peak emission of greenhouse gas is expected
36 around 2040 and then it declines; under RCP 8.5 the emission is expected to increase
37 continuously throughout the 21st century (Van Vuuren, 2011). Kaur and Kaur,2022 also
38 observed that the rise temperature was expected to higher under RCP 8.5 than under
39 RCP 4.5. This paper aims to study the present and future variation in the temperature of
40 Kerala

41
42
43
44
45

2. MATERIAL AND METHODS

2.1. Study area

46 Kerala is the southwestern state of India. Located between 8°18' and 12°48' N latitude
47 and 74°52' and 77°22' E longitude. The state has a total geographical area of 38863 sq.
48 km (Ajithkumar, 2015) Kerala has 14 districts and the study has been done by analyzing
49 the weather data in 14 districts of Kerala. The whole area has been divided into five
50 agro-climatic zones i.e. Northern zone, central zone, high range zone, problematic zone,
51 and southern zone (Fig. 1).



52
53
54

Fig.1. Study area: Kerala

55

2.2. Data and methodology

56
57

The maximum temperature from 1983 to 2021 has been collected from India Meteorological Department (IMD) and the future temperature has been estimated using

58 the climate model, GFDL-CM3 developed at the Geophysical Fluid Dynamics
 59 Laboratory. The GFDL Climate Model version 3 (CM3) is a coupled general circulation
 60 model that has been developed for the atmosphere, ocean, land, and sea ice. The main
 61 aim of CM3 was to concentrate on the novel problem of climate change which includes
 62 interaction between aerosol and cloud, chemistry and climate, and coupling between the
 63 stratosphere and troposphere (Donner, 2011). The GFDL-CM3 model showed a good
 64 ability to simulate the present climate over the central zone of Kerala (Venkat et al.,
 65 2017). To assess the future change in the temperature of Kerala the present
 66 temperature (1980 to 2020) is compared with the future scenarios i.e. mid-century (2021
 67 -50) and end of the century (2051-80) under RCP 4.5 and 8.5 and trend analysis has
 68 been also carried out.

70 3. RESULTS AND DISCUSSION

71 The warming was not found to be uniform in Kerala. Long-term trend analysis of
 72 past weather has been compared with the projected temperature under RCP 4.5 and
 73 8.5.

74 *Trend analysis:* Long-term analysis of maximum temperature (1983 to 2020) had been
 75 done. Results of Mann- Kendall test showed a significant increasing trend in Kerala at a
 76 0.01 significant level. The maximum temperature increased at a rate of 0.03°C per year
 77 in most parts of Kerala (Table 1). The study conducted by Arulbalaji, 2020, reported
 78 a significant decrease in vegetation cover during the period of 1988–2019 in Akkulam-
 79 Veli lake basin of Thiruvananthapuram. This enhanced the warming and an average
 80 increase of temperature by 1.6°C was observed. Hence the changes in the land use or
 81 land cover may lead to climate change and increase in temperature.

82 Table. 1. Long term analysis of maximum temperature

District	Mann-Kendall test	Equation
Kasargod	5.594	$y = 0.0284x + 28.404$ $R^2 = 0.6269$
Kannur	5.819	$y = 0.0301x + 28.096$ $R^2 = 0.66$
Kozhikode	5.818	$y = 0.0316x + 27.647$ $R^2 = 0.6523$
Malappuram	5.897	$y = 0.0316x + 27.858$ $R^2 = 0.6705$
Wayanad	6.088	$y = 0.0327x + 27.632$ $R^2 = 0.7036$
Palakkad	5.841	$y = 0.0317x + 28.365$ $R^2 = 0.6588$
Thrissur	5.886	$y = 0.0314x + 28.417$ $R^2 = 0.6488$
Ernakulam	5.762	$y = 0.0308x + 29.103$ $R^2 = 0.6423$
Idukki	5.987	$y = 0.0314x + 29.665$ $R^2 = 0.6644$
Alappuzha	6.133	$y = 0.0308x + 29.768$ $R^2 = 0.6524$
Kottayam	5.975	$y = 0.0308x + 29.768$ $R^2 = 0.6524$
Pathanamthitta	6.009	$y = 0.0312x + 30.293$ $R^2 = 0.6726$
Kollam	6.076	$y = 0.0313x + 30.571$ $R^2 = 0.6759$

83

Thiruvananthapuram	6.065	$y = 0.0317x + 30.967$ $R^2 = 0.6779$
--------------------	-------	--

84

85

3.1. Analysis of future climate-

86

3.3.1. Northern zone of Kerala

87

Trend analysis: The northern zone of Kerala includes districts like Kasargod, Kannur, Kozhikode and Malappuram. From analyzing the projected climate during mid-century and end century the temperature is expected to increase at the rate of 0.03°C per year. Under RCP 8.5 the temperature is expected to increase at the rate of 0.05°C per year during the mid-century and by the end of the century, warming is expected to increase at the rate of 0.06°C per year.

93

Variation from normal: Under RCP 4.5 during the period 2021-2050 (mid-century) an appreciably above normal (3.1 to 5°C) increase in temperature is expected from normal while during 2051-80 (end of the century) warming is expected to increase in Kozhikode and Malappuram by greater than 5°C from normal. Under RCP 8.5 during mid-century an appreciably above normal (3.1 to 5°C) increase in temperature is expected from normal while during 2051-80 (end of the century) markedly above normal increase i.e. greater than 5°C from normal in maximum temperature is expected (Fig 2a-2d).

94

95


96

97

98


99

100




101


2.a)



102 2 b



103 2 c)
104



105 2d)

106 Fig. 2.a-d Temperature variation in the northern zone of Kerala

107

108

3.3.2. The central zone of Kerala

109

110

111

112

113

114

115

116

117

118

119

120

121


122

123


124

Trend analysis: The central zone of Kerala includes districts like Palakkad, Thrissur and Ernakulam. From analyzing the projected climate during mid-century, the increasing trend is expected to continue at the rate of 0.03°C per year. By the end of the century, the increasing rate is expected to decrease. The temperature is expected to increase at the rate of 0.029°C per year, 0.027°C per year and 0.03°C per year in Thrissur, Ernakulam and Palakkad respectively. Under RCP 8.5 the temperature is expected to increase at the rate of 0.05°C per year during the mid-century and by the end of the century, warming is expected to increase at the rate of 0.06°C per year.


Variation from normal: Under both RCPs during the period 2021-2050 (mid-century) an appreciably above normal (3.1 to 5°C) increase in temperature is expected from normal in Thrissur and Palakkad while in Ernakulam an above normal deviation *i.e.* (1.6 to 3.0°C) is expected. By the end of the century markedly above normal increase in maximum temperature is expected in Thrissur and Palakkad while in Ernakulam an appreciably above normal increase in maximum temperature is expected (Fig 3.a- 3 c).



125 3.a)



126 3 b)



127 3 c)

128

129 Fig. 3.a-c) Temperature variation in the central zone of Kerala

130

131

3.3.3. Problem area zone Kerala

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146


147

148


149

Trend analysis: The problem area zone includes Alappuzha and Kottayam districts of Kerala. An increasing trend in maximum temperature was observed from the long-term analysis of maximum temperature (1983 to 2020). The increasing rate per year was found to be 0.03°C per year. Under projected climate also an increasing trend in maximum temperature was observed. Under RCP 4.5 during the mid-century and end of the century, the increasing trend is expected to continue at the rate of 0.03°C per year in Alappuzha. While in Kottayam the increasing rate is expected to decrease and the increase per year is expected to be 0.02°C. Under RCP 8.5 the temperature is expected to increase by 0.04°C per year during the mid-century. By the end of the century, the temperature is again expected to increase by 0.058°C per year.

Variation from normal: Under RCP 4.5 during the period 2021-2050 (mid-century) an above normal (1.6 to 3°C) increase in temperature is expected from normal while during 2051-80 (end of the century) warming is expected to increase in Kottayam by 3.1 to 5°C from normal. Under RCP 8.5 during mid-century an above normal increase in temperature is expected from normal while during 2051-80 (end of the century) appreciably above normal increase from normal in maximum temperature is expected(Fig 4.a- 4 b)..



150 4.a)
151



152 4.b)

153 Fig. 4.a-b Temperature variation in the problem area zone of Kerala

154


155 **3.3.4. High range zone**

156 *Trend analysis:*Two districts, Wayanad and Idukki come under the high range zone.


157 During the past 40 years, maximum temperature showed an increase of 0.03°Cper

158 year in the high range zones of Kerala. Even though a decrease in temperature is

159 expected at the beginning of the mid-century. Under RCP 4.5 the temperature is
 160 expected to increase by 0.03°C during future scenarios in Idukki and Wayanad.
 161 While under RCP 8.5 maximum temperature is expected to increase at the rate of
 162 0.05 °C per year in mid-century and 0.06°C per year by the end of the century.
 163 *Variation from normal:* Under RCP 4.5, the maximum temperature showed below
 164 normal departure in Idukki in mid future and by end of the century, a normal
 165 departure has been expected. The departure is expected to be normal in Wayanad.
 166 In Wayanad, the temperature decreased by 0.1°C in the mid-century while by end of
 167 the century there was a small increase in maximum temperature by 0.9°C. In Idukki,
 168 the maximum temperature decreased by 1.74°C and 0.81°C during the mid and end
 169 of the century respectively. Under RCP 8.5 in Idukki, the temperature decreased by
 170 1.76°C and 0.05°C during mid-century and end of the century respectively. While in
 171 Wayanad temperature decreased in mid-century by 0.11°C and then by the end of
 172 the century an increase in temperature by 1.73°C was found (Fig 5.a- 5 b).



173 5 a)



174 5 b)

175

176 Fig. 5.a-b. Temperature variation in the high range zone of Kerala

177

178

3.3.5. The southern zone of Kerala

179

180

181

182

183

184

185

186

187

188

Trend analysis: The southern zone of Kerala includes districts like Thiruvananthapuram, Kollam and Pathanamthitta. In the southern zone of Kerala from 1980 to 2020, an increasing trend in maximum temperature was observed similar to other zones. In the future under RCP 4.5 maximum temperature is expected to increase at the rate of 0.03°C per year in mid-century while moving towards the end of the century, the increasing rate of maximum temperature decreases to 0.02°C per year. While under RCP 8.5 maximum temperature is expected to increase at the rate of 0.05°C per year in mid-century. By the end of the century the warming is expected to increase at the rate of 0.06°C per year (Fig 6.a-6 b).

189

190

191


192

193


194

195


Variation from normal: Under RCP 4.5 in the southern zone of Kerala temperature was expected to increase in Pathanamthitta and Kollam. In Thiruvananthapuram, a slight decrease in maximum temperature was found during mid-century by 0.01°C and by end of the century, the temperature was found to be increased by 0.8°C. Under RCP 8.5, an increase in temperature was observed In Kollam and Pathanamthitta the temperature increased by 2.3°C and 3.3°C by end of the century respectively.



196 6 a)
197



198 6 b)



199 6 c)

200

201 Fig. 6.a-c Temperature variation in the southern zone of Kerala

202

203

204

4. CONCLUSION

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

The maximum temperature has increased at the rate of 0.03 °C per year in Kerala from 1980 to 2020. The warming is expected to increase in future simulations under RCP 4.5 and 8.5. In the high-range zone, a decrease in temperature is expected during the beginning of the mid-century, but it continues to show an increasing trend in the projected climate. Under RCP 4.5 in most districts, an increase in temperature by 0.03°C per year is expected in the future. While under RCP 8.5 an increase in temperature the temperature is expected to increase by 0.05°C per year and 0.06°C per year by mid and end century respectively. The increasing temperature trend was confirmed by Mann- Kendall at a 0.01 significant level. Increase temperature has a major effect on the agriculture. Increase in temperature had a detrimental effect on crop production even though due to the higher concentration of CO₂ photosynthesis rate has increased leading to higher growth and plant productivity. But higher temperatures counteract this effect by increasing crop respiration rates and evapotranspiration, changing the flora of weeds, increasing pest infestation and shortening the duration of crops (Malhi et al, 2021).

221
222

COMPETING INTERESTS

223 The authors have declared that no competing interests exist.

REFERENCES

224
225
226
227
228
229
230
231

1. IPCC (2013) Climate change 2013: the physical science basis. In: Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM (eds) Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, United Kingdom and New York, NY. p 1535. <https://doi.org/10.1017/CBO9781107415324>

232
233
234
235

2. Kothawale D. R., Rupa Kumar K On the recent changes in surface temperature trends over India, Geophysical. 2005. Res Lett 32. <http://doi.org/10.1029/2005GL023528>

236
237
238
239
240

3. Kikstra, Jarmo S., Zebedee RJ Nicholls, Christopher J. Smith, Jared Lewis, Robin D. Lamboll, Edward Byers, MaritSandstad et al. "The IPCC Sixth Assessment Report WGIII climate assessment of mitigation pathways: from emissions to global temperatures." *Geoscientific Model Development* 15, no. 24 (2022): 9075-9109.

241
242
243
244

4. Mondal A, Khare D, and Kundu S. Spatial and temporal analysis of rainfall and temperature trend of India. *Theor. Appl.Climatol.* 2015. 122(1–2):143–158. <https://doi.org/10.1007/s00704-014-1283-z>

245
246
247
248

5. Raj, PP Nikhil, and P. A. Azeez. "Variations in temperature in a tropical town: a case study." *International Journal of Global Warming* 26, no. 4 (2022): 417-428.

249
250
251
252

6. Chou, S.C., Lyra, A., Mourão, C., Dereczynski, C., Pilotto, I., Gomes, J., Bustamante, J., Tavares, P., Silva, A., Rodrigues, D. and Campos, D., Assessment of climate change over South America under RCP 4.5 and 8.5 downscaling scenarios. *American J. Clim. Change*, 2014.3(05), p.512. 10.4236/ajcc.2014.35043

253
254
255
256

7. Nayak, Sridhara, and Tetsuya Takemi. "Assessing the impact of climate change on temperature and precipitation over India." *Wadi Flash Floods: Challenges and Advanced Approaches for Disaster Risk Reduction; Sumi, T., Kantoush, S., Saber, M., Eds* (2022): 121-142.

257
258
259
260

8. Zhai, Yuanyuan, Gordon Huang, Xiuquan Wang, Xiong Zhou, Chen Lu, and Zoe Li. "Future projections of temperature changes in Ottawa, Canada through stepwise clustered downscaling of multiple GCMs under RCPs." *Climate Dynamics* 52, no. 5 2019: 3455-3470.

- 261
262
263
264
265
266
267
268
9. Van Vuuren, D.P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G.C., Kram, T., Krey, V., Lamarque, J.F. and Masui, T., The representative concentration pathways: an overview. *Climatic change*, 2011. 109(1), pp.5-31.
10. Kaur, Jatinder, and PrabhjyotKaur. "A critical assessment of changes in climate predicted by four GCMs under different RCP scenarios in Punjab (India)." *MAUSAM* 73.4 (2022): 749-762.
- 269
270
271
11. Ajithkumar, B., Rao, V. U. M., Sreekala, P., Chandrasekhar, Vishnu, Sekhar, E. (2015) *Agroclimatic Atlas of Kerala*.
https://www.researchgate.net/publication/341788375_Agroclimatic_Atlas_of_Kerala
- 272
273
274
275
12. Arulbalaji, P., D. Padmalal, and K. Maya. "Impact of urbanization and land surface temperature changes in a coastal town in Kerala, India." *Environmental Earth Sciences* 79, no. 17 (2020): 1-18.
- 276
277
278
279
280
281
13. Donner, L.J., Wyman, B.L., Hemler, R.S., Horowitz, L.W., Ming, Y., Zhao, M., Golaz, J.C., Ginoux, P., Lin, S.J., Schwarzkopf, M.D., and Austin, J. The dynamical core, physical parameterizations, and basic simulation characteristics of the atmospheric component AM3 of the GFDL global coupled model CM3. *J. Climate*, 2011.24(13):3484-3519. <https://doi.org/10.1175/2011JCLI3955.1>
- 282
283
14. Venkat, J. Assessment of rice (*Oryza sativa* L.) production under climate change scenarios.MSc (Ag) thesis. Kerala Agricultural University. 2017. 163p.
- 284
285
286
15. Malhi, Gurdeep Singh, ManpreetKaur, and PrashantKaushik. "Impact of climate change on agriculture and its mitigation strategies: A review." *Sustainability* 13.3 2021: 1318.
- 287