

## Original Research Article

# Influence of irrigation schedules on yield and nutrient uptake of groundnut varieties

### ABSTRACT

**Aim:** Evaluation of different irrigation schedules on yield and nutrient uptake of groundnut varieties

**Study design:** The experiment was laid out in split plot design with different irrigation schedules in main plot and different groundnut varieties in sub plots and was replicated thrice..

**Place and duration of study:** The field experiment was conducted during *rabi* season of 2021 at the Agricultural College Farm, Bapatla, ANGRAU, Lam, Guntur, Andhra Pradesh.

**Methodology:** The experiment was performed with twelve treatments in split plot design. The main plot comprised three different irrigation schedules and sub plot with four different groundnut varieties (TAG-24, Dheeraj, Kadiri Lepakshi and Kadiri Chitravati). Observations of the crop and soil during the experimentation were recorded at regular intervals. The significance of the treatment impact was examined by the test.

**Results:** Among different irrigation schedules, IW/CPE ratio of 1.0 recorded highest pod and haulm yield which was significantly superior over IW/CPE ratio of 0.6 but found on a par with IW/CPE ratio of 0.8. Among the varieties, Kadiri Lepakshi recorded highest pod yield and haulm yield which was significantly superior over Kadiri Chitravati, Dheeraj and TAG-24 and lowest pod and haulm yield was recorded with TAG-24. Highest N, P and K uptake of plant was recorded with irrigation scheduled at IW/CPE ratio of 1.0 along with Kadiri Lepakshi compared to the other treatments.

**Key words:** *Irrigation schedules, groundnut varieties, IW/CPE, yield, nutrient uptake*

### INTRODUCTION

The most significant oilseed crop in India is groundnut (*Arachis hypogaea* L.). India is the second-biggest producer in the world after China and the largest nation by area. 9.25 million tonnes are produced annually on 4.9 million hectares of land, with an average productivity of 1893 kg ha<sup>-1</sup>. India's top producer, Gujarat, accounts for 43% of the country's total output, followed by Rajasthan (13.76%), Andhra Pradesh (12.28%), Tamil

Comment [S1]: you should enter the problem

Comment [S2]: customize with journal template

Comment [S3]: better complete with numbers from the results of the discussion

Comment [S4]: you should complete it with a reference source

32 Nadu (10.55%), and Karnataka (9.55%). (5.14 per cent).

33 Andhra Pradesh grows groundnut on 1.01 million hectares, generating 0.60 million tonnes  
34 at a productivity of 1497 kg ha<sup>-1</sup> (Ministry of Agriculture and Farmers Welfare, 2020-  
35 2021). The reason why groundnut productivity is so low in comparison to the global  
36 average is primarily due to the fact that it is grown under moisture stress conditions at  
37 different growth stages, regardless of the production environment, irrigation technique,  
38 variety, and other cultivation practices (Thiyagarajan *et al.*, 2010). It is also grown in  
39 acidic soils with low levels of N, P, Ca, S, and B as well as insufficient organic matter  
40 (Noman *et al.*, 2015). Water use efficiency for this crop under irrigated conditions is low  
41 due to improper irrigation management.

42 The reduction in yield will be greater if severe stress occur during the critical crop  
43 growth stages like flowering and pod formation (Saha and Gunri, 2014). Thus the water  
44 management is most important factor because groundnut has specific moisture need due to  
45 the unique feature of developing the pods underground (Baliarsingh and Mahapatra, 2015).  
46 Proper irrigation scheduling helps the crop to put good crop growth and yield  
47 (Thiyagarajan *et al.*, 2010).

48 For scheduling irrigation to groundnut crops in different seasons and soil types,  
49 various approaches have been advocated. The evaporative demand from the atmosphere  
50 has grown in importance as the primary factor in determining crop water requirements, for  
51 which scheduling irrigation to groundnut crops on the basis of a climatological approach  
52 based on the IW/CPE ratio (IW- Irrigation Water, CPE- Cumulative Pan Evaporation) has  
53 been found to be most appropriate at the present time. This method incorporates all of the  
54 weather parameters that influence crop water use and is expected to increase output by at  
55 least 15- 20%. Irrigation scheduling optimization resulted in increased pod yield and water  
56 use efficiency (Taha and Gulati, 2001).

57 Keeping this in view, the present study was undertaken to investigate the influence of  
58 irrigation schedules on yield and nutrient uptake of groundnut varieties.

## 59 MATERIALS AND METHODS

60 The field experiment was conducted at Agricultural College farm, Bapatla during the  
61 *rabi* season of 2021-22. The experiment site was a sandy loam soil with neutral in reaction,  
62 low in available nitrogen and organic carbon content, high in available phosphorous and  
63 medium in available potassium. The total amount of rainfall received during the crop

Comment [S5]: should be completed with a map of the research carried out

64 growth period was 374.7 mm in 21 rainy days. The bulk density of soil at 15 cm depth was  
65  $1.51 \text{ g cm}^{-3}$ . Moisture percentage at field capacity and permanent wilting point was 14.5%  
66 and 7.0%. The experiment was laid out in split plot design with three replications. The main  
67 plots consisting of three irrigation schedules viz., M<sub>1</sub>- IW/CPE ratio of 1.0, M<sub>2</sub>- IW/CPE  
68 ratio of 0.8 and M<sub>3</sub>-IW/CPE ratio of 0.6 and sub plots consisting of four groundnut  
69 varieties viz., V<sub>1</sub>- TAG-24, V<sub>2</sub>-Dheeraj, V<sub>3</sub>-Kadiri Lepakshi and V<sub>4</sub>-Kadiri Chitravati. The  
70 crop was sown at a spacing of 22.5 cm × 10 cm. Recommended N, P and K applied to all  
71 the treatments uniformly @ 30: 40: 50 kg ha<sup>-1</sup>. Nitrogen and phosphorus applied through  
72 urea and SSP, potassium through muriate of potash. Whole quantity of the phosphorus and  
73 half of the nitrogen and potassium applied as basal and remaining half of nitrogen and  
74 potassium as top dressing at 25-30 DAS.

75 Irrigation scheduling was done using a climatological approach (IW/CPE). The open  
76 pan evaporimeter was used to record daily pan evaporation. The total amount of water  
77 applied to the crop was 410 mm, 340 mm, and 300 mm in IW/CPE ratios of 1.0, 0.8, and  
78 0.6, respectively. Four, three, and two irrigations were given to irrigation schedules of  
79 IW/CPE ratio of 1.0, 0.8, and 0.6, respectively along with the pre-sowing irrigation to all  
80 the treatments. In each treatment, the irrigation depth was kept constant at 50 mm per  
81 irrigation. A measured amount of water was given to each treatment using a Parshall flume  
82 with a capacity of 1 cusec (Parshall, 1950). The volume of water to be given for each  
83 treatment is calculated from the formula.

84 
$$\text{Volume} = \text{Area} \times \text{Depth}$$

85 Plant and kernel samples taken at maturity were analyzed for nitrogen (Modified micro  
86 kjeldhal method, Piper, 1966), phosphorus (Vanadomolybdo phosphoric acid method,  
87 Jackson, 1973), potassium (Flame photometer method, Jackson, 1973). From the chemical  
88 analysis data, uptake of the individual nutrient was calculated as shown below. Uptake was  
89 calculated by multiplying the nutrient content by the respective dry weight of kernel and  
90 haulm and then summed up to represent total nutrient uptake at harvest and expressed as kg  
91 ha<sup>-1</sup>.

92 Nutrient content (%) × Dry weight of kernel/haulm (kg ha<sup>-1</sup>)

93 Nutrient uptake (kg ha<sup>-1</sup>) = \_\_\_\_\_

94

100

95 The data on the pod and haulm yield was estimated after harvest of the crop. The data  
96 recorded on various parameters of crop was subjected to statistical scrutiny by the method of  
97 analysis of variance outlined by Panse and Sukhatme (1985).

UNDER PEER REVIEW

98

## 99 RESULTS AND DISCUSSION

100

### 101 Yield of Groundnut

#### 102 Pod yield(kg ha<sup>-1</sup>)

103 Among the irrigation schedules, the IW/CPE ratio of 1.0 (M<sub>1</sub>) produced higher pod yield  
104 (3175 kg ha<sup>-1</sup>) (Table-1) than the IW/CPE ratio of 0.6 (2916 kg ha<sup>-1</sup>) and was comparable to  
105 the IW/CPE ratio of 0.8 (M<sub>2</sub>) (2579 kg ha<sup>-1</sup>). This is most likely due to favourable soil  
106 moisture conditions and improved soil moisture availability throughout the crop growth  
107 period, which significantly stimulated yield attributes and ultimately pod yield. Similar  
108 findings were reported by Shaikh *et al.* (2004), Suresh *et al.* (2013) and Behera *et al.* (2015).  
109 Among the varieties, Kadiri Lepakshi produced the highest pod yield (3607 kg ha<sup>-1</sup>) and was  
110 significantly superior to Kadiri Chitravati, Dheeraj and TAG-24. These increased yield  
111 attributes could be attributed to increased growth parameters such as branch number and  
112 biomass production. The current findings are consistent with those of Mohite *et al.* (2017) and  
113 Naik *et al.* (2018).

#### 114 Haulm yield (kg ha<sup>-1</sup>)

115 The data (Table-1) revealed that irrigation scheduled at IW/CPE ratio of 1.0 (4291 kg  
116 ha<sup>-1</sup>) recorded higher value of haulm yield, which was significantly superior over IW/CPE  
117 ratio of 0.6 (M<sub>3</sub>) (4034 kg ha<sup>-1</sup>) but found statistically on a par with IW/CPE ratio of 0.8 (M<sub>2</sub>)  
118 (3681 kg ha<sup>-1</sup>). However, the lowest haulm yield was recorded with IW/CPE ratio of 0.6.  
119 This might be attributed to maintenance of adequate available soil moisture in the root zone  
120 coinciding with critical growth stages of crop would have helped for proper uptake as well as  
121 utilization of nutrients and created a favourable impact on growth as well as yield components  
122 leading to better haulm yield of the crop. Similar results were also reported by Bandopadhyay  
123 (2005) and Chitodkar *et al.* (2006).

124 Irrigation scheduled at 0.6 IW/CPE treatment recorded the lowest haulm yield (3681  
125 kg ha<sup>-1</sup>). Reduction in plant height, branches per plant, dry matter accumulation and canopy  
126 development due to moisture stress ultimately reduced the haulm yield of groundnut. Similar  
127 results were also reported by Sounda *et al.* (2006).

128 Among the varieties, Kadiri Lepakshi recorded significantly highest haulm yield over  
129 Kadiri Chitravati, Dheeraj and TAG-24. Whereas, Dheeraj and Kadiri Chitravati were

Comment [S6]: It is better to check the data entered again

Comment [S7]: should complete with values from the table

Comment [S8]: It is better to check the data entered again

130 comparable with each other. The highest haulm yield by KadiriLepakshi might be due to the  
131 genetic makeup of the genotype besides the environmental conditions. The results revealed in  
132 the present study are in confirm with findings of Nirmal *et al.* (2015).

### 133 **Nutrient uptake:**

134 N, P and K uptake by groundnut varieties estimated in kernel and haulm at harvest was  
135 significantly influenced by the irrigation schedules, while interaction effect was not  
136 statistically significant.

137 Total nutrient uptake includes nutrient uptake by kernel and haulm. Total N, P and K uptake  
138 was maximum when irrigation was scheduled at 1.0 IW/CPE ratio (104.2 kg ha<sup>-1</sup>, 11.8 kg ha<sup>-1</sup>  
139 and 54.3 kg ha<sup>-1</sup>, respectively) and was significantly superior than IW/CPE ratios of 0.8 and  
140 0.6. When compared to other treatments, irrigation with an IW/CPE ratio of 0.6 (78.1 kg ha<sup>-1</sup>,  
141 8.2 kg ha<sup>-1</sup> and 44.8 kg ha<sup>-1</sup>, respectively) resulted in the lowest N, P and K uptake. Because  
142 N, P and K uptake in plants is a function of yield and concentration, significant uptake by the  
143 plant may have resulted in higher yields. A consistent and adequate supply of moisture  
144 throughout the crop growth period could be one reason for increased nutrient availability for  
145 higher uptake and progressive utilisation by the crop, which in turn modified to  
146 produce incremental photosynthates for better partitioning of dry matter from source to sink.  
147 These results are in conformity with the findings of Patel *et al.* (2009), Naresha *et al.* (2018)  
148 and Verma *et al.* (2015).

149 Highest uptake of N, P and K found with Kadiri Lepakshi (109.4 kg ha<sup>-1</sup>, 12.7 kg ha<sup>-1</sup>  
150 and 58.1 kg ha<sup>-1</sup>, respectively) and was significantly superior over Kadiri Chitravati (V<sub>4</sub>),  
151 Dheeraj (V<sub>2</sub>) and TAG-24 (V<sub>1</sub>). The lowest uptake of nitrogen was found with TAG-24 (70.4  
152 kg ha<sup>-1</sup>, 7.3 kg ha<sup>-1</sup> and 41.6 kg ha<sup>-1</sup>, respectively) variety. N, P and K uptake of varieties is  
153 mostly governed based on the amount of dry matter produced besides the concentration of  
154 nutrient in various plant parts which corroborated the findings of Mohapatra and Dixit (2010)  
155 and Yadav *et al.* (2015). The higher uptake of N, P and K was recorded with Kadiri Lepakshi,  
156 this might be due to the inherent characteristic feature of the Kadiri Lepakshi variety to absorb  
157 greater quantity of nutrients, and among the varieties, variation in uptake may also be due to  
158 nitrogen content of different cultivars.

159

160

161  
162  
163  
164  
165  
166  
167

**Table 1. Pod yield and haulm yield (kg ha<sup>-1</sup>) of groundnut varieties as influenced by irrigation schedules.**

Comment [S9]: you should enter data source and year

Treatments	Pod yield (kg ha <sup>-1</sup> )	Haulm yield (kg ha <sup>-1</sup> )
<b>IRRIGATION SCHEDULES (M)</b>		
M <sub>1</sub> : IW/CPE ratio of 1.0	3175	4291
M <sub>2</sub> : IW/CPE ratio of 0.8	2916	4034
M <sub>3</sub> : IW/CPE ratio of 0.6	2579	3681
SEm ±	84.8	74.5
CD (p=0.05)	333	293
CV (%)	10.2	6.5
<b>GROUNDNUT VARIETIES (V)</b>		
V <sub>1</sub> : TAG-24	2074	3424
V <sub>2</sub> : Dheeraj	2694	3835
V <sub>3</sub> : Kadiri Lepakshi	3607	4647
V <sub>4</sub> : Kadiri Chitravati	3185	4101
SEm±	110.3	122.2
CD(p=0.05)	328	363
CV(%)	11.5	9.2
INTERACTION (M × V)	NS	NS

168  
169  
170  
171  
172

173  
 174  
 175  
 176  
 177  
 178  
 179

**Table 2. Nitrogen uptake (kg ha<sup>-1</sup>) of groundnut varieties as influenced by irrigation schedules.**

Treatments	Nitrogen uptake at harvest (kg ha <sup>-1</sup> )		
	Haulm uptake	Kernel uptake	Total uptake
<b>IRRIGATION SCHEDULES (M)</b>			
M <sub>1</sub> : IW/CPE ratio of 1.0	43.9	60.3	104.2
M <sub>2</sub> : IW/CPE ratio of 0.8	41.5	50.0	91.5
M <sub>3</sub> : IW/CPE ratio of 0.6	37.6	40.6	78.1
SEm ±	0.50	1.01	0.97
CD (p=0.05)	1.97	3.98	3.82
CV (%)	4.24	6.98	3.69
<b>GROUNDNUT VARIETIES (V)</b>			
V <sub>1</sub> : TAG-24	34.9	35.4	70.4
V <sub>2</sub> : Dheeraj	40.3	45.6	85.9
V <sub>3</sub> : Kadiri Lepakshi	46.3	63.0	109.4
V <sub>4</sub> : Kadiri Chitravati	42.4	57.1	99.5
SEm±	0.78	1.70	2.08
CD(p=0.05)	2.31	5.04	6.18
CV(%)	5.69	10.12	6.84
INTERACTION (M × V)	NS	NS	NS

Comment [S10]: you should enter data source and year

180  
 181  
 182  
 183

184  
185  
186  
187

**Table 3. Phosphorus uptake (kg ha<sup>-1</sup>) of groundnut varieties as influenced by irrigation schedules.**

Treatments	Phosphorus uptake at harvest (kg ha <sup>-1</sup> )		
	Haulm uptake	Kernel uptake	Total uptake
<b>IRRIGATION SCHEDULES (M)</b>			
M <sub>1</sub> : IW/CPE ratio of 1.0	4.7	7.1	11.8
M <sub>2</sub> : IW/CPE ratio of 0.8	4.0	6.2	10.2
M <sub>3</sub> : IW/CPE ratio of 0.6	3.3	5.0	8.2
SEm ±	0.09	0.10	0.20
CD (p=0.05)	0.37	0.40	0.79
CV (%)	8.17	5.84	6.88
<b>GROUNDNUT VARIETIES (V)</b>			
V <sub>1</sub> : TAG-24	2.9	4.4	7.3
V <sub>2</sub> : Dheeraj	3.9	5.5	9.3
V <sub>3</sub> : Kadiri Lepakshi	5.1	7.6	12.7
V <sub>4</sub> : Kadiri Chitravati	4.2	6.9	11.0
SEm±	0.15	0.18	0.25
CD(p=0.05)	0.44	0.54	0.75
CV(%)	11.04	8.98	7.5
INTERACTION (M × V)	NS	NS	NS

**Comment [S11]:** you should enter data source and year

188  
189  
190  
191  
192  
193

194  
195  
196  
197  
198

**Table 4. Potassium uptake (kg ha<sup>-1</sup>) of groundnut varieties as influenced by irrigation schedules.**

Treatments	Potassium uptake at harvest (kg ha <sup>-1</sup> )		
	Haulm uptake	Kernel uptake	Total uptake
<b>IRRIGATION SCHEDULES (M)</b>			
M <sub>1</sub> : IW/CPE ratio of 1.0	40.6	13.7	54.3
M <sub>2</sub> : IW/CPE ratio of 0.8	38.3	12.0	50.3
M <sub>3</sub> : IW/CPE ratio of 0.6	35.2	9.6	44.8
SEm ±	0.56	0.36	0.64
CD (p=0.05)	2.20	1.43	2.50
CV (%)	5.09	10.73	4.42
<b>GROUNDNUT VARIETIES (V)</b>			
V <sub>1</sub> : TAG-24	33.5	8.05	41.6
V <sub>2</sub> : Dheeraj	36.4	10.6	46.9
V <sub>3</sub> : Kadiri Lepakshi	43.2	14.9	58.1
V <sub>4</sub> : Kadiri Chitravati	39.2	13.6	52.6
SEm±	0.94	0.41	1.04
CD(p=0.05)	2.80	1.21	3.09
CV(%)	7.43	10.38	6.27
INTERACTION (M × V)	NS	NS	NS

**Comment [S12]:** you should enter data source and year

199 **Conclusion:**

200 From the present investigation it can be concluded that highestpod and haulm yield of  
201 groundnut was recorded with IW/CPE ratio of 1.0 (M<sub>1</sub>) which was significantly superior over  
202 IW/CPE ratio of 0.6 but was on par with IW/CPE ratio of 0.8. Among the varieties, Kadiri  
203 Lepakshi recorded significantly higher Pod and haulm yield over Kadiri Chitravati, Dheeraj  
204 and TAG-24. N, P and K uptake of groundnut varieties was found significantly highest with

**Comment [S13]:** complete with numbers from the results of the discussion

205 IW/CPE ratio of 1.0 along with Kadiri Lepakshi variety compared to other treatments. The  
206 lowest N, P and K uptake was recorded with 0.6 IW/CPE ratio along with TAG-24 variety.

## 207 **Literature cited**

208 Bandyopadhyay, P.K., Mallick, S and Rana, S.K. 2005. Water balance and crop coefficients of  
209 summer-grown peanut (*Arachis hypogaea* L.) in a humid tropical region of  
210 India. *Irrigation Science*. 23(4): 161-169.

211  
212 Baliarsingh, A and Mahapatra, P.K. 2015. Effect of date of sowing and irrigation on yield and  
213 moisture use dynamics of summer groundnut. *Environment and Ecology*. 33:175-179.

214  
215 Behera, B.S., Mohit, D.A.S., Behera, A.C and Behera, R.A. 2015. "Weather based irrigation  
216 scheduling in summer groundnut in Odisha condition". *International Journal of*  
217 *Agricultural Science and Research*. 5(5): 247-259.

218 Chitodkar, S.S., Choudhar, P.M., Patil, H.E and Raunal, P.U. 2006. Studies on irrigation  
219 regimes, mulches and antitranspirant on yield and water management of summer  
220 groundnut (*Arachis hypogaea* L.). *International Journal of Agricultural Sciences*. 2(2):  
221 496-498.

222 Jackson, M.L. 1973. *Soil chemical analysis*. Prentice Hall India Private Limited New Delhi pp:  
223 498.

224 Ministry of Agriculture and Farmers Welfare Government of India. [www.indiastat.com](http://www.indiastat.com) 2020-  
225 2021.

226  
227 Mohapatra, A.K.B and Dixit, L. 2010. Integrated nutrient management in rainy season  
228 groundnut. *Indian Journal of Agronomy*. 55(2):123-127.

229 Mohite, U.A., Mohite, A.B and Jadhav, Y.R. 2017. Effect of sowing windows on growth and  
230 yield of groundnut varieties during *kharif* season. *Contemporary Research in India*. 7:  
231 189-192.

232 Naik, A.K., Pallavi, N and Sannathimmappa, H.G. 2018. Performance of different spanish-type  
233 groundnut varieties suitable under Central dry zone of Karnataka, India. *International*  
234 *Journal of Current Microbiology and Applied Sciences*. 7: 1394-1397.

235 Naresha, R., Laxminarayana, P., Devi, K.S and Narender, J. 2018. Effect of moisture regimes  
236 and phospho-gypsum levels on yield, nutrient uptake and soil nutrient balance of *rabi*  
237 groundnut. *International Journal of Agriculture, Environment and Biotechnology*.  
238 10(4): 489-498.

239  
240 Nirmal De, Meena, R.S., Yadav, R.S., Reager, M.L., Meena, V.S., Verma, J.P., Verma, S. K  
241 and Kansotia, B. C. 2015. Temperature use efficiency and yield of groundnut varieties  
242 in response to sowing dates and fertility levels in western dry zone of India. *American*  
243 *Journal of Experimental Agriculture*. 7(3): 170-177.

Comment [S14]: customize with journal template

- 244 Noman, H.M., Rana, D.S and Rana, K.S. 2015. Influence of sulphur and zinc levels and zinc  
245 solubilizer on productivity, economics and nutrient uptake in groundnut. *Indian Journal*  
246 *of Agronomy*. 60(2): 301-306.
- 247 Panse, V. Gand Sukhatme, P.V. 1985. Statistical Methods for Agricultural Workers. *Indian Council*  
248 *of Agricultural Research*. New Delhi. 205-210
- 249 Parshall, R.L. 1950. Measuring water in irrigation channels with parshall flumes  
250 and small weirs. USDA, Circular No. 843. Pp: 500.
- 251 Patel, G.N., Patel, P.T., Patel, P.H., Patel, D.M., Patil, D.K and Patil, R.M. 2009. Attributes,  
252 yield, quality and uptake of nutrients by summer groundnut (*Arachis hypogaea* L.) as  
253 influenced by sources and levels of sulphur under varying irrigation schedules. *Journal*  
254 *of Oilseeds Research*. 26(2): 119-122.
- 255 Piper, C. S. 1966. *Soil and Plant Analysis*. Hans Publishers, Bombay. 15-39.
- 256 Saha, D and Gunri, S.K. 2014. Effect of polythene mulching, irrigation regimes and fertilizer  
257 dose on summer groundnut nodulation, pod yield and nitrogen availability in  
258 soil. *Environment and Ecology*. 32: 1301-1303.
- 259 Shaikh, A.A., Nimbalkar, C.A and Jawale, S.M. 2004. Effect of irrigation scheduling and  
260 mulches on yield contributing characters of summer groundnut. *Journal of Maharashtra*  
261 *Agricultural University*. 299(2): 163-166.
- 262 Sounda, G., Mandal, A., Moinuddin, G and Mondal, K. 2006. Effect of irrigation and mulch on  
263 yield, consumptive use of water and water use efficiency of summer groundnut. *Journal*  
264 *of Crop and Weed*. 2(1): 29-32.
- 265 Suresh, K., Balaguravaiah, D., Ramulu, V and Sujani Rao, C.H. 2013. Comparative efficiency  
266 of sprinkler irrigation over check basin irrigation in groundnut at different irrigation  
267 schedules. *International Journal of Plant Animal and Environmental Sciences*. 3(2): 9-  
268 13.
- 269 Taha, M and Gulati, J.M.L. 2001. Influence of irrigation on yield and moisture utilization of  
270 groundnut (*Arachis hypogaea* L.). *Indian Journal of Agronomy*. 46 (4): 523-527.
- 271 Thiyagarajan, G., Ranghaswamy, M.V., Rajkumar, D and Kumaraperumal, R. 2010. Deficit  
272 irrigation effects on groundnut with micro sprinklers. *Madras Agriculture Journal*  
273 97: 40-42.
- 274 Verma, S.K., Singh, S.B., Prasad, S.K., Meena, R.N and Meena, R.S. 2015. Influence of  
275 irrigation regimes and weed management practices on water use and nutrient uptake in  
276 wheat. *Bangladesh Journal of Botany*. 44(3): 437-442.
- 277 Yadav, G.S., Datta, M., Babu, S., Saha, P and Singh, R. 2015. Effect of sources and levels of  
278 phosphorus on productivity, economics, nutrient acquisition and phosphorus-use

279 efficiency of groundnut under hilly ecosystems of North-East India.*Indian Journal of*  
280 *Agronomy*. 60(2): 307-311.

281

282

283

284

285

286

287

288

289

290

UNDER PEER REVIEW

