

## Data Article

### **Effect of Organic and Liquid Manures on Yield and Economic Attributes of Urdbean.**

#### **Abstract**

A field experiment was carried out in 2020 kharif season, on loamy sand soil at the S.K.N. College of Agriculture's Agronomy Farm in Jobner (Rajasthan). It included two foliar sprays of each of four liquid manures (control, panchgavya @ 5%, vermiwash @ 10%, and cow urine @ 5%), totaling sixteen treatment combinations. It included four treatments of organic manures (control, 100% RDP through FYM, through vermicompost, and through poultry manure) and two foliar sprays of each of four liquid manures. Three replications of the experiment were set up using a Randomized Block Design. 'Krishna' variety of urdbean was sown as on July 13, 2020, a test crop with a 30-cm row spacing. The results showed that two foliar sprays of 5% panchgavya were the most effective liquid manure treatment in raising the maximum seed production (942 kg/ha), straw yield (2,121 kg/ha), and biological yield (3,063 kg/ha) of urdbean. The highest levels of nitrogen, phosphorous, and potassium were found in seed and straw under this liquid manure treatment, which was comparable to two foliar sprays of 10% vermiwash. However, in terms of economics, it was discovered that applying 100% RDP through FYM was the most lucrative strategy, yielding the highest net returns of '36,314/ha and a B:C ratio of 2.68. Foliar spraying with 5% panchgavya also generated an extra 36,798/ha in net returns over the control, which had the highest B:C ratio of 2.37.

**Keyword:** Urdbean, liquid manure, net returns, organic manures.

#### **Introduction**

Urdbean [*Vigna mungo* (L.) Hepper] is India's third-most significant pulse crop. It comes from the wild plant *Phaseolus sub lobatus* and is indigenous to India. It is typically eaten in the form of "Dal." It is the main ingredient in "papad," "idly," and "dosa." It is high in phosphoric acid and contains protein (25%), carbs (60%), fat (1.3%), and fat. Pulses provide a significant fraction of the world's population with nutritional protein, and they also help to avert climate change by enhancing soil health because to their amazing nitrogen-fixing abilities and by adding a lot of residue to the soil. The greatest user (29% of global consumption), producer (25-28% of global output), and importer of pulses worldwide is India. During the 2019–2020 growing season in India, pulses would occupy close to 29.36 million hectares, producing 23.02 million tonnes at an average productivity of 779 kg/ha (Anonymous, 2019-20).

It accounts for 10% of India's overall pulse production and 13% of the country's total pulse area. It is grown on an average of 655 kg/ha on an area of 5.44 mha in India, producing 3.56 mt (Anonymous, 2019). States with significant urdbean production include Madhya Pradesh, Rajasthan, Andhra Pradesh, Uttar Pradesh, Tamil Nadu, Maharashtra, Jharkhand, and Gujarat. Pulses are grown on a total of 38.42 lakh hectares in Rajasthan, where they are produced in 17,80 lakh tonnes at an average productivity of 463 kg/ha. In Rajasthan, urad

bean cultivation covers 0.95 million hectares, producing 0.34 million tonnes on an average per ha (Anonymous 2019-20). It is primarily farmed in dry and semi-arid regions of Rajasthan, including Jhalawar, Kota, Bundi, Baran, Chittorgarh, Udaipur, and Ajmer.

Among organic manures, FYM is a good source of plant nutrients, being high in organic matter and containing 0.5% N, 0.25% P, and 0.5% K as well as a variety of micronutrients in varying amounts. It breaks down into organic acids, which make nutrients available to plants in growth. It aids in protecting soils from abrupt chemical changes. Additionally, FYM serves as a source of energy for the expansion of soil bacteria.

Earthworms are used to create vermicompost, an organic source of nutrients that can be recycled from organic waste. It contains a lot of nutrients, including 7.61% Ca, 0.56% Mg, 0.11% Zn, and 1.33% Fe, as well as 1.67% N, 1.2% P, and 1.0% K. Along with adding organic carbon to the soil and aiding in the slow and efficient release of nutrients to the rooting zone of crop plants, it also contains a healthy proportion of exchangeable Ca, Mg, Na, and other elements.

Manure from poultry is largely resistant to microbial deterioration. However, due to its bulky character and the fact that inorganic fertilizer is so expensive, it is no longer within the means of farmers with little resources. It is necessary for creating and maintaining the best soil physical condition, which is required for plant growth (Rahman, 2004). In that order, it has 1.30 % N, 1.80 % P, and 0.80 % K.

Cow secretions like dung, urine, milk, curd, and ghee are used to make inexpensive, environmentally friendly organic fermented preparations called panchgavya and cow urine. These preparations are excellent sources for foliar fertilizing crops. It is a powerful stimulator of plant development that raises the biological effectiveness of crops. It is utilized to energize the soil, safeguard plants from disease, and improve the nutritional level and caliber of plants' food. Its NPK concentration is low, coming in at 0.10, 0.017, and 0.019%, respectively. Along with irrigation water, it is applied to the soil and utilized as a foliar spray, as well as for treating seeds and seedlings, among other things.

Nitrogen, phosphate, potassium, magnesium, calcium, sulphate, and chloride are all abundant in cow urine. It is composed primarily of water (95%) and various minerals, including salts, enzymes, and urea (2.5%). In addition to acting as a plant hormone, it has been claimed to help plants that are lacking in certain micronutrients. It has a 6.8 to 21.1 g N content per liter (Nagar et al., 2016).

Numerous growth-regulating chemicals are present in vermiwash made from earthworm beds (Nielson, 1965). After water has passed through several layers of earthworm culture units, it is collected as a clear, pale yellow liquid extract of organic waste components. 1.34, 0.30, and 2.0% of N, P, and K, respectively, are present in vermiwash. As a foliar spray, it works incredibly well. Additionally, the earthworms' enzymes and secretions found in the vermiwash encourage crop development and productivity. Inorganic fertilizers are becoming prohibitively expensive, making them unaffordable for small and marginal farmers. When inorganic fertilizers are used, the number of helpful organisms declines, the soil's natural processes for regenerating nutrients stop, the soil turns barren, and the fertility

of the soil declines. The use of these organic liquid formulations will increase soil microbial population and activity to a greater level, which will benefit crop development and yield.

## Materials and Methods

During the 2020 kharif season, a field experiment named "Effect of Organic and Liquid Manures on Growth and Yield of Urdbean [*Vigna mungo* (L.) Hepper]" was carried out at the agronomy farm of the S.K.N. College of Agriculture, Jobner. Jobner is located 45 kilometres (km) west of Jaipur, at 26°05' North latitude, 75°28' East longitude, and 427 metres above mean sea level. The region is located in Rajasthan's agro-climatic Zone-IIIa (Semi-arid Eastern Plain Zone). The data showed that the crop season saw 256.7 mm of rainfall, the majority of which was recorded in August. During the Urdbean growth season, the mean daily maximum and minimum temperatures varied from 30.2 to 40.9 C<sup>0</sup> and 14.3 to 25.4 C<sup>0</sup>, respectively. Similar to that, the mean daily relative humidity ranged from 43% to 87%. There were between 1.8 and 8.0 hours of sunshine per day on average. Before layout, soil samples were collected from the experimental area in five random locations, ranging in depth from 0 to 15 cm. A representative composite sample was then created by combining and processing all of the soil samples. To determine its physico-chemical properties, the homogeneous composite soil sample underwent mechanical, physical, and chemical analyses. The experimental field had a loamy sand texture, a pH of 8.20, an EC of 1.41, an alkaline reaction, a poor organic carbon content of 0.15, a low available nitrogen level of 128.0, a medium available phosphorus content of 16.12, and a high potassium content of 153.4.

A drought-tolerant urdbean cultivar called Krishna was employed in the trial. The treatment consisted of 4 organic manures, [(control 100% RDP through FYM 100% RDP through vermicompost 100% RDP through poultry manure)]; it was set up in a factorial randomised block design. and four liquid manures (Control, Panchgavya (two sprays at 5%), Vermiwash (two sprays at 10%), and cow urine (two sprays at 5%), totaling sixteen treatments, were used. This experiment was repeated three times. The trial field was extensively prepared by ploughing and then deep-ploughing with a tractor-drawn disc plough. Following planking to achieve fine tilth, treatments such as well-rotted FYM, vermicompost, and poultry manure were applied and well assimilated into the soil. Using a hand plough and a seed rate of 20 kg/ha, the seeds were sown according to the "Kera" method, with a row spacing of 30 cm. Azospirillum was uniformly applied to crop seed at a rate of 750 g/ha, and VAM and

Trichoderma were applied to soil at a rate of 5 kg/ha for all treatments. To enable the best plant population, thinning was carried out 15–20 days after sowing, with a 10 cm intra-row spacing. Through the sprinkler system, the crop received two life-saving irrigations. Using the battery-operated knapsack sprayer in the early morning and late at night, liquid manures, including panchgavya, vermiwash, and cow urine at 5%, 10%, and 5% respectively, and water at 500 l/ha, were sprayed twice: once at 25 DAS and again 15 days later at the flowering initiation stage. Vermiwash 50 liters/ha, panchgavya 25 liters/ha, and cow urine 25 liters/ha were the amounts of liquid manures used for each spraying. The crop was cut down on September 24, 2020, in order to evaluate the biological, seed, and stalk yields. Harvesting from the net plot area was completed after leaving two rows on each side and 0.5 m across each row on both sides (3.0 m x 1.8 m). Bundles were hauled from each plot to the threshing floor for appropriate sun drying after being properly knotted and labelled.

## **Result**

### **4.4.4 Economics of the treatments**

#### **4.4.1 Net returns**

**Organic manures:** The data shown in table 1 and fig. 1 make it clear that all organic manure treatments greatly outperformed the control in terms of net returns. The most lucrative strategy yielded the best net returns of '36,314/ha and applied 100% RDP through FYM. By a margin of 3,862 and 10,037/ha over control, it improved the net returns, respectively. However, it was discovered that an additional net return of '8,910/ha was obtained that was notably similar to the control. 100% RDP through vermicompost enhanced the net returns by 23.7% over control, making it comparable to poultry manure.

**Liquid manures:** Due to the application of various liquid manures by foliar spray, net returns were also greatly increased (Table 1 and Fig 1). The two foliar sprays of 5% panchgavya yielded the greatest net yields of \$36 798/ha, which is a considerable increase of 11.1 and 47.2 percent over the two sprays of 5% cow urine and control, respectively. Statistics with panchgavya spray were shown after two foliar sprays of 10% vermiwash also produced increased net returns of '10,344 over the control. On the other side, a 5% sprinkle of cow pee might boost net returns by 32.4% over control.

#### **4.4.2 B:C ratio**

**Organic manures:** The B:C ratio in urdbean was dramatically improved by applying several organic manures (Table ). When 100% RDP was applied using FYM, the B:C ratio reached

its peak at 2.68, which was 17.0% higher than control. 100% RDP through poultry manures was the next superior therapy in this regard, giving 2.25 rupees for every rupee invested.

**Liquid manures:** Different liquid manures applied to the leaves significantly improved the B:C ratio. The highest B:C ratio was recorded by a foliar spray of 5% panchgavya, which was closely followed by two foliar sprays of 10% vermiwash and 5% cow urine.

The most profitable amount of nutrients in urdbean were discovered to be from organic manures such as vermicompost and poultry manure. A considerable improvement of 10.52 and 23.65 percent over 100% RDP through FYM and control, respectively, which was on par with 100% RDP through chicken manure, was achieved by vermicompost, yielding net yields of "32492/ha." In order to determine net return, the yields of seeds and straw are multiplied by their selling prices, and the overall cost of cultivation, including treatment costs, is then subtracted. Because this treatment had a lower additional cost than 100% RDP by vermicompost, where the B:C ratio further decreased to 1.96, the highest B:C ratio was reached with 100% RDP with FYM (2.68). Both Kumar et al. (2013) and Kumawat et al. (2013) showed similar outcomes for mungbean and urdbean, respectively.

#### **5.2.4 Economics**

Urdbean's net return (₹ 36798/ha) and B:C (2.37) ratio rose after being sprayed with panchagavya twice compared to the control and other treatments (Table 1). The higher seed (942 kg/ha) and straw production (2121 kg/ha) under the two sprays of panchagavya in the current study can be used to explain the higher net return. Both groundnut and blackgram by Kumawat et al. (2009) and Somasudaram et al. (2009) reported similar outcomes.

#### **Conclusion**

The most effective method for increasing urdbean seed and straw yields (969 and 2165 kg/ha) was to apply 100% RDP using vermicompost. While the application of 100% RDP through FYM yielded the highest net returns of ₹36,314/ha with a B:C ratio of 2.68 due to the lower market price. The most effective treatments, however, were two foliar sprays of 5% panchgavya, which produced the highest seed and straw yields. (942 and 2,121 kg/ha) with the maximum net returns of ₹ 36,798/ha and B:C ratio (2.37) of urdbean

**Table 1 : Effect of organic and liquid manures on net returns and B: C ratio of urdbean.**

<b>Treatments</b>	<b>Net returns (₹/ha)</b>	<b>B: C Ratio</b>
<b>Organic manures</b>		
Control	26,277	2.29
100% RDP through FYM	36,314	2.68
100% RDP through vermicompost	32,492	1.96
100% RDP through poultry manure	35,187	2.25
SEm±	9,79	0.06
CD (P=0.05)	2,827	0.17
<b>Liquid manures</b>		
Control	25,005	2.14
Panchgavya (2 sprays @ 5%)	36,798	2.37
Vermiwash (2 sprays @ 10%)	35,349	2.34
Cow urine (2 sprays @ 5%)	33,117	2.34
SEm±	979	0.06
CD (P=0.05)	2827	0.17
CV (%)	10.41	8.86

**Table 2 : Effect of organic and liquid manures on seed, straw and biological yield of urdbean (kg/ha) and harvest index (%)**

Treatments	Yield attributes		
	Number of pods/plant	Number of seeds/pod	Test weight (g)
<b>Organic manures</b>			
Control	19.11	5.22	29.86
100% RDP through FYM	22.65	5.98	32.32
100% RDP through vermicompost	24.42	6.36	33.90
100% RDP through poultry manure	24.13	6.16	33.76
SEm±	0.55	0.16	0.64
CD (P=0.05)	1.59	0.45	1.86
<b>Liquid manures</b>			
Control	18.88	5.18	29.87
Panchgavya (2 sprays @ 5%)	24.47	6.34	34.07
Vermiwash (2 sprays @ 10%)	24.13	6.28	33.79
Cow urine (2 sprays @ 5%)	22.84	5.92	32.11
SEm±	0.55	0.16	0.64
CD (P=0.05)	1.59	0.45	1.86
CV (%)	8.46	9.18	6.87

**Table 3 : Effect of organic and liquid manures on yield attributes of urdbean**

Treatments	Yield (kg/ha)			
	Seed yield	Straw yield	Biological yield	Harvest index (%)
<b>Organic manures</b>				
Control	679	1564	2243	30.27
100% RDP through FYM	845	1935	2780	30.39
100% RDP through vermicompost	969	2165	3134	30.91
100% RDP through poultry manure	929	2036	2965	31.33
SEm±	22	51	82	0.87
CD (P=0.05)	63	146	237	NS
<b>Liquid manures</b>				
Control	704	1592	2296	30.63
Panchgavya (2 sprays @ 5%)	942	2121	3063	30.73
Vermiwash (2 sprays @ 10%)	917	2052	2970	30.85
Cow urine (2 sprays @ 5%)	858	1935	2793	30.68
SEm±	22	51	82	0.87
CD (P=0.05)	63	146	237	NS
CV (%)	8.78	9.13	10.21	9.78



## References:

- Chhonkar, P.K. 2002. Soil Research in India. Some oversights and failures. *Journal of the Indian Society of Soil Science* **50**(4): 382-432.
- Choudhary, G.L. and Yadav, L.R. 2011. Effect of fertility levels of foliar nutrient on productivity of cowpea. *Journal of Food Legume* **24**(1): 67-68.
- Kudi, V.K., Singh, J.K., Choudhary, M., Koodi, H.L. and Jat, R. 2017. Effect of *Rhizobium*, PSB and fertility levels on nutrient content and their removal and yield of blackgram under custard apple based agri-horti system in Vindhyan region of Uttar Pradesh. *International Journal of Chemical Studies* **5**(3): 378-381.
- Kumar, R., Baba, A.Y., Manoranjan, K, Bhusan, A. and Kulveer, S. 2020. Assessment of organic and inorganic source of nutrients on yield and yield traits of black gram (*Vigna mungo* L.). *Journal of Pharmacognosy and Phytochemistry* **9**(3): 611-613.
- Kumar, R., Singh, Y., Choudhary, H.R. and Yadav, R.I. 2013. Nutrient uptake and profitability of *kharif* greengram [*Vigna radiata* (L.) Wilczek] as influenced by phosphorus levels and PSB under custard apple (*Annona squamosa*) based agri-horti-system. *Environment & Ecology* **31**(3): 1344-1346.
- Kumawat, P.K., Tiwari, R.C., Golada, S.L., Garhwal, R.K. and Choudhary, R. 2013. Effect of phosphorus sources, levels and biofertilizer on yield attributes, yield and economics of blackgram (*Phaseolus mungo*). *Legume Research* **36**: 70-73.
- Kumawat, R.N., Mahajan, S.S. and Mertia, R.S. 2009a. Growth and development of Groundnut (*Arachis hypogea*) under foliar application of panchgavya and leaf extracts of endemic plants. *Indian Journal of Agronomy* **54**(3): 324-331.
- Mathur, N., Singh, J., Bohra, S., Bohra, A., Solanki, R. and Vyas, A. 2008. Response of mothbean genotypes to nutrient management under arid conditions of Indian Thar desert. *Journal of Food Legumes* **21**: 69-70.
- Meena, J.S., Verma, H.P. and Pancholi, P. 2015. Effect of fertility levels and biofertilizer on growth and yield of cowpea on sandy loam soil of Rajasthan. *An Asian Journal of Soil Science* **10**(1): 55-58.

- Rajkhowa, D.J., Saikia, M. and Rajkhowa, K.M. 2002. Effect of vermicompost with and without fertilization of greengram. *Legume Research* **25**(4): 295-296.
- Shikha, J., Khaddar, V.K., Choudhary, S.K., Phadris, S. and Raehra, N. 2004. Effect of organic and chemical fertilizers on the growth yield attributes and yield of soybean-wheat cropping sequence. *Research on Crops* **5**(1): 22-30.
- Singh, K., Manohar, R.S., Choudhary, Yadav, A.K. and Sangwan, A. 2015. Response of different sources and levels of phosphorus on yield, nutrient uptake and net return on mungbean under rainfed condition. *Indian Journal of Agricultural Research* **35**: 263-268.
- Singh, A., Singh, V.K., Rana, N.S., Kumar, S., Panwar, G.S. and Kumar, Y. 2008. Response of urdbean to farmyard manure and phosphorus application under urdbean-wheat cropping sequence. *Journal of Food Legumes* **21**(2): 119-121.
- Singh, A. and Singh, J. 2016. Yield, nutrient uptake and soil fertility balance as influenced by organic and inorganic nutrient sources in urdbean- wheat cropping system. *International Journal of Agricultural Invention* **1**(1): 24-29.
- Sitaram, T., Sharama, S.K. and Reager, M.L. 2013. Growth attributes and nutrient uptake of greengram as influenced by vermicompost and zink in arid western Rajasthan. *Advance Research Journal of Crop Improvement* **4**(1): 65-69.
- Somasundarm, E., Meena, S., Sankaran, N. and Thiagarajan, T.M. 2003. Effect of panchgavya on growth and yield of mungbean. *National Symposium on Resource Management for Ecofriendly Crop Production (26-28)*: 71-72.
- Thiyageshwari, S., Rani, P. and Perumal, R. 2000. Changes in available phosphorus and grain yield of urdbean under integrated nutrient management in inceptisol. *Agropedology* **10**(1): 40-43.
- Yadav, O.S. 2001. Effect of nitrogen sources and biofertilizer on growth and quality of cowpea. M.Sc. (Ag.) Thesis, RAU, Bikaner.

UNDER PEER REVIEW