

# Integrated Pest Management of Brown Plant Hopper, *Nilaparvata lugens* (Stal) in rice of Mancherial District, Telangana State, India

## 1- Original Research Article:

### Abstract

A Front Line Demonstration (FLD) on Integrated Pest Management (IPM) of brown plant hopper, *Nilaparvata lugens* (Stal) in rice has been conducted in the adopted villages of Mancherial district of Telangana state during the *kharif* (June – December), season of 2018–19, 2019–20 and 2020–21 respectively to create awareness among the farming community on the IPM practices to manage the pest. The adoption of IPM practices includes formation of alleyways, recommended dose of nitrogen fertilizer, alternate wetting and drying, spraying of need based insecticides like Dinotefuran 20 SG @ 0.4 g or Pymetrozine 50 WDG @ 0.6 g l<sup>-1</sup> of water were carried out. The study reveals that the lowest hoppers incidence was witnessed in the demonstrated plot with 1.84, 12.18 and 19.0 adults hill<sup>-1</sup> in tillering stage and 7.76, 24.95 and 52.38 adults hill<sup>-1</sup> in panicle stage over farmers practice (3.33, 35.40 and 25.42 adults hill<sup>-1</sup> in tillering stage and 19.84, 50.30 and 63.47 adults hill<sup>-1</sup>). The average yields of IPM module demonstrated plot was 7130, 5513 and 5065 kg ha<sup>-1</sup> whereas in farmer practice the yield was 6733, 4851 and 4768 kg ha<sup>-1</sup> during *Kharif* (June–December), 2018–19, 2019–20 and 2020–21 respectively with an increased yield of 5.88%, 13.64% and 6.22% during corresponding *Kharif* (June–December), 2018–19, 2019–20 and 2020–21. Further the Cost-Benefit ratio of 1.88, 2.68 and 1.78 in the technology demonstrated plots whereas in farmers practice the recorded Cost-Benefit ratio of 1.43, 1.94 and 1.48 during corresponding *Kharif* (June–December), 2018–19, 2019–20 and 2020–21 respectively.

**Keywords:** Rice, Brown Plant Hopper, Integrated Pest Management, Alleyways, Insecticides, Alternate wetting and drying

## 1. Introduction

Rice (*Oryza sativa* L.) is an

important cereal crop of the world and second largest cultivated crop after wheat [1] and providing a staple food for more than half of the world population [2]. The crop is attacked by 300 insect pests in the different stages of the crop and among them only 23 species causes notable damage [3]. Among which brown plant hopper, *Nilaparvata lugens* (Stal), (Hemiptera: Delphacidae) is one of the most economically significant insect-pests of rice in most of the Asian countries causing a noticeable damage to the crop by reducing the yields ranges from 10%–90% and in Asia the loss have been estimated as more than \$300 million annually [4]. If timely control measures are not taken up, there may be a chance of total crop loss within a very short period [5]. In addition to this it also transmits the ragged stunt virus, striped virus and grassy stunt virus [6,7]. The search and evaluation of new insecticide molecules must also be a systematic practice so as to develop safer and effective alternatives to minimize the

brown plant hoppers damage [8,9]. This continuous and indiscriminate use of one insecticide has resulted in the rapid development of insecticide resistance and exhaustion of most insecticide options in many rice-growing regions [10] including organophosphates, carbamates, pyrethroids, neonicotinoids and phenylpyrazoles [11]. Insecticides remain, as a significant weapon in the integrated pest management (IPM) system because of its speedy, efficient, economical and easy to use against pests [12,13,14]. To overcome these lapses KVKs acts as a Knowledge and Resource Centre at district level to demonstrate the technologies [15] and the output of the research is disseminated to farmers through conduction of frontline demonstrations about the developed technologies [16]. For this IPM technologies need to be practiced in cluster approach to manage the pest [17]. Integrated pest management (IPM) practices is one of the eco-friendly approaches,

which can be applied to regulate the indiscriminate use of insecticides to control rice insect pests [18,19]. Hence, the following integrated pest management module under front line demonstrations were formulated with all integrated approaches like formation of alleyways, recommended dose of nitrogen fertilizer, alternate wetting and drying are being added along with chemical control to manage brown plant hopper, *Niaparvatha lugens* (Stal) in the present study conducted by Krishi Vigyan Kendra, Bellampalli, Mancherial District, which was undertaken during kharif (June–December), 2018–19, 2019–20 and 2020–21 respectively to manage the brown plant hopper, *Niaparvatha lugens* (Stal) by instigating integrated pest management practices in the farmer fields of villages in the various mandals of Mancherial district of Telangana state.

## 2. Materials and Methods

The present study was carried out in the various adopted villages

of KVK, Bellampalli in the Mancherial district during *kharif* (June–December), 2018–19, 2019–20 and 2020–21. In this study, 30 farmers (10 farmers/year) were selected for the demonstration of the technology over the three years. The improved technology were imposed, consisting formation of alleyways, recommended dose of nitrogen fertilizer, alternate wetting and drying, spraying of need based insecticides like Dinotefuran 20 SG @ 0.4 g or Pymetrozine 50 WDG @ 0.6 g l<sup>-1</sup> of water in the demonstrated plots. Whereas in the farmers practice, indiscriminate spraying of Acephate 75 SP @ 1.5 g, Imidacloprid 17.8 SL @ 0.25 ml, Buprofezin 25 SC @ 1.6 ml and Imidachloprid+Ethiprole 40 WG @ 0.25 g l<sup>-1</sup> etc. of which results in the heavy infestations of *N. lugens* (Farmers practice). The hopper population (nymph or adult) hill<sup>-1</sup> was recorded during the vegetative and reproductive stage of the crop after each spray

and 7 days after each application and the percent reduction over control of hoppers was calculated. Finally, the grain yield (kg ha<sup>-1</sup>) and cost - benefit ratios of demonstrated plots and farmers practice were also recorded and calculated.

### 3. Results and Discussion

Results revealed that the incidence of *N. lugens* was lower in the demonstrated plots with 1.84 nymphs or adults hill<sup>-1</sup> in the vegetative stage and 7.76 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif*, 2018–19 and with 12.18 nymphs or adults hill<sup>-1</sup> in the vegetative stage and 24.95 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif*, 2019–20 and with 19.0 nymphs or

adults hill<sup>-1</sup> in the vegetative stage and 52.38 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif*, 2020–21. The higher infestation was observed in farmer practices with 3.33 nymphs or adults hill<sup>-1</sup> in the vegetative stage and 19.84 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif* (June–December), 2018–19 and with 35.40 nymphs or adults hill<sup>-1</sup> in the vegetative stage and 50.30 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif* (June–December), 2019–20 and with 25.42 nymphs or adults hill<sup>-1</sup> in the vegetative stage and 63.47 nymphs or adults hill<sup>-1</sup> in reproductive stage during *kharif* (June – December), 2020–21 (Table 1).

**Table 1: Brown plant hoppers incidence in relation with weather during *kharif*, 2018–19, 2019–20 and 2020–21**

Month	Average Population Hill <sup>-1</sup>						Meteorological Observation								
	2018–19		2019–20		2020–21		Temperature °C (Max.)			Temperature °C (Min.)			Rain fall (mm)		
	Demo	Farmer's Practice	Demo	Farmer's Practice	Demo	Farmer's Practice	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21	2018–19	2019–20	2020–21
Vegetative Period	1.84	3.33	12.18	35.40	19.00	25.42	31.3	30.6	30.3	23.1	23.4	30.6	346.7	379.0	263.8
Reproductive Period	7.76	19.84	24.95	50.30	52.38	63.47	32.0	30.4	30.2	17.8	18.5	18.2	12.12	44.1	35.9

Vegetative period (August–September) and Reproductive Period (October–December)

The total cost of cultivation incurred, gross and net returns & B:C ratio in this study to assess the economic impact of technology of

IPM module and farmer practice. The data in table 2 revealed that the yield of IPM module followed plot was 7130, 5513 and 5065 kg ha<sup>-1</sup>

whereas in farmer practice the yield was 6733, 4851 and 4768 kg ha<sup>-1</sup> during (June–December), 2018–19, 2019–20 and 2020–21 respectively. The economic analysis results revealed that the paddy crop recorded higher gross income from IPM module were ₹10337, ₹101981 and ₹88642 ha<sup>-1</sup> as related to ₹97636, ₹89744 and ₹83440 ha<sup>-1</sup> in farmers practice during *kharif* (June – December), 2018–19, 2019–20 and 2020–21 respectively. The B:C Ratio in IPM module was 1.88, 2.68 and 1.78 while in farmer practice plot was 1.43, 1.94 and 1.27 during *kharif* (June – December), 2018–19, 2019–20 and 2020–21 respectively. The results of the current study are in line with the findings [20,21,22,23]. IPM module shown positive results in respect of yield

and economics of rice. It was marked from the results that B:C Ratio of rice crop in IPM module was higher as compared to farmer practice in both the years. Because of non-adoption of IPM module for brown plant hopper management in rice crop resulted in lower B:C Ratio in farmer practice. Thus, promising B:C Ratio and higher net returns in IPM module showed the economic sustainability of the demonstrated technology and influenced the farmers on the utility of technology provided at actual farming situation. Therefore, it is concluded that the technology demonstrated treatments can be recommended in large scale to manage brown plant hoppers in rice fields in ensuing cropping seasons (Table 2).

**Table 2: Economic impact of experiment during *kharif* (June–December), 2018–19, 2019–20 and 2020–21**

Particulars	Yield (Kg ha <sup>-1</sup> )			Percent increase in yield over check (%)			Cost of cultivation (₹ ha <sup>-1</sup> )			Gross returns (₹ ha <sup>-1</sup> )			B:C Ratio		
	2018-19	2019-20	2020-21	2018-19	2019-20	2020-21	2018-19	2019-20	2020-21	2018-19	2019-20	2020-21	2018-19	2019-20	2020-21
IPM Module	7130	5513	5065	5.88	13.64	6.22	54950	38068	49915	103378	101981	88642	1.88	2.68	1.78
Farmer's Practice	6733	4851	4768	-	-	-	68238	46353	65653	97636	89744	83440	1.43	1.94	1.27

#### 4. Conclusion

In IPM module, documented higher paddy yield with net returns of ₹48428, ₹63913 and ₹38727 ha<sup>-1</sup> which was about 5.88%, 13.64% and 6.22% with corresponding B:C ratios of 1.88, 2.68 and 1.78 higher than the non IPM module with ₹29398, ₹43391 and ₹17787 ha<sup>-1</sup> during *kharif*, 2018–19,

2019–20 and 2020-21 respectively. The IPM based practices were found effective in comparison to farmer practice. From the above study, it can be concluded that by adopting IPM based brown plant hopper management strategies in in can be efficiently managed instead of practicing chemical control measure alone.

#### 5. References-

- Hosseini SZ, Barbaeian JNA, Bagheri NA. Study of silicon effect on plant growth and resistance to stem borer in rice. *Communication in Soil Science and Plant analysis*. 2012; 43 (21): 2744–2751.
- Kulagod SD, Hegde M, Nayak GV, Vastrad AS, Hugar PS, Basavanagoud K. Evaluation of insecticides and bio-rationals against yellow stem borer and leaf folder on rice crop. *Karnataka Journal of Agricultural Sciences*. 2011; 24(2): 244–246.
- Pasalu IC, Katti G. Advances in ecofriendly approaches in rice IPM. *Journal of Rice Research*. 2006; 1(1): 83–90.
- Min S, Lee SW, Choi BR, Lee SH, Kwon DH. Insecticide resistance monitoring and correlation analysis to select appropriate insecticides against *Nilaparvata lugens* (Stal), a migratory pest in Korea. *Journal of Asia-Pacific Entomology*. 2014; 17: 711–716.
- Cheng J. Rice plant hopper problems and relevant causes in China. In: Heong, K.L., Hardy, B. (Eds.). *Plant Hoppers: New Threats to the Sustainability of Intensive Rice Production Systems in Asia*. International Rice Research Institute, Los Baños, Philippines. 2009. 157–177.

6. Gurr GM, Liu J, Read DMY, Catindig JLA, Cheng JA, Lan LP *et al.* Parasitoids of Asian rice plant hopper (Hemiptera: Delphacidae) pests and prospects for enhancing biological control by ecological engineering. *Annals of Applied Biology*. 2011;158: 149–176.
7. Bottrell DG, Schoenly KG. Resurrecting the ghost of green revolutions past: the brown plant hopper as a recurring threat to high-yielding rice production in tropical Asia. *Journal of Asia-Pacific Entomology*. 2012; 15(1): 122–140.
8. Lakshmi VJ, Krishnaiah NV, Katti G, Pasalu IC, Chirutkar PM. Screening of insecticides for toxicity to rice hoppers and their predators. *Oryza*. 2010; 47(4): 295–301.
9. Ratnakar V, Jhansi Lakshmi V, Srinivas C, Jagadeeshwar R, Satendra K, Mangrauthia. Bioefficacy of commonly used insecticides against rice brown planthopper, *Nilaparvata lugens* (Stål). *The Journal of Research PJTSAU*. 2019; 47(4): 28–32.
10. Wang YH, Wu SG, Zhu YC, Chen J, Liu FY, Zhao XP *et al.* Dynamics of imidacloprid resistance and cross-resistance in the brown plant hopper *Nilaparvata lugens*. *Entomologia Experimentalis et Applicata*. 2009; 131: 20–29.
11. Matsumura M, Morimura SS. Recent status of insecticide resistance in Asian rice plant hoppers. *Japan Agriculture Research*. 2010; 44(3): 225–230.
12. Karunaratne SHPP, Weerakoonl KC, Nungaliyadda L, Manuweera GK. Susceptibility of rice insect pests and their natural enemies to commonly used insecticides. *Journal of the National Science Foundation of Sri Lanka*. 2007; 35(2): 97–102.
13. Xue J, Bao YY, Li BL, Cheng YB, Peng ZY, Liu H *et al.* Transcriptome analysis of the brown planthopper *Nilaparvata lugens*. *PLoS ONE*. 2010; 5(12): e14233.
14. Perry T, Batterham P, Daborn PJ. The biology of insecticidal activity and resistance. *Insect Biochemistry and Molecular Biology*. 2011; 41 (7): 411–422.
15. Sharma RK, Bhati DS, Sharma SK. Impact of frontline demonstrations on rapeseed

- mustard growers. *Journal of Progressive Agriculture*. 2017; 8(1): 115-118.
16. Singh DK, Gautam US, Singh RK. Study on yield gap and level of demonstrated crop production technology in sagar district. *Indian Research Journal of Extension Education*. 2007; 7(2&3): 94–95.
17. Shankar M, Aariff Khan MA, BalazziiNaaiik RVT, Sumalini K. Extension interventions for enlightening tribal farmers for enhancing cotton production in Nalgonda district, Telangana. *International Journal of Bio-resource and Stress Management*. 2022; 13 (4): 365 – 371.
18. Singh AK, Pratap SP, Rajpoot SKS. Evaluation of IPM module against major rice insect pests of rice in St Kabir Nagar district of Uttar Pradesh. *International Journal of Current Microbiology and Applied Sciences Special Issue*. 2018; 7: 4400-4404.
19. Trivedi TP, Ahuja DB. Integrated pest management: approaches and implementation. *Indian Journal of Agricultural Sciences*. 2011; 81(11): 981-993.
20. Singh VD, Gangwar RK, Prem G, Choudhary R, Kumar A, Kumar R. Efficacy of granular insecticide against yellow stem borer (*Scirpophaga incertulas*) on basmati rice. *Journal of Krishi Vigyan*. 2017; 5(2): 63-66.
21. Saravanakumar S. Ecological engineering: a potential way for controlling pests and improving paddy productivity. *Indian Journal of Extension Education*. 2020; 56(1): 9-12.
22. Kumar S, Nath S, Kannaujia SK, Gautam AD, Singh BP. Assessment of the integrated pest management against insect pests of paddy in eastern Uttar Pradesh. *Journal of Krishi Vigyan*. 2020; 8(2): 8-11.
23. Balasubramaniam M. Evaluation of frontline demonstrations on integrated pest and disease management (IPDM) practices in paddy. *International Journal of Farm Sciences*. 2022; 12(1): 119-121.

\*\*\*