

Evaluation of maize genotypes against banded leaf and sheath blight incited by *Rhizoctonia solani* f. sp. *sasakii* Exner (Cantharellales: Ceratobasidiaceae) under artificial epiphytotic conditions, India

ABSTRACT

Zea mays L. (Poales) is one of the most important cereal crops belongs to *Poaceae* family with very high potential for yield. Production of maize is hampered by a major constraint and devastating disease banded leaf and sheath blight caused by *Rhizoctonia solani* f. sp. *sasakii*. The experiment was conducted to evaluate the maize genotypes under field conditions at CCS Haryana Agricultural University, Regional Research Station, Karnal (India). A set of fifty seven genotypes (hybrids/inbred lines) were screened against banded leaf and sheath blight during *kharif* 2020 and 2021 under artificial epiphytotic conditions. The severity of banded leaf and sheath blight was recorded following 1-9 scale and genotypes evaluated categorized into four groups *viz.*, resistant, moderately resistant, moderately susceptible and susceptible based on their disease reaction. Out of fifty seven genotypes screened against BLSB, eight inbred lines *viz.*, HKI 161, HKI-163, HKI-193-2, HKI-288-2, HKI-295, HKI-488, HKI-1344 and HKI 1352 (58-9-2)-2 found resistant and seventeen maize inbred lines were moderately resistant, whereas two hybrids (HQPM-4 and HKI-1040-7 x HKI 659-3) showed resistant reaction and ten maize hybrids were moderately resistant. Thus, these resistant/ moderately resistant maize genotypes can be further use in maize breeding program to develop durable BLSB resistant varieties.

Keywords: Banded leaf and sheath blight, *Rhizoctonia solani* f. sp. *sasakii*, inbred lines, hybrids

1. INTRODUCTION

Maize (*Zea mays* L.) is a member of *Poaceae* family, originated in the Mexico, Central America and the most important cereal crops of the tropical and sub-tropical regions of the world agricultural ecosystem. Maize is known as ‘Queen of the Cereals’ or ‘Miracle C 4 crop’ because of its genetic makeup conserve highest yield potentiality comparable to the other cereals and gaining popularity as evidenced by its large scale consumption as food, feed, fodder for animals and as source of industrial raw materials *viz.*, starch, syrup, dextrose, gelatin, protein, alcoholic beverages, oil, pharmacy, cosmetics, bio-fuel (ethanol), food sweeteners etc. Globally maize area of about 193.7 million hectares is being produced together by over 170 countries with annual production of approximately 1147.7 million MT and average productivity is about 5.75 ton/ha [5]. It occupies an area about 9.89 million hectares having production of 31.65 million tones with average productivity of 3.20 ton/ha in India during 2020-21 [4]. In Haryana, *kharif* maize is cultivated over an area of 6.63 thousand hectares with average production of 20.06 thousand tones achieving productivity of 3.02 ton/ha [4].

Despite its high yield potential, maize is affected by several biotic and abiotic stresses that affect the

tropical and subtropical maize yield production. Globally, about one hundred and twelve diseases of maize has been reported from different locations, of these sixty five diseases to occur in India incited by pathogens belonging to fungi, bacteria, viruses and nematodes groups. Out of these, banded leaf and sheath blight disease of maize caused by fungus, *Rhizoctonia solani* f. sp. *sasakii* Exner, (Tel: *Thanatephorus sasakii* (Shirai) Tu and Kimbro) belonging to anastomosis group (AG1-IA) is considered as the major constraint responsible for yield losses. The disease causes severe huge impact on yield in several Asian countries [17]. In India, the disease was first reported from Tarai region of Uttar Pradesh and was recognized as a minor disease [15], later on disease was appeared in the epidemic form in humid and warm foot hill regions of Mandi district in Himachal Pradesh in early 1970 [22].

The banded leaf and sheath blight is known as highly destructive disease in Himachal Pradesh, Assam, Meghalaya, Uttar Pradesh, Bihar, Nagaland, Jammu Kashmir, Haryana, Uttarakhand, Punjab, Sikkam, Madhya Pradesh, Delhi, Rajasthan, Orissa, Andhra Pradesh and West Bengal (India) [2], [16], [19]. The magnitude of grain yield losses may vary between 11-40 per cent with 71 percent disease index in India, up to 100 per cent at the ear rot phase in Indonesia and 44-66 per cent in Myanmar depending upon prevailing agro climatic conditions [13], [19], [20]. In India, losses in terms of grain yield may estimate to the range of 11-40 per cent, even to 100 per cent especially in Haryana due to continue rains in the months of July and August [12]. In this study, we investigated the reaction of maize genotypes against banded leaf and sheath blight disease incited by soil-borne fungal pathogen *R. solani* f. sp. *sasakii*.

2. MATERIALS AND METHODS

2.1 Experimental setup and field preparation

In the present investigation, fifty seven maize genotypes (Hybrids/inbred lines) were screened against banded leaf and sheath blight (BLSB) at research area of Chaudhary Charan Singh Haryana Agricultural University, Regional Research Station, Karnal during *kharif* seasons 2020 and 2021 under artificial epiphytotic conditions. These conditions were created through inoculations of fungal coated barley grains inserted between second and third sheath of maize during evening hours. Two lines of each genotype consist of 3 m length with a row to row and plant to plant spacing of 75 cm and 20 cm, respectively were sown during both the *kharif* seasons. Experiments were laid out in randomized complete block design with three replications and all the recommended packages and practices of CCS Haryana Agricultural University, Hisar were followed [3].

2.2 Preparation of mass culture of *R. solani* f. sp. *sasakii*

Barley grains were soaked in water for 24 hours and dispensed 40 g seeds in 250 ml Erlenmeyer flask after removing excess of water and autoclaved at 121.6°C at 15 lbs for 20 minutes. Each flask was inoculated with 5 mm mycelium disc of *R. solani* isolates, derived from the actively growing cultures on

the potato dextrose agar (PDA) plates and incubated at 25 ±2°C in biochemical oxygen demand (BOD) incubator for ten days. The fully grown fungal mycelium on barley grains were dispensed out from flask on a paper for drying at 15°C. The impregnated grains were utilized for inoculation and stored in paper bags or in the flasks for later use [1]. The barley grains are widely use for preparation of mass culture of *R. solani*, provides appropriate nutrients to pathogen for increasing inoculums density.

2.3 Artificial inoculation method

Inoculation on 30-35 days old maize plants was performed during evening hours and the moist conditions prevailed either due to rain or through irrigation. The grain inoculums were inserted between the leaf sheath and stalk at lower second or third internodes level of plant [1].

2.4 Observations recorded

The observations were recorded using 1-9 scale (Table 1) [7] and categorized them into different disease reaction on the basis of average disease score as follows:

Table 1: Banded leaf and sheath blight disease rating scale in maize (India)

Rating scale	Disease reaction	PDI (%)
1.0-3.0	Resistant (R)	0.00-33.33
4.0-5.0	Moderately resistant (MR)	44.44-55.55
6.0-7.0	Moderately susceptible (MS)	66.66-77.77
8.0-9.0	Susceptible (S)	88.88-99.99

PDI- Per cent disease intensity, Source- Hooda et al. (7)

3. RESULTS AND DISCUSSION

3.1 Screening of maize genotypes (hybrids/inbred lines) against banded leaf and sheath blight

Host plant resistance is considered as one of the outstanding ways for managing the diseases to a sustainable agriculture with cost effective and eco-friendly approach. All the genotypes (hybrids/inbred lines) were showed greater extent of variations in disease reaction during both the *kharif* seasons 2020 and 2021. Data indicated that all the genotypes showed resistant to susceptible reaction against banded leaf and sheath blight of maize. During *kharif* 2020 and 2021 the mean of disease score of banded leaf and sheath blight of maize varied from 2.2 to 8.4 (Table 2).

A total of forty four state inbred lines of maize were screened against banded leaf and sheath blight (BLSB). Out of which eight maize inbred lines namely HKI 161, HKI-163, HKI-193-2, HKI-288-2, HKI-295, HKI-488, HKI-1344 and HKI 1352 (58-9-2)-2 were found resistant (disease score ≤ 3.0) to banded leaf and sheath blight disease and seventeen maize inbred lines *viz.*, MBR-139, HKI-164-7-6, HKI-191-2-5, HKI-193-1, HKI-323, HKI-335, HKI-488-1RG, HKI-536C, HKI-1105, HKI-1128, HKI-1332, HKI-1345, HKI-1354-7, HKIL-287, HKI-1042-NP 19-ER-1 and PC-1 ER-4 were moderately resistant (disease score 3.1-5.0). Eighteen maize inbred lines (HKI-488T-3, HKI-536YN, BC- 570 ER-1, HKI-1011, HKI-1015-6, HKI-1025, HKI-1040-4, HKI-1105-ER-5, HKI-1348-8-2, HKI-1352, HKI-1651 (14-3-1), HKI-1657, HKI-1659, HKI-1664, NP-80ER-2, PC-3 ER-1, PC4-6 and PC 1221-4) were

moderately susceptible (disease score 5.1-7.0) whereas, one inbred line PC 1473-5 exhibited susceptible reaction (disease score ≥ 7.0) against BLSB (Table 3).

Similarly, thirteen maize hybrids were screened against banded leaf and sheath blight (BLSB) during *kharif* 2020 and 2021. Among them, two hybrids *viz.*, HQPM-4 and HKI-1040-7 x HKI 659-3 were found resistant (disease score ≤ 3.0) to banded leaf and sheath blight disease and ten maize hybrids *viz.*, HQPM-1, HM-8, HM-11, HKI288-2x HKI 323, HKI 193-2x L-287, HKI 327 Tx L-287, HKI-1105-6x LM-17, HKI-1354-7x HKI-1344, HKI-1105 x IMR 307 and HKI-184 x HKI-1105 were moderately resistant (disease score 3.1-5.0). One maize hybrid (HKI 288-2x LM-17) showed moderately susceptible reaction against BLSB whereas, none of maize hybrid exhibited susceptible reaction (Table 4). The variable response of the genotypes for *R. solani* f. sp. *sasakii* has also been observed by earlier researches. However, there have been limited reports on genetics of resistance to banded leaf and sheath blight in maize [18]. The developments of resistant hybrids through classical plant breeding have been slow because unavailability of banded leaf and sheath blight resistance source [14, [6]. Lal [8] screened one hundred inbred lines and sixty hybrids of maize against banded leaf and sheath blight during *kharif* 2011, out of which eighteen inbred and fourteen hybrids were resistant. Malik et al. [10] also evaluated thirty inbred lines and seven hybrids during *kharif* 2013-14. Out of thirty, only four inbred lines (HKI-488WG, HKI-C-141, HKI-1040-5 and HKI-1347-4LT) were resistant, whereas (HKI-139, HKI-1352, HKI- 1378, HKI-150, HKI-766RG, HKI-MBR-139, HKI-323 and HKI-325-17A) were found moderately resistant. In case of hybrids, HM 11 was resistant whereas HQPM 1, HQPM 7, HM 6 and HKI 1040-7 x HKI 1128 were found moderately resistant. Similarly, Madhavi et al. [9] also screened twenty two maize genotypes out of which one inbred line PFSR9-2 was resistant against BLSB, whereas PFSR6-1, PFSR6-2 and PFSR18 were moderately resistant. In case of hybrids, one hybrid DHM 117 showed resistant reaction while DKC9145, DKC9133 and KMH3110 were found moderately resistant. Thakur et al. [22] and Meena et al. [11] also corresponds the variable pathological response in maize inbred lines and hybrid against BLSB pathogen under artificial epiphytotic conditions.

Table 2: Disease rating and disease reaction of maize genotypes against banded leaf and sheath blight under field conditions during *kharif* 2020 and 2021

Sr. No	Inbred lines	Disease score		Avg. rating score	Disease reaction	Sr. No	Inbred lines	Disease score		Avg. rating score	Disease reaction
		2020	2021					2020	2021		
1	HKI-161	2.8	2.6	2.7	R	31	HKI-536C	3.6	3.8	3.7	MR
2	HKI-163	2.5	2.4	2.5	R	32	HKI-536YN	5.6	6.2	5.9	MS
3	HKI-193-1	4.6	4.9	4.8	MR	33	HKI-1015-6	6.4	6.7	6.6	MS
4	HKI-193-2	2.6	2.5	2.6	R	34	HKI-1040-4	5.6	5.9	5.8	MS
5	HKI-488-1RG	3.9	4.6	4.3	MR	35	HKI-1105-ER-5	6.8	6.5	6.7	MS
6	HKI-1025	5.4	6.5	6.0	MS	36	HKI-1332	4.2	4.7	4.5	MR
7	HKI-1128	4.8	4.6	4.7	MR	37	HKI-1348-8-2	5.3	6.4	5.9	MS
8	HKI-1344	2.4	2.7	2.6	R	38	HKI 1352 (58-9-2)-2	2.8	2.6	2.7	R
9	HKI-191-2-5	4.2	4.4	4.3	MR	39	HKI-1659	6.4	6.9	6.7	MS
10	HKI-288-2	2.3	3.6	3.0	R	40	PC-1 ER-4	3.8	4.6	4.2	MR
11	HKI-323	3.5	4.5	4.0	MR	41	PC-3 ER-1	5.4	6.8	6.1	MS
12	HKI-335	4.6	4.8	4.7	MR	42	PC 1221-4	6.5	5.8	6.2	MS
13	HKI-488	2.7	2.4	2.6	R	43	HKI-1011	6.8	6.4	6.6	MS
14	BC-570ER-1	6.3	6.9	6.6	MS	44	PC 4-6	5.9	6.4	6.2	MS
15	HKI-1105	3.5	3.8	3.7	MR						
						Sr. No	Hybrids	Disease score		Avg. rating score	Disease reaction
								2020	2021		
16	HKI-1345	4.1	4.5	4.3	MR	1	HQPM-1	3.6	3.9	3.8	MR
17	HKI-1352	6.4	6.8	6.6	MS	2	HQPM-4	2.4	2.2	2.3	R
18	HKI-1354-2	4.8	4.6	4.7	MR	3	HM-8	3.5	3.8	3.7	MR
19	HKI-1354-7	3.5	4.2	3.9	MR	4	HM-11	4.1	4.5	4.3	MR
20	HKI-1651 (14-3-1)	5.9	6.8	6.4	MS	5	HKI 288-2x HKI 323	4.8	4.6	4.7	MR
21	HKI-1657	6.4	6.6	6.5	MS	6	HKI 193-2x L-287	4.5	4.8	4.7	MR
22	HKI-1664	6.6	5.9	6.3	MS	7	HKI 288-2x LM-17	5.6	6.2	5.9	MS
23	HKIL-287	4.2	4.6	4.4	MR	8	HKI 327 T x L-287	3.7	4.2	4.0	MR
24	HKI-1042-NP 19-ER-1	3.8	3.4	3.6	MR	9	HKI-1105-6 x LM-17	4.5	5.3	4.9	MR
25	NP-80ER-2	6.7	6.4	6.6	MS	10	HKI-1040-7 x HKI 659-3	2.4	2.7	2.6	R
26	PC 1473-5	7.6	8.4	8.0	S	11	HKI-1354-7 x HKI-1344	4.3	4.5	4.4	MR
27	MBR-139	4.3	4.8	4.6	MR	12	HKI-1105 x IMR 307	3.8	4.1	4.0	MR
28	HKI-164-7-6	3.9	4.5	4.2	MR	13	HKI-184 x HKI-1105	4.1	4.6	4.4	MR
29	HKI-295	2.4	2.6	2.5	R						
30	HKI-488T-3	6.8	6.4	6.6	MS						

Table 3: Screening of inbred lines against banded leaf and sheath blight (*R. solani* f. sp. *sasakii*) under field conditions during *kharif* 2020 and 2021

Disease score	Disease reaction	No. of inbreds	Inbreds
< 3.0	Resistant	8	HKI 161, HKI-163, HKI-193-2, HKI-288-2, HKI-295, HKI-488, HKI-1344, HKI 1352 (58-9-2)-2
3.1-5.0	Moderately Resistant	17	MBR-139, HKI-164-7-6, HKI-191-2-5, HKI-193-1, HKI-323, HKI-335, HKI-488-1RG, HKI-536C, HKI-1105, HKI-1128, HKI-1332, HKI-1345, HKI-1354-7, HKIL-287, HKI-1042-NP 19-ER-1, PC-1 ER-4
5.1-7.0	Moderately Susceptible	18	HKI-488T-3, HKI-536YN, BC- 570 ER-1, HKI-1011, HKI-1015-6, HKI-1025, HKI-1040-4, HKI-1105-ER-5, HKI-1348-8-2, HKI-1352, HKI-1651 (14-3-1), HKI-1657, HKI-1659, HKI-1664, NP-80ER-2, PC-3 ER-1, PC4-6, PC 1221-4
7.1-9.0	Susceptible	1	PC 1473-5

Table 4: Screening of hybrids against banded leaf and sheath blight (*R. solani* f. sp. *sasakii*) under field conditions during *kharif* 2021 and 2021

Disease score	Disease reaction	No. of hybrids	Hybrids
< 3.0	Resistant	2	HQPM-4, HKI-1040-7 x HKI 659-3
3.1-5.0	Moderately Resistant	10	HQPM-1, HM-8, HM-11, HKI288-2x HKI 323, HKI 193-2x L-287, HKI 327 Tx L-287, HKI-1105-6x LM-17, HKI-1354-7x HKI-1344, HKI-1105 x IMR 307, HKI-184 x HKI-1105
5.1-7.0	Moderately Susceptible	1	HKI 288-2x LM-17
7.1-9.0	Susceptible	0	NIL

4. CONCLUSION

This study emphasized the identification of resistance source among maize genotypes (hybrids /inbred lines) against banded leaf and sheath blight disease. This study revealed eight inbred lines *viz.*, HKI 161, HKI-163, HKI-193-2, HKI-288-2, HKI-295, HKI-488, HKI-1344 and HKI 1352 (58-9-2)-2 and two hybrids (HQPM-4 and HKI-1040-7 x HKI 659-3) of maize found resistant against banded leaf and sheath blight can be further utilize for developing resistant maize varieties and help to develop management strategies for the control of banded leaf and sheath blight. These resistant /moderately resistant genotypes of maize could be used for identifying durable resistance, quantitative trait loci analysis/as donor parents in breeding programs.

ACKNOWLEDGEMENT

All the sort of assistance rendered by CCS Haryana Agricultural University, Regional Research Station, Karnal for the above study is gratefully acknowledge.

REFERENCES

1. Ahuja SC, Payak MM. A field inoculation technique for evaluating maize germplasm to banded leaf and sheath blight. *Indian Phytopathol.* 1978;31(5):17-20.
2. Ahuja SC, Payak MM. Banded leaf and Sheath blight of maize. In: Agnihotri VP; Sarbhoy AK, Kumar D (eds.) *Perspectives in Mycology and Plant Pathology*, Malhotra Publishing House, New Delhi, India. 1988;1-186.
3. Anonymous. Package and practices of *kharif* crops. Directorate of extension education. CCS HAU, Hisar. 2021;40-49.
4. Anonymous. <https://www.indiastat.com/data/agriculture/maize-17199>. 2022.
5. FAO. <https://www.fao.org/faostat/en/#data/QI>. 2020.
6. Han YP, Xing YZ, Cheng ZX, Gu SL, Pan XB, Chen XL, Zhang QF. Mapping QTL for horizontal resistance to sheath blight in an elite rice restorer line, Minghui. *Acta Gen Sinca.* 2002;29(7): 622-626.
7. Hooda KS, Bagaria PK, Khokhar M, Kaur H, Rakshit S. Mass screening techniques for resistance to maize diseases. ICAR Indian Institute of Maize Research, PAU Campus, Ludhiana. 2018;93.
8. Lal M. Studies on banded leaf and sheath blight of maize caused by *Rhizoctonia solani* f. sp. *sasakii* Exner. M.Sc. thesis, CCSHAU, Hisar. 2013;18-46.
9. Madhavi M, Narayan PR, Ranga R, Sudarshan MR. Evaluation of maize genotypes against banded leaf and sheath blight disease incited by *Rhizoctonia solani* f. sp. *sasakii* (Kuhn) Exner. *J Res. ANGRAU.* 2012;40: 20-23.
10. Malik VK. Performance of maize germplasm against BLSB disease incited by *Rhizoctonia solani* f. sp. *sasakii*. *Indian Phytopathol.* 2016;69(1): 97-98.
11. Meena BR, Yerasu S.R, Gupta N, Singh J. Resistance assessment and biochemical responses of maize genotypes against *Rhizoctonia solani* f. sp. *sasakii* Exner causing banded leaf and sheath blight disease. *Australasian Plant Pathol.* 2020;50: 41-49.
12. Mehra R, Kamboj MC, Mehla JC, Lal M, Chand M. Status of maize disease and their management in Haryana. In *Proceeding of National Seminar on Sustainable and Food Security: Challenges in Changing Climate*, CCSHAU, Hisar. 2012;217.
13. Muis A. Evaluation of *Bacillus subtilis* BR23 applied as spray suspension against banded leaf and sheath blight (*Rhizoctonia solani* Khun) in corn. *Seminor Ilmiah dan Pertemuan Tahunan PEI dan PFI XVIII Komda Sul-Sel, Indonesia.* 2007;28-34.
14. Pan YB, Rush MC. Studies in the US on genetics and breeding for resistance to rice sheath blight. *J Jiangsu Agriculture Coll.* 1997;18(10): 57-63.
15. Payak MM, Renfro BL. Diseases of maize new to India. *Indian Phytopathological Society Bulletin* 3, IARI, New Delhi. 1966;14-18.
16. Rani DV, Reddy NP, Devi UG. Banded leaf and sheath blight of maize incited by *Rhizoctonia solani* f. sp. *sasakii* and its management. *Int J Pharma and Biol Arch,* 2013;4(4): 52-60.
17. Sharma G, Saxena SC. Integrated management of banded leaf and sheath blight of maize (*Zea mays* L.) caused by *Rhizoctonia solani* (Kuhn). *Adv Plant Sci.* 2002;15(1): 107-113.

18. Sharma RC, Vasal SK, Gonzalez F, Batsa BK, Singh, NN. Redressal of banded leaf and sheath blight of maize through breeding, chemical and biocontrol agents. In: Proceeding of the 8th Asian Regional Maize Workshop: New Technologies for the New Millennium, Bangkok. 2002;391-397.
19. Singh BM, Sharma YR. Evaluation of maize germplasm to banded sclerotial disease and assessment of yield loss. *Indian Phytopathol.* 1976;29: 129-132.
20. Sudjono M.S. Effectiveness of antagonists against sheath blight and ear rot caused by *Rhizoctonia solani* Kuhn. Proceeding Kongres National XII dan Seminar IImiah PFI. Jogjakarta, 1995;545-549.
21. Thakur N, Lata S, Sharma BK, Devlash R. Evaluations of maize genotypes against banded leaf and sheath blight under natural and artificial epiphytotic conditions. *Himachal J Agri Res.* 2018;44: 17-24.
22. Thakur SM, Sharma SL, Munjal RL. Correlation studies between incidences of banded sclerotial disease maize. *Indian J Mycol Plant Pathol.* 1973;3: 180-181.