

Original Research Article

Effect of tillage methods, farmyard manure and potassium rates on some soil chemical properties and nutrient contents in cassava in Kagera, Tanzania

ABSTRACT

A study was conducted in Bukoba, Missenyi and Biharamulo districts, Tanzania to determine the effect of tillage methods, farmyard manure (FYM) and potassium rates on soil pH and the concentrations of N, P and K in the soils and cassava leaves. The treatments were arranged in the Randomized Complete Block Design (RCBD) using the split-plot design with three replications. Tillage methods (flat tillage, open ridging and tied ridging), were the main plots, and the fertilizer rates [farmyard manure (FYM) alone at 4 MT ha⁻¹ alone or FYM alone at 8 MT ha⁻¹, nitrogen (40 kg N ha⁻¹) + phosphorus (30 kg P ha⁻¹) + potassium at 40, 80 or 120 kg K ha⁻¹ and the combination of FYM alone at 4 MT ha⁻¹ or FYM alone 8 MT ha⁻¹ + potassium at 40, 80 or 120 kg K ha⁻¹] and the control, were the sub-plots. The inherent and post-harvest composite soil samples for determining soil pH and the concentration of N, P and K in each site were collected, processed and analysed. Post-harvest soil samples and cassava leaf samples were collected from the control plots and plots that received the combination of FYM at 8 MT ha⁻¹ and potassium at 40 or 120 kg K ha⁻¹ during the second cropping season. The leaf samples were oven dried at 70 °C, grounded to pass through 0.5 mm sieve and analyzed. The results indicate that there was no a significant (P = .46) difference in the soil pH and the concentrations of N, P and K in the soils and cassava leaves among the tillage methods. There was a significant (P < .001) difference in the soil pH and the concentrations of N, P and K in the soils and cassava leaves between the control and the combined use of FYM and potassium rates. However, there was no a significant (P = .08) difference in the concentrations of N, P and K in the soils among the combined use of FYM and potassium rates, but there was a significant (P < .001) difference in the concentrations of N and K in cassava leaves among the combined use of FYM and potassium rates. The combined use of FYM at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ or FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹, increased the soil pH and the concentration of N and K in the soil. In addition, the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ gave significantly (P < .001) higher concentrations of N and K in cassava leaves than the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹. Therefore, combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ is desirable for increasing the concentration of N and K in the soil and in cassava leaves. However, for the resource-poor farmers who cannot afford the high rate of K, the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ could be used.

Keywords: Farmyard manure, flat tillage, open ridging, tied ridging, potassium fertilizer, soil pH, nutrients concentration, cassava leaves.

1. INTRODUCTION

The combined use of organic and inorganic fertilizers plays an important role in the improvement of soil physico-chemical properties [1] such as water holding capacity, soil structure, cation exchange capacity (CEC), and lowering bulk density [2], which results in the availability of macro and micro nutrients to plants [3]. The soil properties such as pH, organic

Comment [O1]: It would have been better if, in addition to the things examined in the soil and plant leaves, the amount of plant growth and plant yield were also measured
If you have the morphological data of the plant, add it because assuming this research is done, what effect does it have on the growth and performance of the plant?
This is a question that you have to come to the conclusion of and therefore add the morphological and functional analysis of the plant.

Comment [f2]: unclear the right form is $p \leq 0.05$

Comment [f3]: In statistics, this expression is not common, more than the probability level of one and five percent

Comment [f4]: this statment must be rewrite again and edited in scientific pattern

matter (OM) contents and available forms of macro-nutrients like nitrogen (N) phosphorus (P) and potassium (K) are significantly affected by long-term use of mineral fertilizers and organic manures [4]. For example, [5] reported significantly increase in N, P and K concentrations in the soil due to the combined use of organic and inorganic fertilizers. Incorporation of manures in the soil has beneficial effects on soil health by improving physico-chemical properties and supplying nutrients like nitrogen (N), phosphorus (P), and potassium (K) [6]. However, neither use of organic manure nor chemical fertilizers alone can achieve sustainable crop yields under different cropping system where the nutrient depletion and turnover in soil-plant system is high. Although the use of chemical fertilizers improves crop productivity, continuous use of chemical fertilizers alone is associated with decline in some soil properties and crop yields over time [7] due to high solubilization, fast release of nutrients in the soil, and nutrients loss through leaching. For this reason, therefore, an integrated use of inorganic fertilizers and organic manures is a sustainable approach for efficient nutrient usage, which enhances efficiency of the chemical fertilizers while reducing nutrient losses [8].

Nutrient uptake by the crops depends on both the ability of the roots to absorb nutrients and nutrients concentration at the surface of the roots. As the plant grow, roots spread out laterally and vertically by taking advantage of areas within the soil that have more water and nutrients depending on the stage of plant growth [9]. Combined use of organic and inorganic fertilizers increases nutrient uptake by the crops. For example, [10] reported increased N uptake by maize crop when compost was mixed with mineral N fertilizer as compared to compost or N mineral fertilizer alone due to improving soil physico-chemical properties and mineralization of the compost. However, in dry soil, nutrient uptake by the plants become low and nutrient levels in plant tissues may be lower than normal [11].

Tillage practices influence soil temperature, moisture and aeration, which eventually affect nutrient uptake by plants [12]. Fertilizer placement also influences nutrient availability and depending upon soil and climate conditions, either enhances or reduces nutrient uptake [9]. Therefore, this study aimed at determining the effects of tillage methods, farmyard and potassium rates, on soil pH and concentrations of N, P and K in the soil and cassava leaves following two consecutive years of planting cassava on tied ridges, open ridges and flat tillage, and the use of different rates of farmyard manure and potassium fertilizers in Bukoba, Missenyi and Biharamulo districts in the Kagera region, Tanzania.

2. MATERIAL AND METHODS

2.1 Description of the study area

2.1.1 Location of the study area

Kagera Region is located in the north-western corner of Tanzania on the western shore of Lake Victoria between latitudes 1°00' and 3°45' south of Equator and between longitudes 30°25' and 32°40' east of Greenwich. It is the fifteenth largest region in Tanzania with an area of about 3 568 600 ha of land, which accounts for approximately 3.3% of Tanzania's total land area. Out of the region's area, 10 173 ha are covered by water of the Lake Victoria, Ikimba and Burigi, and of the river Kagera and Ngono. Administratively, the region has seven districts, namely Biharamulo, Bukoba, Karagwe, Kyerwa, Missenyi, Muleba and Ngara, and borders four countries, namely Uganda, Rwanda, Burundi, and Kenya across Lake Victoria [13, 14]. However, this study was conducted in three districts, namely Bukoba, Missenyi and Biharamulo. The selection of these districts were based on the representative of agro-ecological zones of Kagera region and the potential for cassava production. The representative study sites were Tanzania Agricultural Research Institute (TARI), Maruku Centre in Butairuka village (Bukoba district), Mabuye Primary School in Mabuye village

(Missenyi district) and Rukaragata Farmers' Extension Centre in Rukaragata village (Biharamulo district).

Bukoba district covers an area of 284,100 ha and is situated between latitudes 1° 00' and 3° 00' S and between longitudes 30° 45' and 31° 00' E with altitude between 1200 - 1400 meters above sea level. Missenyi district covers an area of 270 875 ha and is situated between latitudes 1° 00' and 1° 30' S and between longitudes 30° 48' and 31° 49' E with altitude between 1100 - 1400 meters above sea level. Biharamulo district covers an area of 374 400 ha and is situated between latitudes 2° 15' and 3° 15' S and between longitudes 31° 00' and 32° 00' E with altitude ranging from 1100 - 1700 meters above sea level (masl) [13, 14]. Based on rainfall, three agro-ecological zones namely high, medium and low rainfall zones are found in Kagera region [15, 16, 14], which in this study are represented by Bukoba district (high rainfall), Missenyi district (medium rainfall) and Biharamulo district (low rainfall).

Comment [O5]: correct this word

2.1.2 Climate and soils of the study area

The districts in Kagera region experience bimodal rainfall distribution between September and December (short rains) and between March and June (long rains). The mean annual rainfall ranges from 900 - 2400 mm in Bukoba district, 600 - 2000 mm in Missenyi district and 700 - 1000 mm in Biharamulo district [15, 17]. The mean annual temperature ranges from 16 - 28 °C, Missenyi having higher annual temperature (28 °C) than Bukoba and Biharamulo (26 °C). In terms of soil texture, the soils range from sandy clay loam to sandy clay and clay [15, 18]. However, the soils of the study area indicate that P, K and Mg deficiencies were widely spread in Bukoba district while N and S deficiencies were widely spread in Missenyi district and N, P and K deficiencies were widely spread in Biharamulo district [18].

Comment [O6]: these information must be gives in only table

2.2 Site selection

This study was conducted in Bukoba, Missenyi and Biharamulo districts. In each district, one ward and one village in each ward were selected. In each selected village, one site was selected for the establishment of the experimental trial. The selected experimental sites were Tanzania Agricultural Research Institute (TARI)-Maruku Centre, Mabuye Primary School, and Rukaragata Extension Centre in Bukoba, Missenyi and Biharamulo districts, respectively. The locations of the experimental trial sites are presented in Figure 1.

2.3 Experimental layout and treatments application

Three field experimental trials, one in each study site were established in two consecutive seasons (2018/19 and 2019/20) in Bukoba, Missenyi and Biharamulo districts. In each trial site, land was prepared by clearing of bushes and trees before trial establishment, followed by ploughing and harrowing. Ridges were prepared by heaping up the soil to about 60 cm within 1 m wide (0.5 m from each side of the ridge top) using hand hoe. Plot size was 6 m x 5 m and the separations between plots and blocks were 1.5 m and 2 m apart, respectively. The treatments were arranged in Randomized Complete Block Design (RCBD) with three replications using the split-plot Design; with tillage methods (flat tillage, open ridging and tied ridging) as the main plots and fertilizer rates [FYM at 4 MT ha⁻¹, 8 MT ha⁻¹, (N + P at 40 kg N ha⁻¹ and 30 kg P ha⁻¹) + potassium at 40, 80 or 120 kg K ha⁻¹, combination of FYM at 4 MT ha⁻¹ or 8 MT ha⁻¹ + potassium at 40, 80 or 120 kg K ha⁻¹] and the control as the subplots (Table 1). The combinations of N at 40 kg N ha⁻¹ + P at 30 kg P ha⁻¹ [19] + potassium at 40, 80, or 120 kg K ha⁻¹; were applied as inorganic fertilizer treatments and FYM at 4 MT ha⁻¹ or 8 MT ha⁻¹ + potassium at 40, 80 or 120 K kg ha⁻¹ were applied as the combinations of organic and inorganic fertilizer treatments.

Comment [O7]: arrangement is correct edited

Comment [O8]: it is better said kg per hectar

Comment [O9]: The treatment should be clearly defined
There is no word or




Figure 1. Location of experimental trial sites in Bukoba, Missenyi and Biharamulo districts

Source: [14]

Farmyard manure was applied at planting along the planting rows in the flat tillage treatment and along the ridges in the open and tie-ridging treatments followed by incorporation into the soils. The applied farmyard manure in each experimental site was collected from one farmer. In all districts, the distance from the source of manure to the experimental sites ranged from 20–30 km. Inorganic fertilizers, namely di-ammonium phosphate (DAP) for N and P and muriate of potash (MOP) for K were applied in two splits; the first split at one month after planting for allowing fibrous roots development on cassava cuttings for nutrients uptake. The

second split of inorganic fertilizer was applied at three months after planting by banding the fertilizers round each cassava plant. Improved cassava variety (*Mkumba*), was the test variety, which was planted at a spacing of 1 m x 1 m in each treatment. The experimental plots were maintained free from weeds, throughout the growing period, and repeated in the following season while maintaining the same plots [20].

Table 1. Experimental treatments in the split-plot design

Flat tillage	Main plots	
	Open ridge tillage	Tied ridge tillage
	Sub plots	
Co	Co	Co
FYM ₄	FYM ₄	FYM ₄
FYM ₈	FYM ₈	FYM ₈
K ₄₀ N ₄₀ P ₃₀	K ₄₀ N ₄₀ P ₃₀	K ₄₀ N ₄₀ P ₃₀
K ₈₀ N ₄₀ P ₃₀	K ₈₀ N ₄₀ P ₃₀	K ₈₀ N ₄₀ P ₃₀
K ₁₂₀ N ₄₀ P ₃₀	K ₁₂₀ N ₄₀ P ₃₀	K ₁₂₀ N ₄₀ P ₃₀
FYM ₄ K ₄₀	FYM ₄ K ₄₀	FYM ₄ K ₄₀
FYM ₄ K ₈₀	FYM ₄ K ₈₀	FYM ₄ K ₈₀
FYM ₄ K ₁₂₀	FYM ₄ K ₁₂₀	FYM ₄ K ₁₂₀
FYM ₈ K ₄₀	FYM ₈ K ₄₀	FYM ₈ K ₄₀
FYM ₈ K ₈₀	FYM ₈ K ₈₀	FYM ₈ K ₈₀
FYM ₈ K ₁₂₀	FYM ₈ K ₁₂₀	FYM ₈ K ₁₂₀

CO = control (no fertilizer application); FYM₄ = farmyard manure at 4 MT ha⁻¹; FYM₈ = farmyard manure at 8 MT ha⁻¹; K₄₀N₄₀P₃₀ = potassium at 40 kg K ha⁻¹, nitrogen at 40 kg N ha⁻¹ and phosphorus 30 kg P ha⁻¹; K₈₀N₄₀P₃₀ = potassium at 80 kg K ha⁻¹, nitrogen at 40 kg N ha⁻¹ and phosphorus 30 kg P ha⁻¹; K₁₂₀N₄₀P₃₀ = potassium at 120 kg K ha⁻¹, nitrogen at 40 kg N ha⁻¹ and phosphorus 30 kg P ha⁻¹; FYM₄K₄₀ = farmyard manure at 4 MT ha⁻¹ and potassium at 40 kg K ha⁻¹; FYM₄K₈₀ = farmyard manure at 4 MT ha⁻¹ and potassium at 80 kg K ha⁻¹; FYM₄K₁₂₀ = farmyard manure at 4 MT ha⁻¹ and potassium at 120 kg K ha⁻¹; FYM₈K₄₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹; FYM₈K₈₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 80 kg K ha⁻¹; FYM₈K₁₂₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹.

2.4 Data collection

The soil samples for determining soil pH and the concentrations of N, P and K, in the inherent soils were collected before trial establishment during the first season. The post-harvest soil samples for determining the effect of applied treatments on the soil pH and the concentrations of N, P and K in the soils after two-consecutive years of treatments application were collected during the second season. The soil samples were randomly collected at 0-20 cm depth using a zig-zag pattern over the whole field area using a soil auger from at least 20 spots, mixed thoroughly to get one composite soil from each experimental site, air-dried, grounded, sieved through 2 mm sieve, packed, and labelled for laboratory analysis. The soil pH and the concentrations of N, P and K for the inherent soils in each site are presented in Table 2. In addition, the post-harvest composite soil samples from the control plots and the plots that received the combination of FYM at 8 MT ha⁻¹ and potassium at 40 and 120 kg K ha⁻¹ in each site were collected from each replication using the soil auger by taking the soil at 0-20 cm depth from the planting rows or ridges within the plots of each tillage method, mixed thoroughly to get one composite soil sample, air-dried, grounded, sieved through 2 mm sieve, packed, and labelled for laboratory analysis for soil pH, N, P and K.

Cassava leaf samples were collected at 4 months after planting during the second cropping season to determine the contents of N, P and K in the cassava leaves. At least 20 standard

leaves (the fourth young fully-expanded leaves from the top-blades) were collected from the control, the combined use of high rate of FYM (8 MT ha⁻¹) and low rate of K (40 kg K ha⁻¹), and the combined use of high rate of FYM (8 MT ha⁻¹) and high rate of K (120 kg K ha⁻¹) from each replication in each site. The collected leaf samples were oven dried at 70 °C, grounded so fine that they can pass through 0.5 mm sieve using TecatorCyclotec 1093 Sample Mill, packed in paper bags, and well labelled for N, P and K analysis.

It should be noted that the decision to collect and analyze the leaf samples and post-harvest soil samples from the control, the combination of FYM at 8 MT ha⁻¹ + K at 40 kg K ha⁻¹ and the combination FYM at 8 MT ha⁻¹ + K at 120 kg K ha⁻¹, for soil pH, N, P and K was made due to financial constraint, which limited the analysis of the leaf and soil samples from all applied treatments, in all study sites. Thus, the analyzed soil and leaf samples helped to understand the status of the soil pH, and the concentrations of N, P and K in the soils and plant leaves at high rate of FYM, and at low and high rates of K after two-consecutive years of treatments application. This then, enabled to understand the effects of applied treatments on the soil pH and the concentration of N, P and K in the soil and cassava leaves.

Table 2: Soil pH and N, P and K concentrations in the experimental soils before treatments application in Bukoba, Missenyi and Biharamulo districts

District	Experimental site	Soil pH _w (1:2.5)	TN (%)	Bray – 1 P (mg kg ⁻¹)	Exch. K (cmol(+) kg ⁻¹)
Bukoba	TARI- Maruku	5.1 sta	0.41 m	1.38 l	0.15 l
Missenyi	Mabuye Primary School	6.0 ma	0.13 l	35.32 h	0.51 m
Biharamulo	Rukaragata Extension Centre	5.3 sta	0.18 l	4.96 l	0.16 l

Source: [18]

Chemical property: TN = total nitrogen, Bray-1 P = extractable Bray-1 phosphorus, K = exchangeable potassium; Rating: sta = strong acid; ma = moderate acid; l = low; h = high; m = medium [29].

In addition, the farmyard manure applied in each site was collected from farmers, processed, packed in paper bags, well labelled and analyzed for N, P and K. The contents of N, P and K in the applied farmyard manure and the amount of N, P and K added annually in the soils in each cropping season due to farmyard manure application in each experimental site are presented in Table 3.

Table 3: Contents of N, P and K in the applied farmyard manure and annual addition of N, P and K in the soil from farmyard manure

Experimental site	Cropping season	Content of N, P and K in FYM (%)			Addition of N, P and K (kg ha ⁻¹) in soil from FYM					
		N	P	K	4 MT		8 MT			
		N	P	K	N	P	K	N	P	K
Bukoba	2018/19	0.50	0.09	1.21	20.00	3.56	48.40	40.00	7.12	96.80
	2019/20	0.52	0.09	1.25	20.80	3.44	50.00	41.60	6.88	100.00
Missenyi	2018/19	0.54	0.12	1.51	21.60	4.80	60.40	43.20	9.60	120.80
	2019/20	0.56	0.11	1.54	22.40	4.40	61.60	44.80	8.80	123.20
Biharamulo	2018/19	0.58	0.08	1.78	23.20	3.04	79.20	46.40	6.08	142.40
	2019/20	0.57	0.08	1.83	22.80	3.00	73.20	45.60	6.00	146.40

Source: [20]

2.5 Data analysis

2.5.1 Analysis of soil samples

Laboratory analysis for soil pH and concentrations of N, P and K in the soils was done at Sokoine University of Agriculture (SUA) Soil Science Laboratory following standard soil analysis procedures as described by [21] and [22].

2.5.2 Analysis of leaf samples

Leaf samples sieved through 0.5 mm (20 mesh) sieve were digested using wet digestion method, whereby, leaf samples for extraction of P and K were digested using the nitric acid-hydrogen peroxide ($\text{HNO}_3 - \text{H}_2\text{O}_2$) wet digestion procedure. The solutions (digests) for determining P were prepared following colour development by the molybdenum blue method [19, 23] and the contents of P in the digests (leaf samples) were determined using an ultraviolet-visible (UV/VIS) spectrophotometer [24]. Concurrently, the solutions (digests) for determining K were prepared and the contents of K in the digests were determined using flame emission spectrometer. The contents of N in the leaf samples were determined using the macro-Kjeldahl digestion method [19, 25]. The contents of N, P and K in cassava plant leaves obtained after laboratory analyses were compared by the required standard levels of those nutrients in cassava leaves based on the ratings by [26].

2.6 Statistical data analysis

Data from leaf and soil samples were subjected to analysis of variance (ANOVA) based on the statistical model for the split-plot design [27] (Equation 1) using GENSTAT 15th Edition Statistical Packages. Means differences were separated using the Tukey's test [Honestly Significant Difference (HSD)] at $P = .05$ level of significance.

$$Y_{ijk} = \mu + \beta_i + A_j + \delta_{ij} + B_k + AB_{jk} + \epsilon_{ijk} \quad (1)$$

$$i = 1, 2, 3, \dots, r$$

$$j = 1, 2, 3, \dots, a$$

$$k = 1, 2, 3, \dots, b$$

Where: Y_{ijk} = Response level/Yield, μ = General effect or general error mean, β_i = Blocking effect, A_j = Main plot effect, δ_{ij} = Main plot random error (Error a), B_k = Sub-plot effect, AB_{jk} = Interaction effect between the main plot and the sub-plots, ϵ_{ijk} = Sub-plot random error (Error b), i = replication/blocking (r), j = main plot (a), k = sub-plot (b)

3. RESULTS AND DISCUSSION

3.1 Effect of applied treatments on soil chemical properties

3.1.1 Effect of tillage methods on soil pH and concentration of nitrogen (N), phosphorus (P) and potassium (K) in the soils in Bukoba, Missenyi and Biharamulo districts

The results on the effects of flat tillage, open ridging and tied ridging on soil pH and the concentration of N, P and K in the post-harvest soil in Bukoba, Missenyi and Biharamulo districts are presented in Table 4. Soil pH ranged from 5.3 - 5.4, which is strongly acid (Bukoba district), from 6.1- 6.2, which is slightly acid (Missenyi district) and from 5.0 - 5.1 which is very strongly acid (Biharamulo district). The concentration of N in the soil ranged from 0.34 - 0.37%, which is medium (Bukoba district), was 0.10%, which is low (Missenyi district) and ranged from 0.11 - 0.12%, which is low (Biharamulo district). Extractable P in the soil ranged from 1.08 - 1.19 mg kg⁻¹, which is low (Bukoba district), from 26.56 - 33.11 mg kg⁻¹, which is high (Missenyi district) and from 0.98 - 1.40 mg kg⁻¹, which is low (Biharamulo district). Exchangeable K in the soil ranged from 0.21 - 0.34 mg kg⁻¹, which is low to medium (Bukoba district), from 0.40 - 0.58 mg kg⁻¹, which is medium (Missenyi district) and from 0.20 - 0.23 mg kg⁻¹, which is low (Biharamulo district) [28, 29, 30].

Comment [O10]: this range must be controlled

Comment [O11]: this results not required must be gives in table

In all the study sites, there was no significant ($P = .46$) difference in the recorded soil chemical properties among the tested tillage methods, which signified that the use of flat tillage, open ridging or tied ridging had no effects on the soil pH and the concentration of N, P and K in the soils.

Table 4: Soil pH and concentration of N, P and K in the soil with respect to different tillage methods in Bukoba, Missenyi and Biharamulo districts

Treatment	Location											
	1	2	3	1	2	3	1	2	3			
	Soil pH _w (1:2.5)			Total N (%)			Bray-1 P (mg kg ⁻¹)			Exchangeable K (cmol(+) kg ⁻¹)		
Flat tillage	5.3 ^a sta	6.1 ^a sla	5.0 ^a vsta	0.34 ^a m	0.10 ^a l	0.11 ^a l	1.08 ^a l	26.56 ^a h	0.98 ^a l	0.21 ^a l	0.40 ^a m	0.20 ^a l
Open ridging	5.4 ^a sta	6.2 ^a sla	5.0 ^a vst	0.36 ^a m	0.10 ^a l	0.12 ^a l	1.17 ^a l	28.43 ^a h	1.40 ^a l	0.33 ^a m	0.55 ^a m	0.21 ^a l
Tied ridging	5.4 ^a sta	6.2 ^a sla	5.0 ^a vsta	0.37 ^a m	0.10 ^a l	0.12 ^a l	1.19 ^a l	33.11 ^a h	1.25 ^a l	0.34 ^a l	0.58 ^a m	0.23 ^a l
SED	0.08	0.05	0.06	0.02	0.01	0.00	0.11	2.57	0.26	0.04	0.08	0.02
CV (%)	4.50	2.70	3.50	13.60	34.10	12.60	28.70	26.20	30.60	31.10	32.90	28.20

Means within a column (for a particular nutrient) followed by the same letter(s) are not significantly ($P = .05$) different according to Turkey's HSD Test

SED = standard error of differences of means; CV = coefficient of variation; Location: 1 = Tanzania Agricultural Research Institute (TARI), Maruku Centre in Bukoba district, 2 = Mabuye primary school in Missenyi district, 3 = Rukaragata extension Centre in Biharamulo district; Chemical property: TN = total nitrogen, Bray-1 P = extractable Bray-1 phosphorus, K = exchangeable potassium; Rating: vsta = very strongly acid, sta = strongly acid, sla = slightly acid, vl = very low l = low, h = high, m = medium [28, 29,30].

3.1.2 Effect of combined use of farmyard manure and potassium fertilizers on soil pH and concentration of N, P and K in the post-harvest soil in Bukoba, Missenyi and Biharamulo districts

The results on the effects of combined use of FYM at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ or FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ on soil pH and concentration of N, P and K in the post-harvest soils in Bukoba, Missenyi and Biharamulo districts are presented in Table 5. Soil pH ranged from 5.1 - 5.5, which is strongly acid (Bukoba district), from 5.8 - 6.3, which is slightly acid (Missenyi district) and from 4.8 - 5.4 which is very strongly acid to strongly acid (Biharamulo district). Total nitrogen ranged from 0.20 - 0.49%, which is low to medium (Bukoba district), from 0.08 - 0.14%, which is very low to low (Missenyi district) and ranged from 0.09 - 0.20%, which is very low to low (Biharamulo district). Extractable P in the post-harvest soil ranged from 0.77 - 1.30 mg kg⁻¹, which is low (Bukoba district), from 13.28 - 33.54 mg kg⁻¹, which is medium to high (Missenyi district) and from 0.90 - 2.14 mg kg⁻¹, which is low (Biharamulo district). Exchangeable K in the post-harvest soil ranged from 0.13 - 0.34 (cmol(+) kg⁻¹), which is low to medium (Bukoba district), from 0.24 - 0.87 (cmol(+) kg⁻¹), which is low to high (Missenyi district) and from 0.11 - 0.33 (cmol(+) kg⁻¹), which is low to medium (Biharamulo district) [28, 29, 30].

Comment [O12]: it is good

Comment [O13]: this results not required

In all study sites, there was a significant ($P < .001$) difference in the recorded soil chemical properties between the control and fertilizer treatments, signifying that combined use of FYM at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ or FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ had significant effects on the soil pH and on the concentration of N, P and K in the soils. However, there was no a significant difference in the recorded soil chemical properties between combined use of FYM at 8 MT ha⁻¹ + potassium at 120 kg K ha⁻¹ and FYM at 8 MT ha⁻¹ + potassium at 40 kg K ha⁻¹. The low values of soil chemical properties were recorded in the control while the high values were recorded in the combined use of FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹. When comparing the status of the recorded chemical properties in the soil before treatments application (Table 1) and after treatments application (Table 5), the results indicate a slight increase or decrease in the respective soil chemical properties (Table 6).

The results also indicate that in Bukoba district, the control treatment did not cause any change in soil pH while the combined use of FYM at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ or the combined use of FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ increased soil pH by 7.27%. In Missenyi and Biharamulo districts, the control treatment decreased soil pH by 3.45% and 10.42%, respectively, whereas, combined use of FYM at 8 MT ha⁻¹ + potassium

at 40 kg K ha⁻¹ and combined use of FYM at 8 MT ha⁻¹ + potassium at 120 kg K ha⁻¹ increased soil pH by 4.76% and 1.85%, respectively. These results conform to the findings by [12] who reported significant increase in soil pH due to combined use of farmyard manure and inorganic N and P fertilizers as compared to the control, inorganic N fertilizer alone or inorganic P fertilizer alone. The decrease in soil pH in the control (no fertilizer application) was probably attributed to mineralization of inherent soil organic matter, which releases hydrogen ions in the soil [31]. The increase in soil pH upon use of FYM and potassium fertilizers was attributed to ability of farmyard manure to absorb or bind hydrogen ions in its humic forms [12, 32]. Other reason was continued addition of potassium (one of basic cations) from both FYM and muriate of potash (MOP) fertilizer.

Table 5: Soil pH and concentration of N, P and K in the soil with respect to combined use of farmyard manure and potassium rates in Bukoba, Misenyi and Biharamulo districts

Treatment	Location											
	1	2	3	1	2	3	1	2	3	1	2	3
	Soil pH _w (1:2.5)			Total N (%)			Bray-1 P (mg kg ⁻¹)			Exchangeable K (cmol(+) kg ⁻¹)		
Co	5.1 ^a sta	5.8 ^a ma	4.8 ^a vsta	0.20 ^a l	0.08 ^a vl	0.09 ^a vl	0.77 ^a l	13.28 ^a m	0.90 ^a l	0.13 ^a l	0.24 ^a l	0.11 ^a l
FYM ₈ + K ₄₀	5.5 ^b sta	6.3 ^b sla	5.4 ^b sta	0.49 ^b m	0.14 ^b l	0.20 ^b l	1.30 ^b l	37.54 ^b h	2.14 ^b l	0.31 ^b m	0.83 ^b h	0.31 ^b m
FYM ₈ + K ₁₂₀	5.5 ^b sta	6.3 ^b sla	5.4 ^b sta	0.49 ^b m	0.14 ^b l	0.20 ^b l	1.30 ^b l	37.53 ^b h	2.14 ^b l	0.34 ^b m	0.87 ^b h	0.33 ^b m
SED		0.06	0.06	0.02	0.01	0.00	0.12	2.27	0.26	0.04	0.08	0.02
CV (%)	4.50	2.70	3.50	13.60	34.10	12.60	28.70	26.20	30.60	31.10	32.90	28.20

Means within a column (for a particular nutrient) followed by the same letter(s) are not significantly ($P = .05$) different according to Turkey's HSD Test

SED = standard error of differences of means; CV = coefficient of variation; Location: 1 = Tanzania Agricultural Research Institute (TARI), Maruku Centre in Bukoba district, 2 = Mabuye primary school in Misenyi district, 3 = Rukaragata extension Centre in Biharamulo district; Treatment: Co = control (no fertilizer application), FYM₈K₄₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹, FYM₈K₁₂₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹; Chemical property: TN = total nitrogen, Bray-1 P = extractable Bray-1 phosphorus, K = exchangeable potassium; Rating: vsta = very strongly acid, sta = strongly acid, ma = medium acid, sla = slightly acid, vl = very low l = low, h = high, m = medium [28, 29, 30].

Table 6: Percentage change in soil pH and the contents of N, P and K in the soil with respect to combined use farmyard manure and potassium rates in Bukoba, Misenyi and Biharamulo districts

Treatment	Location											
	1	2	3	1	2	3	1	2	3	1	2	3
	Soil pH _w (%)			Total N (%)			Bray-1 P (%)			Exchangeable K (%)		
Co	0.0	-3.45	-10.42	-105.0	-62.50	-100.0	-79.22	-165.96	-451.11	-15.38	-112.38	-45.45
FYM ₈ + K ₄₀	7.27	4.76	1.85	16.33	13.33	10.0	-6.15	5.91	-131.78	51.61	38.55	48.39
FYM ₈ + K ₁₂₀	7.27	4.76	1.85	16.33	13.33	10.0	-6.15	5.89	-131.78	55.88	41.38	51.52

Location: 1 = Tanzania Agricultural Research Institute (TARI), Maruku Centre in Bukoba district, 2 = Mabuye primary school in Misenyi district, 3 = Rukaragata extension Centre in Biharamulo district; Treatment: Co = control (no fertilizer application), FYM₈K₄₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹, FYM₈K₁₂₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹; Chemical property: TN = total nitrogen, Bray-1 P = extractable Bray-1 phosphorus, K = exchangeable potassium

In all study sites, there was a decrease in total nitrogen (TN), extractable P and exchangeable K in the control treatment caused by the plant uptake and other soil losses such as leaching, since there was no addition of any fertilizer in the control treatment.

However, there was an increase in TN in the fertilizer treatments, with respect to the combined use of FYM at 8 MT ha⁻¹ + potassium at 40 kg K ha⁻¹ or the combined use of FYM at 8 MT ha⁻¹ + potassium at 120 kg K ha⁻¹. This was attributed to the addition of nitrogen in the soil from FYM as compared to the control. Comparable results were reported by [33] and [34] and conform to the findings by [1], [35] and [36] who reported significantly increase in total N due to combined use of farmyard manure and inorganic N and P or N, P and K fertilizers. The results also indicate that higher values of N were recorded in Bukoba site than in Missenyi and Biharamulo sites. This was because N was not a limiting nutrient in Bukoba site but was a limiting nutrient in Missenyi and Biharamulo sites [18], as a result; cassava plants took up most of the added nitrogen in the soil from the applied FYM in Missenyi and Biharamulo sites. This was supported by the low levels of N recorded in the post-harvest soils in Missenyi and Biharamulo sites as compared to medium level of N recorded in the post-harvest soil in Bukoba site (Table 5) according to the ratings by [29] and [30].

In Missenyi site, the results indicate an increase in extractable P in the fertilizer treatments while in Bukoba and Biharamulo sites there was a decrease in extractable P in the fertilizer treatments. This was because, P was not a limiting nutrient in Missenyi site but was a limiting nutrient in both Bukoba and Biharamulo sites [18], whereas, the added P from FYM in the soils in Bukoba and Biharamulo sites, were taken up by the cassava plants for growth and development. Thus, the increase in extractable P in Missenyi sites, was attributed to continued addition of P in the soil from FYM due to the combined use of FYM at 8 MT ha⁻¹ + potassium at 40 kg K ha⁻¹ or the combined use of FYM at 8 MT ha⁻¹ + potassium at 120 kg K ha⁻¹ as also reported by [1]. Other researchers [37, 38], reported significant increase in available P in the soil due to the combined use of FYM and inorganic N, FYM and P and K fertilizers or FYM and inorganic N and P fertilizers. Although the amount of P added from FYM was low in all sites (Table 3), there was high amount of inherent soil P in Missenyi site (Table 2), which was taken up by cassava plants while the unused part remained in the soil. This was supported by the high levels of extractable P recorded in the post-harvest soils in the fertilizer treatments in Missenyi site as compared to the low levels and the decreased extractable P in the post-harvest soils in the fertilizer treatments recorded in Bukoba and Biharamulo sites.

Moreover, the results indicate an increase in exchangeable K in the fertilizer treatments in all study sites, due to continued addition of K into the soil from both FYM and potassium fertilizer. Other researchers [39], reported significant increase in exchangeable K in the soil upon the combined use of farmyard manure and inorganic N, P and K fertilizers due to mineralization of FYM and solubilization of inorganic N, P and K fertilizers. It should be noted that farmyard manure used in this study had high content of K and thus the amount of K added in the soils was also high as compared to N and P. The results also indicate that there were medium levels of exchangeable K in the post-harvest soils in Bukoba and Biharamulo sites, and high level of exchangeable K in the post-harvest soil in Missenyi site, which signified that not all the amount of K added in the soils was taken up by cassava plants, instead, the unused portion remained in the soils. Therefore, the results from this study revealed that the combined use of FYM and potassium fertilizer increased the concentration of K in the soils. This was supported by the high levels and increased exchangeable K in the post-harvest soils in the fertilizer treatments in Missenyi site, and medium levels and increased exchangeable K, in the post-harvest soils in the fertilizer treatments in Bukoba and Biharamulo sites (Tables 5 and 6).

3.2 Effect of the applied treatments on the contents of nitrogen (N), phosphorus (P) and potassium (K) in cassava leaves in Bukoba, Missenyi and Biharamulo districts

3.2.1 Effects of tillage methods on the contents of N, P and K in the soils in cassava leaves Bukoba, Missenyi and Biharamulo districts

The results on the effects of flat tillage, open ridging and tied ridging on the concentration of N, P and K in cassava leaves soil in Bukoba, Missenyi and Biharamulo districts are presented in Table 7. The results indicate that the concentration of N in cassava leaves ranged from 3.07 - 3.26% (Bukoba district), from 3.14 - 3.44% (Missenyi district) and from 2.91 - 3.04% (Biharamulo district). The concentration of P in cassava leaves ranged from 0.16 - 0.20% (Bukoba district), from 0.26 - 0.27% (Missenyi district) and from 0.21 - 0.23% (Biharamulo district). Moreover, the concentration of K in cassava leaves ranged from 0.62 - 0.67% (Bukoba district), from 0.66 - 0.70% (Missenyi district) and from 0.38 - 0.45% (Biharamulo district).

In all study sites, the results indicate that all the tillage methods used in this study gave very deficient to deficient levels of N, P and K in cassava leaves according to the ratings by [25] (Table 2). There was no significant ($P = .55$) difference in the concentration of N, P and K in cassava leaves with respect to different tillage methods, implying that different tillage methods (flat tillage, open ridging and tied ridging) used in this study had no effects on the concentration of N, P and K in cassava leaves.

Comment [O14]: the reader your articles understand your work from table therefore not required

Comment [O15]: this range not common in analysis data it is better correct tp $p=0.05$

Table 7: The concentration of N, P and K in cassava leaves with respect to tillage methods in Bukoba, Missenyi and Biharamulo districts

Treatment	Location								
	1			2			3		
	Nitrogen (%)			Phosphorus (%)			Potassium (%)		
Flat tillage	3.16 ^a vd	3.44 ^a vd	3.04 ^a vd	0.16 ^a vd	0.27 ^a de	0.21 ^a vd	0.67 ^a vd	0.67 ^a vd	0.42 ^a vd
Open ridging	3.26 ^a vd	3.35 ^a vd	2.91 ^a vd	0.16 ^a vd	0.26 ^a de	0.22 ^a vd	0.62 ^a vd	0.66 ^a vd	0.45 ^a vd
Tied ridging	3.07 ^a vd	3.14 ^a vd	2.94 ^a vd	0.20 ^a vd	0.26 ^a de	0.23 ^a vd	0.65 ^a vd	0.70 ^a vd	0.38 ^a vd
SED	0.12	0.18	0.13	0.02	0.02	0.02	0.03	0.05	0.03
CV (%)	11.3	16.50	12.90	25.10	23.60	19.00	11.70	22.10	18.00

Means within a column (for a particular nutrient) followed by the same letter(s) are not significantly ($P = .05$) different according to Turkey's HSD Test

SED = standard error of differences of means; CV = coefficient of variation; Location: 1 = Tanzania Agricultural Research Institute (TARI), Maruku Centre in Bukoba district, 2 = Mabuye primary school in Missenyi district, 3 = Rukaragata extension Centre in Biharamulo district; Rating: vd = very deficient, de = deficient [26].

3.2.2 Effects of combined use of farmyard manure and potassium fertilizers on the concentration of N, P and K in cassava leaves in Bukoba, Missenyi and Biharamulo districts

The results on the concentration of N, P and K in cassava leaves with respect to the control treatment and combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 or 120 kg K ha⁻¹ in Bukoba, Missenyi and Biharamulo districts are presented in Table 8. The results indicate that the concentration of N in cassava leaves ranged from 1.43 - 4.53% (Bukoba district), from 1.00 - 4.77% (Missenyi district) and from 1.46 - 4.09% (Biharamulo district). The concentration of P in cassava leaves ranged from 0.13 - 0.20% (Bukoba district) and from 0.18 - 0.29% (Missenyi and Biharamulo districts). Moreover, the results indicate that the concentration of K in cassava leaves ranged from 0.25 - 1.07% (Bukoba district), from 0.17 - 1.20% (Missenyi district) and from 0.07 - 0.75% (Biharamulo district). According to the ratings by [40], the control treatment and combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 or 120 kg K ha⁻¹ gave very deficient to deficient levels of N, P and K in cassava leaves in all studied sites.

Comment [O16]: similar above comment

The observed very deficient to deficient levels of N, P and K in the cassava leaves in all study sites, was probably, attributed to the slow release of nutrients from FYM in the soil during the decomposition. This was because, at 4 months after planting (leaves sampling time), FYM might have not undergone complete decomposition to release enough nutrients into the soil solution for plant uptake. Other researchers, for example, [41] and [42], reported low use efficiency of farmyard manure due its slow release of nutrients into the soil upon decomposition.

Table 8: The concentration of N, P and K in cassava leaves with respect to combined use of farmyard manure and potassium rates in Bukoba, Missenyi and Biharamulo districts

Treatment	Location								
	1			2			3		
	Nitrogen (%)			Phosphorus (%)			Potassium (%)		
Co	1.43 ^a vd	1.00 ^a vd	1.46 ^a vd	0.13 ^a vd	0.18 ^a de	0.18 ^a vd	0.25 ^a vd	0.17 ^a vd	0.07 ^a vd
FYM ₈ K ₄₀	3.33 ^b vd	3.15 ^b vd	3.34 ^b vd	0.19 ^b vd	0.27 ^b de	0.26 ^b de	0.62 ^b vd	0.66 ^b vd	0.42 ^b vd
FYM ₈ K ₁₂₀	4.53 ^c de	4.77 ^c de	4.09 ^c de	0.20 ^b vd	0.29 ^b de	0.29 ^b de	1.07 ^c de	1.20 ^c de	0.75 ^c vd
SED	0.12	0.18	0.13	0.02	0.02	0.02	0.03	0.05	0.03
CV (%)	11.3	16.50	12.90	25.10	23.60	19.00	11.70	22.10	18.00

Means within a column (for a particular nutrient) followed by the same letter(s) are not significantly ($P = .05$) different according to Turkey's HSD Test

SED = standard error of differences of means; CV = coefficient of variation; Location: 1 = Tanzania Agricultural Research Institute (TARI), Maruku Centre in Bukoba district, 2 = Mabuye primary school in Missenyi district, 3 = Rukaragata extension Centre in Biharamulo district; Treatment: Co = control (no fertilizer application, FYM₈K₄₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹, FYM₈K₁₂₀ = farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹; Rating: vd = very deficient, de = deficient [25].

In all study sites, the results indicated a significant ($P < .001$) difference in the concentration of N, P and K in cassava leaves between the control and the combined use of farmyard manure and potassium rates, with low values of N, P and K recorded in the control. There was a significant ($P < .001$) difference in the concentration of N and K in cassava leaves between the combined use of FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ and the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹. Whereby, the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ gave significantly ($P < .001$) higher concentrations of N and K in cassava leaves than the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹. However, in all sites, there was no significant ($P = 0.44$) difference in the concentration of P in cassava leaves between the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ and the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹. This was probably attributed to the low amount of P added in the soil from the applied farmyard manure as oppose to the amounts of N and K added in the soil from the same farmyard manure [20].

4. CONCLUSION

Planting of cassava on the ridges or on the flat tillage had no effects on the soil pH and the concentrations of N, P and K in the soil and cassava leaves. The combined use of FYM at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ or the combined use of FYM at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ increased the soil pH and the concentration of N and K in the soil but did not increase the concentration of P in the soil and in cassava leaves following two consecutive years of applying FYM and potassium fertilizer. In addition, the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ gave higher concentrations of N and K in cassava leaves than the combined use of farmyard manure at 8 MT ha⁻¹ and

Comment [O17]: is it 0.01? if it is, this correct. this probability range said that in 0.01 not significant therefore your results must be in to side normal curve farthest from means even ferthest of 0.01.

potassium at 40 kg K ha⁻¹. Therefore, from the results of this study, planting cassava on the flat tillage or on ridges together with the use of the combination of farmyard manure at 8 MT ha⁻¹ and potassium at 120 kg K ha⁻¹ is desirable for increasing the concentrations of N and K in the soil and in cassava leaves and thus recommended. However, for the resource-poor farmers who cannot afford the high rate of K, the combined use of farmyard manure at 8 MT ha⁻¹ and potassium at 40 kg K ha⁻¹ could be used. Since the results indicate low content of P in the applied farmyard manure (especially in Bukoba and Biharamulo districts) where P is a limiting nutrient, we also recommend basal application of inorganic P-containing fertilizer for increasing the concentration of P in the soil and in cassava leaves.

REFERENCES

- [1] Dhaliwal SS, Naresh RK, Mandal A, Walia MK, Gupta RK, Singh, R, Dhaliwal, MK. Effect of manures and fertilizers on soil physical properties, build-up of macro and micronutrients and uptake in soil under different cropping systems: a review. *Journal of Plant Nutrition*, 2019;42(20):2873-2900.
- [2] Bulluck LR, Brosius M, Evanylo GK, Ristaino JB. Organic and synthetic fertility amendments influence soil microbial, physical and chemical properties on organic and conventional farms. *Applied Soil Ecology*, 2002;19(2):147-160.
- [3] Zeidan MS. Effect of organic manure and phosphorus fertilizers on growth, yield and quality of lentil plants in sandy soil. *Research Journal of Agriculture and Biological Sciences*, 2007;3:748-52.
- [4] Rutkowska B, Szulc W, Sosulski T, Stępień W. Soil micronutrient availability to crops affected by long-term inorganic and organic fertilizer application. *Plant, Soil and Environment*, 2014;60(5):198-203.
- [5] Hao XH, Liu SL, Wu JS, Hu RG, Tong CL, Su YY. Effect of long-term application of inorganic fertilizer and organic amendments on soil organic matter and microbial biomass in three subtropical paddy soils. *Nutrient Cycling in Agro-ecosystems*, 2008;81(1):17-24.
- [6] Dhaliwal SS, Walia SS. Integrated nutrient management for sustaining maximum productivity of rice-wheat system under Punjab conditions. *Journal of Research*, 2008;45(1): 12-16.
- [7] Hepperly P, Lotter D, Ullsh CZ, Seidel R, Reider C. Compost, manure and synthetic fertilizer influences crop yields, soil properties, nitrate leaching and crop nutrient content. *Compost Science and Utilization*, 2009;17(2):117-126.
- [8] Schoebitz M, Vidal G. Microbial consortium and pig slurry to improve chemical properties of degraded soil and nutrient plant uptake. *Journal of Soil Science and Plant Nutrition*, 2016;16(1):226-236.
- [9] Jones C, Olson-Rutz K, Dinkins CP. Nutrient uptake timing by crops: to assist with fertilizing decisions. Montana, USA. Montana State University; 2015.
- [10] Mahmoud E, Nasser AE, Robin P, Nouraya AC, Lamyaa AE. Effect of different organic and inorganic fertilizers on cucumber yield and some soil properties. *World Journal of Agricultural Sciences*, 2009;5:408-414.
- [11] Sanchez CA, Doerge TA. Using nutrient uptake patterns to develop efficient nitrogen management strategies for vegetables. *Horticultural Technology*, 199;9(4): 601-606.

- [12] Munyabarezi I. Effect of farmyard manure and mineral fertilizers on maize yield and soil properties in Huye and Bugesera districts of Rwanda. Thesis for award of Master of Science Degree at the Kenya University, Nairobi, Kenya; 2014.
- [13] United Republic of Tanzania (URT). Kagera Region Investment Guide. KageraTanzania: Kagera Regional Commissioner Office;2019.
- [14] MerumbaMS, Msanya BM, Semu E,Semoka JM. Pedological characterization and suitability assessment for cassava production in Bukoba, Missenyi and Biharamulo districts, Tanzania. American Journal of Agriculture and Forestry.2020a;8(4):144-166.
- [15] Oosterom AP, Ngailo JA, Kileo RO, Mbogoni JDJ, Msangi AS, Andriesse W, et al and Van Kekem, A. J. (1999). Land Resources of Biharamulo district, Kagera region, Tanzania. International Activities Report 75. Wageningen The Netherlands: Winand Staring Centre; 1999.
- [16] Bajjukya FP, Folmer ECR. Agro-Ecological Zonation of the Kagera region. In: Planning the Future: Past, Present and Future Perspectives of Land Use in KageraRegion. FolmerECR, Schouten C,Bajjukya FP, editors.Bukoba, KageraTanzania; 1999.
- [17] United Republic of Tanzania (URT). National Sample Census of Agriculture 2007/2008 Regional ReportofKageraRegion. Volume V. Dar es SalaamTanzania: National Bureau of Statistics; 2012.
- [18] Merumba MS, Semu E, Semoka JM, Msanya BM. Soil Fertility Status in Bukoba, Missenyi and Biharamulo districts in Kagera Region, Tanzania. International Journal of Applied Agricultural Sciences, 2020b;6(5): 96-117.
- [19] Shekiffu CY. Improving soil productivity in cassava-based system in the Coast Region of Tanzania: Phosphorus and Potassium requirements under mono-cropping and intercropping. Thesis for Award of PhD Degree at Sokoine University of Agriculture. Morogoro Tanzania; 2011.
- [20] Merumba MS, Semoka JM,Semu E, Msanya BM, Kibura JK. Effect of tillage methods, farmyard manure and potassium rates on cassava yield and root quality in Kagera, Tanzania. International Journal of Plant and Soil Science,2022; 34(13):33-50.
- [21] Moberg JP. Soil Analysis Manual. Revised edition. Copenhagen Denmark:The Royal Veterinary and Agricultural University, Chemistry Department;2001.
- [22] Okalebo JR, Gathua KW, Woomer PL. Laboratory Methods of Soil and Plant Analysis: A Working Manual No. 128. 2nd ed. Nairobi, Kenya. Tropical Soil Biology and Fertility Institute of the International Centre for Tropical Agriculture; 2002.
- [23] Murphy J, Riley JP. Modified single solution method for determination of phosphate in natural waters. AnalyticaChimicaActa, 1962;27:31-36.
- [24] Chapman HD. Cation exchange capacity. In: Black CA, editor. Methods of Soil Analysis. American Institute of Agronomy. 1965;9:891-901.
- [25] BremnerJM,Mulvaney CS. Total nitrogen. In:Black et al, editors.Methods of Soil Analysis. Part 2. Agronomy Monograph 9. MadisonWisconsinUSA.American Society of Agronomy; 1982.
- [26] Howeler RH. Cassava Mineral Nutrition and Fertilizer. In: Hillocks RJ, Thresh JM, Belloti AC, editors. Cassava Biology, Production, and Utilization. WallingfordUnited Kingdom: Centre for Agriculture and Bioscience International Publishing; 2002.

- [27] Montgomery DC. Design and Analysis of Experiments. 8th ed. New York: John Wiley and Sons Incorporation; 2013.
- [28] EuroConsult. Agricultural Compendium for Rural Development in the Tropics and Subtropics. 3rd ed. Amsterdam. Elsevier Science Publishers; 1989.
- [29] Baize D. Soil Science Analyses. A Guide to Current Use. West Sussex. John Wiley and Sons Ltd; 1993.
- [30] Msanya BM, Kimaro DN, Kimbi GG, Kileo EP, Mbogoni JDJ. Land resources inventory and suitability assessment for the major land use types in Morogoro Urban district, Tanzania: Soils and Land Resources of Morogoro Rural and Urban, Volume 4. Morogoro. Tanzania. Department of Soil Science, Sokoine University of Agriculture; 2001.
- [31] Brady NC, Weil RR. The Nature and Properties of Soil. 15th ed. Essex England. Pearson Education; 2017.
- [32] Tisdale SL, Nelson WL, Beaton JD, Havlin JL. Soil Fertility and Fertilizer. 5th ed. New Delhi India. Prentice-Hall of India; 1995.
- [33] Lemanowicz J, Siwik-Ziomek A, Koper J. Effects of farmyard manure and nitrogen fertilizers on mobility of phosphorus and sulphur in wheat and activity of selected hydrolases in soil. International Agrophysics, 2014;28(1):49-55.
- [34] Mahmood F, Khan I, Ashraf U, Shahzad T, Hussain S, Shahid M, Abid M, Ullah S. Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. Journal of Soil Science and Plant Nutrition, 2017;17(1):22-32.
- [35] Hemalatha S, Chellamuthu S. Impacts of long-term fertilization on soil nutritional quality under finger millet-maize cropping sequence. Journal of Environmental Research and Development, 2013;7(4):1571-1576.
- [36] Shahid M, Shukla AK, Bhattacharyya P, Tripathi R, Mohanty S, Kumar A, Lal B, Gautam P, Raja R, Nayak AK. Micronutrients (Fe, Mn, Zn and Cu) balance under long-term application of fertilizer and manure in a tropical rice-rice system. Journal of Soils and Sediments, 2016;16(3):737-747.
- [37] Singh G, Singh SHER, Singh SS. Integrated nutrient management in rice and wheat crop in rice-wheat cropping system in lowlands. Annals of Plant and Soil Research, 2013;15(1):1-4.
- [38] Tana T, Woldesenbet M. Effect of combined application of organic and mineral nitrogen and phosphorus fertilizer on soil physico-chemical properties and grain yield of food barley (*Hordeum vulgare L.*) in Kaffa Zone, South-western Ethiopia. Momona Ethiopian Journal of Science, 2017;9(2):242-261.
- [39] Jatav MK, Sud KC, Trehan SP. Effect of organic and inorganic sources of phosphorus and potassium on their different fractions under potato-radish cropping sequence in a brown hill soil. Journal of Indian Society of Soil Science, 2010; 58(4):388-393.
- [40] Howeler R H. (2002). Cassava Mineral Nutrition and Fertilization. In: Cassava: Biology, Production and Utilization. Hillocks, R.J, Thresh MJ, Bellotti AC, editors. Wallingford, Oxon, United Kingdom. Centre for Agriculture and Bioscience International Publishing; 2002.
- [41] Sanginga N, Wooster PL, editors. Integrated Soil Fertility Management in Africa: Principles, Practices and Development Process. Nairobi Kenya. Tropical Soil Biology and Fertility Institute of the International Centre for Tropical Agriculture; 2009.

- [42] Bayu W, Rethman NFG, Hammes PS, Alemu G. Effects of farmyard manure and inorganic fertilizers on sorghum growth, yield and nitrogen use in a semi-arid area of Ethiopia, *Journal of Plant Nutrition*, 2006; 29 (2):391-407.

ACRONYMS, ABBREVIATIONS

AGRA	Alliance for Green Revolution in Africa
ANOVA	Analysis of variance
CEC	Cation exchange capacity
CV	Coefficient of variation
°C	Degree Celsius
DAP	Di-ammonium phosphate
et al.	and others
FYM	Farmyard manure
GENSTAT	General statistics
g	Gram
GPS	Global positioning system
HNO ₃	Nitric acid
H ₂ O ₂	Hydrogen peroxide
HSD	Honestly Significant Difference
kg ha ⁻¹	Kilogram per hectare
masl	meter above sea level
m	meter
mm	millimeter
MOP	Muriate of potash
MT ha ⁻¹	Metric ton per hectare
OM	Organic matter
N, P, K, Mg, S	Nitrogen, phosphorus potassium, magnesium, sulphur
%	Percent
pH	Negative logarithm of hydrogen ion (H ⁺) concentration
RCBD	Randomized Complete Block Design
SUA	Sokoine University of Agriculture
TARI	Tanzania Agricultural Research Institute
TN	Total nitrogen
URT	United Republic of Tanzania
UV/VIS	Ultra-violet visible spectrophotometer