

Original Research Article

Isolation and characterization of endophytic plant growth promoting *Kosakoniaradicincitans* to stimulate wheat growth in saline soil

Among abiotic stresses, salinity is a significant limiting factor affecting agricultural productivity, survival, and production, resulting in significant economic losses. The goal of this study was to isolate the endophytic *Kosakoniaradicincitans* bacteria, characterize it and study of its plant growth-promoting activities (PGP) to determine the effect of its inoculation on wheat plants under salinity stress conditions. One endophytic bacterial isolate was obtained from root nodules of faba bean (*Vicia faba*) plants grown in the salt-affected clay soil of Egypt. The isolate was identified and characterized using 16S rRNA gene sequencing. This isolate phylogenetically was closely related to *Kosakoniaradicincitans* strain DSM 16656 with accession number OM980222.1. The isolate was evaluated for its ability to promote plant growth *in vitro*. Results revealed that the bacterial isolate of *Kosakoniaradicincitans* produced indole-3-acetic acid ($40.44 \mu\text{gml}^{-1}$) and exopolysaccharides (14.2 g/L), fixed nitrogen and could solubilize phosphate and potassium (273 mg/L and 42.8 $\mu\text{g/ml}$) respectively. In a field trial, this bacterial isolate reduced proline, and showed the best results for chlorophyll, grain yield, and 100-grain weight of wheat plants grown in the salt-affected soil as compared to the control plants. It can be concluded that wheat plants may benefit from the use of *Kosakoniaradicincitans* as biofertilizers in sustainable farming techniques.

Comment [O.I.1]: The goal of this study was to isolate and characterize the endophytic *Kosakonia radincitans* bacteria and their plant growth-promoting (PGP) activities to determine the effect of its inoculation on wheat plants under salinity stress conditions.

Comment [O.I.2]: Was phylogenetically related

Keywords: Endophytic bacteria, *Kosakoniaradicincitans*, Wheat, 16S rRNA

1. INTRODUCTION

Globally, rising salt levels are a serious concern and a significant barrier to food production [1]. Due to the rising annual need for food by a population that is expanding worldwide, the issue of soil salinity is critical. [2] confirmed that agriculture causes environmental damages, owing to large fertilizer inputs and land use practices. Living microbe cells called "biofertilizers" can replace artificial fertilizers because they give plants nutrition, prevent illnesses from spreading through the soil, and enhance the health and quality of the soil [3].

Endophytic N_2 -fixing bacteria are of particular interest because they spend at least part of their lives inside plants and cause no visible harm to the host [4]. Despite the high specificity between Rhizobia and legumes, non-rhizobia have been discovered inside the root nodules, including *Pseudomonas fluorescense*, *Bacillus* sp., *Enterobacter* sp., *Klebsiella* sp., and *Paenibacillus* sp. [5],[6],[7]. Many studies confirmed the diversity of bacteria other than rhizobia that are associated with root nodules [8], [9]. [10] proved that *Enterobacter* and *Klebsiella*, members of the family Enterobacteriaceae known to live endophytically in a variety of crops' root nodules.

Comment [O.I.3]: Italicize

Comment [O.I.4]: Italicize

To comprehend the ecological significance of the relationship between endophytes and their host plants, studies on the interactions between them are crucial [11]. These microorganisms could serve as biocontrol agents, release phytohormones like indole-3-acetic acid, cytokinins, and gibberellins, fix nitrogen so that it becomes available to the plant via cellular degeneration or active nitrogenous compound release [12].

Kosakoniaradicincitans (DSM 16656) is a gram-negative, rod-shaped bacterium that stimulates plant growth and has been identified in the newly recognized genus *Kosakonia* of the Enterobacteriaceae [13]. Previously class

Comment [O.I.5]: Italicize

Enterobacter, the genus *Kosakonia* was recently separated as a new genus [14]. The 16S rRNA gene is present in almost all bacteria and is usually used to their identification [15].

Comment [O.I.6]: Italicize

Comment [O.I.7]: Italicize

Many *Kosakonia* species have been isolated from different plants and have demonstrated the ability to promote plant growth [16]. The isolate of *K. radicinans*(DSMT 16656) from wheat has plant growth promotion (PGP) [16] and nitrogen-fixing abilities [17]. [18] also demonstrated that *K. radicinans* is capable of biologically fixing atmospheric nitrogen, producing hormones, and solubilizing rock phosphates. *K. radicinans*DSM 16656 has shown distinct enhancement in growth, yield, and product quality in both glasshouse and field trials [19] [16], highlighting the strain's potential for various cultivation management systems.

Comment [O.I.8]: Italicize

Since wheat makes up most of the Egyptian diet, it is regarded as the country's first strategic food crop. Wheat straw is also a significant source of fodder. [20]. This crop has been affected by water stress leading to decline in grain yield and quality. Different strategies to diminish the negative effects of salt stress have been developed to improve seed quality and production of wheat [21].

The present work intends to isolate the endophytic bacterial strain *Kosakonia radicinans* from the nodules of faba bean, characterize it through 16S rRNA based molecular technique and study of its plant growth-promoting activities as well as to determine the effect of its inoculation on wheat plants under salinity stress conditions.

Comment [O.I.9]: Present it in passt tense

2. MATERIAL AND METHODS

2.1 Isolation of endophytic bacteria

Nodules were chosen at random from roots of three faba bean plants grown in the salt-affected clay soil (Sahl El-Hussinia, El-Sharkia Governorate, Egypt), washed with sterile distilled water to remove soil particles, and surface sterilized with 95% alcohol for 30 s and 0.1 % (w/v) HgCl₂ for 2 min before being rinsed 6-8 times with sterile distilled water to completely remove HgCl₂. For the isolation of endophytic bacteria, the surface sterilized nodules were crushed and streaked on yeast-extract-mannitol agar (YEM) as indicated by [22],[23] and defined growth medium (DM) as indicated by [24]. Single colonies were purified further by streaking on the same medium repeatedly for 3 days at 28°C. The isolates were kept at -20°C in glycerol (20% v/v).

2.2 DNA isolation and 16S rRNA amplification

Isolation and purification of DNA was carried out according to the method described in [25]. The total genomic DNA was extracted from bacterial species and quality was evaluated on 1% agarose gel. Isolated DNA was amplified with 16S rRNA gene, the PCR amplification was carried out in 20 µl reaction solution containing 27F/1492R universal primers [26] as shown in Table 1. The amplification was performed: initial denaturation at 95°C for 5 min followed by 35 cycles comprised of denaturation at 94°C for 45 sec, 55° C for 60 sec for annealing, extension at 72° C for 60 sec and the final extension at 72°C for 30 min. The PCR included a positive control (*E. coli* genomic DNA) and a negative control. The PCR products resolved by electrophoresis in 1.5 % agarose gel.

Comment [O.I.10]: Italicize

2.3 DNA Sequencing and Phylogenetic analysis

The PCR amplicon purification was carried out by column-based method using Montage PCR Clean-up kit (Macro gen). The purified PCR products were sequenced by using two primers, as described in Table 1. Sequencing was performed using Big Dye terminator cycle sequencing kit and the products were resolved on an Applied Biosystems model 3730XL automated DNA sequencing system (Applied Bio Systems, USA). Identification was performed by (GenBank) using the Basic Local Alignment Search Tool (BLAST). Phylogenetic analysis was done using the MEGA X program [27].

Comment [O.I.11]: Biosystems

Table 1. The primer sequences used for 16S rRNA gene amplification.

Primers	Sequences
16S rRNA	27F 3' AGA GTT TGA TCM TGG CTC AG 5'
	1492R 3' TAC GGY TAC CTT GTT ACG ACT T 5'
Sequencing	518F 3' CCA GCA GCC GCG GTA ATA CG 5'
	800R 3' TAC CAG GGT ATC TAA TCC 5'

2.4 Plant growth promoting attributes

Indole acetic acid (IAA) production

According to the modified [28] technique, isolate was used for IAA production. The test isolate was inoculated into Luria-Bertani (LB) broth that had been supplemented with tryptophan. In a shaker incubator, a flask was incubated at 35°C for three days at 150 rpm. The well-developed isolate culture was centrifuged at 8000 rpm for 20 min. Then, 2 mL of Salkowski reagent and three drops of orthophosphoric acid were added to 1 mL of the supernatant. For one to two hours, the mixture was incubated in darkness. At 530 nm, the isolate's absorbance was observed. The IAA concentration in cell-free supernatant was determined colorimetrically using a standard curve prepared from authentic IAA (Sigma Chemical, USA).

Exopolysaccharides (EPS) production

The bacterial isolate was inoculated into a conical flask containing 100 ml of yeast extract mannitol (YEM) broth to estimate EPS production. The inoculated flasks were incubated for 72 hours at 30°C and 200 rpm on a rotary shaker. The culture broth was centrifuged at 3500 g after incubation, and the supernatant was mixed with two volumes of acetone. Centrifugation at 3500 g for 30 min was used to collect the crude polysaccharides that had developed. After being washed with distilled water and acetone alternately, the EPS was transferred to filter paper and weighed after being dried overnight at 105°C. [29].

Nitrogen fixing ability

Nitrogen fixing ability was examined using Jensen's medium and Bromothymol Blue (BTB) as a color indicator according to [30].

Estimation of phosphate solubilization

Qualitative determination of phosphate solubilization was performed on Pikovskaya's agar plate (PVK) medium [31]. Isolate was spot inoculated and incubated at 28 ± 2°C, and the size of the halo corresponding to phosphate solubilization was measured after 3-7 days of incubation. Phosphate solubilization was expressed as solubilizing efficiency (SE%) [32], [33]. Quantitative estimation of tri-calcium phosphate solubilization was performed by growing the bacterial strain in Pikovskaya's broth. The concentration of the soluble phosphate was determined from the culture supernatant.

$$\text{Phosphate solubilization efficiency (\%)} = \frac{\text{Solubilization diameter (S)} \times 100}{\text{growth diameter}}$$

Estimation of potassium solubilization

The spotting method was used to investigate potassium solubilization by bacterial isolate on Aleksandrov medium plates containing insoluble mica powder as a potassium source [34]. Plates were incubated for 5 days at 28°C±2. The ability of bacterial isolate to form solubilization zones was used to detect potassium solubilization. Quantitative estimation of potassium release was performed by growing bacterial isolate in Aleksandrov broth medium and incubated for 5 days at 28°C±2. Following the incubation, the broth culture was filtered through Whatman No. 1 filter paper and centrifuged for 20 minutes at 12,000 rpm. The soluble K content in the supernatant was measured using flame photometer [35].

2.5 Field Experiment

A field experiment was conducted at the farm El-Rowad village in Sahl El-Hussinia, El-Sharkia Governorate, Egypt during the winter growing season of 2018. Wheat grains (*Triticum aestivum* L. cv. Sakha 93) were obtained from Field Crops Research Institute, Agricultural Research Centre, Giza, and were inoculated with vermiculite-based inoculants that had been gamma irradiated. Three treatments were tested and a randomized completed design with three replicates was used to determine the role of isolated bacteria in alleviating salt stress in wheat plants grown in a saline soil as compared to the control and *Azotobacter chroococcum* (a ref PGPB and nitrogen fixer for nonlegumes) that was obtained from Biofertilizers Production Unit, Agricultural Microbiology Research Department at Soils, Water and Environment Research Institute, Agricultural Research Centre, Giza. The treatments were as follow:

1. Control (Recommended dose of NPK)
2. *Azotobacter chroococcum*+ half dose of NPK
3. *Kosakoni aradicincitans*+ half dose of NPK

Proline determination

Extraction and determination of proline in shoot of wheat plants (60 days from sowing) were performed according to the method of [36]. Proline content was measured by spectrophotometer at 520 nm and calculated as μ mole proline / g of fresh weight material using a standard curve prepared from proline.

Estimation of total chlorophyll content

The chlorophyll content was determined at 60 days after planting using portable chlorophyll meter (SPAD-502) [37].

Determination of N, P, K contents and protein

Comment [O.I.12]: Nonlegumes

Total contents of nitrogen, phosphorus and potassium of the harvested wheat grains were determined according to [38], and the protein was calculated by multiplying total nitrogen content by 5.75 [39].

Soil analysis

The main soil properties of the experimental soil were determined as shown in Table (2) according to the methods described by [40]. Soil pH and total soluble salt (EC) were measured in soil paste according to [41]. Available N was measured according to the modified Kjeldahl method [40]. Available P was determined colorimetrically according to Olsen's method [41]. Available K was that determined using the Flame-Photometer and Available micronutrients were determined as described by [42].

Comment [O.I.13]: Delete

Table (2): Physical and chemical properties of the soil sample before planting

Course sand (%)	Fin sand (%)	Silt (%)	Clay (%)	Texture	O.M (%)	CaCO ₃ (%)		
4.2	15.2	35.5	45.3	Clay	0.47	11.5		
pH (1:2.5)	EC (dS/m)	Cations (meq ⁻¹)				Anions (meq ⁻¹)		
		Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ²⁻
8.12	12.45	22.5	35.5	68.5	1.48	4.5	95.5	27.98
N	P	K	Zn	Mn	Fe	Cu		
PPM								
107.1	2.95	171.16	0.868	2.16	1.85	0.61		

2.6 Statistical analysis

The significance of variation in the Yield, nutrients uptake and protein contents such as N, P, K, protein, Proline and chlorophyll contents in relation to the different treatments were assessed using one-way analysis of variance technique (ANOVA). The differences were tested using Tukey test. The applications of these techniques were according to SPSS program ver. 20. [43].

Comment [O.I.14]: yield

Comment [O.I.15]: Please cross check the spelling

3. RESULTS AND DISCUSSION

3.1 Appearance of bacterial isolate

From root nodules of faba bean plants, gram-negative, rod-shaped bacterium was isolated. When cultivated on YEMA containing Congo red, the isolate did not absorb the color red.

3.2 Molecular characterization of the isolates by 16S rRNA gene

Molecular characterization of the isolate *Kosakoniaradicincitans* strain DSM 16656 was achieved using the 16S rRNA gene. The amplified size designed for the 16S rRNA gene, about 1500 bp, was used (Table1). The PCR products were separated in gel, and the DNA bands conforming to the expected size of the gene were purified from the gel for sequencing. With the use of the forward and reverse primers, the product was sequenced, resulting in a 940 bp long sequence. This isolate's partial sequence was aligned with the partial sequences of the neighbor-joining sequences in GenBank and then deposited under accession number MO980222.1 and described by the phylogenetic tree (AM3) in Fig. 1.




Figure 1. Phylogenetic tree based on 16S rDNA sequence analysis indicates the phylogenetic relationship of the bacterial isolate.

3.3 In Vitro Assessment of plant-growth-promoting activities

Bacterial isolate of *Kosakonia radicincitans* was examined for its plant growth promoting attributes in terms of indole-3-acetic acid (IAA) production, exopolysaccharides production, nitrogen fixation, phosphate (P) and potassium (K) solubilization (Table 3 and Fig. 2). Results indicated that the isolate was able to produce indole acetic acid (40.44 µg/ml) in the cultural filtrate. This agrees with [44] who proved that *K. radicincitans* produced a significant IAA and the capacity to produce IAA suggested that the isolate could be used to regulate growth.

Concerning the releasing amounts of EPS, it revealed that the bacterial isolate could produce EPS (14.2 g/L). These results agree with [45] who used EPS producing strain of *Kosakonia cowanii* for biosynthesis of EPS by several agro-industrial residues.

Jensen's medium, which is nitrogen-free, was created to find and grow bacteria that fix nitrogen. Several nitrogen fixing endophytic bacteria is acid producing and change the color of media from greenish blue to yellow. The *in vitro* test on Jensen's nitrogen free media showed that the new bacterial isolate could grow on Jensen's medium and was acid producer. This agrees with [46] who suggested that *Kosakonia* sp. was an associative nitrogen-fixing bacteria or endophytic nitrogen-fixing bacteria.

It is known that microorganisms can solubilize phosphorous, making it easily accessible to plants and lowering the demand for phosphate fertilizers [47]. In the current study, the isolate showed halo zone on the Pikovskaya's agar medium that recorded solubilization efficiency up to 100%, and P liberated in the medium was 273 mg/L indicating its ability to solubilize tricalcium phosphate (Table 3 and Fig. 2). This agrees with [24] who proved that osmotic stress on *K. radicincitans* cells increased their capacity to solubilize phosphate to 357.7 mg L⁻¹ from 290.3 mg L⁻¹ in non-stressed cells.

The ability to solubilize potassium by the bacterial isolate was tested. The isolate was able to solubilize potassium effectively and recorded higher solubilization efficiency up to 266% as shown in (Table 3 and Fig. 2). Quantitative estimation of K solubilization was performed and the amount of K liberated in the broth medium was 42.8µg/ml. This may be happened due to the production of organic and inorganic acids during growth of bacterial isolate which is a major mechanism for mineral solubilization as indicated by [48], where there was a decreasing in final pH reaching a value 5.3. This agrees with [49] who proved that pH decrease resulting from the release of organic acids could be related to P & K solubilization. These PGP activities may be due to sharing the bulk of genes that involved in plant growth-promotion as indicated with [46] who found that there were many genes related to promoting plant growth in *Enterobacter* FY-07.

Comment [O.I.16]: This result was in agreement with

Comment [O.I.17]: Are in agreement

Comment [O.I.18]: changes

Comment [O.I.19]: agreed

Comment [O.I.20]: result is in agreement

Comment [O.I.21]: delete

Comment [O.I.22]: decrease

Comment [O.I.23]: pH,

Table (3). Production of IAA, EPS, solubilization of phosphate and potassium of the isolate

Isolate	IAA ($\mu\text{g ml}^{-1}$)	EPS (g/l)	Nitrogen fixation	Final pH	Phosphate solubilization		Potassium solubilization	
					Solubilization efficiency (SE)%	P- liberated (mg/l)	Solubilization efficiency (SE)%	K- liberated ($\mu\text{g/ml}$)
AM3	40.44	14.2	+	5.3	100	273	266	42.8




Figure (2). (a) Isolate grown on Congo red YEMA medium (b) IAA production, (c) Exopolysaccharides production (d) Phosphate solubilization (e) Potassium solubilization by bacterial isolate.

3.4 Proline content in wheat

Under salt stress, most of plant species show a striking rise in their proline content [50]. It is obvious from the present results that application of new isolate onto wheat plants lowered the proline accumulation in these plants as compared to control and inoculant of *Azotobacterchroococcum* a ref bacterial strain (Fig 3). These results are consistent with [51] who confirmed that proline content in maize increased under saline stress however it was decreased by inoculation with plant growth promoting bacteria. These findings may be due to the synthesis indole acetic acid or exopolysaccharides by the bacterial isolate as reported by [52] and [53] respectively, where foliar applications of IAA and inoculation with EPS producing bacteria as *A. chroococcum* reduced proline content in the leaves of maize plants under salt stress.




Figure 3. Proline content as affected by bacterial treatments

3.5 Total chlorophyll content

Results in Fig. 4 show that maximum content of chlorophyll in wheat leaves was obtained using bacterial isolate of *Kosakonia radicincitans* as compared to control or *Azotobacter* sp. In this respect [54] showed an increase in chlorophyll content and photosynthesis rates in barely with inoculation by phosphate solubilizing bacteria (PSB).




Figure 4. Total chlorophyll in wheat plant as affected by treatments

3.6 Yield, nutrients uptake and protein contents in wheat

There was a significant increase over *A. chroococcum* and the control in grain yield of wheat by the application of the new isolate of *K. radicincitans* (Table 4) that gave the highest value (17.6 Ardab/fed). Moreover, 100-grains weight of wheat was significantly influenced by the application of new bacterial isolate followed by *A. chroococcum* compared to the control but there was non-significant increase in straw yield of wheat in case of using the new isolate. These outcomes might be attributable to the isolate's capacity to create IAA, as suggested by [55] who found that the administration of IAA boosted wheat grain production.

In terms of the concentration of the nutrients in wheat plants as a result of treatments, data in (Table 4) revealed that the contents of N, P, K, and protein are not significantly different by application of new bacterial isolate + half dose of NPK as compared to *A. chroococcum* + half dose of NPK and the control treatment (Recommended dose of NPK). This could be due to the ability of the bacterial isolate to fix nitrogen and solubilize both phosphate, and potassium. Upon the inoculation of diverse plant species, [56] demonstrated that numerous *Kosakonia* bacterial strains could enhance plant development and increase yields where they could fix atmospheric nitrogen using nitrogenase that was present inside the bacteria [57]. [18] confirmed the capacity of *K. radicincitans* to synthesize hormones, physiologically fix atmospheric nitrogen, and dissolve rock phosphates. In this context, According to [58], *K. radicincitans* MUSA4 possessed a number of traits that aided in the promotion of plant growth, such as the production of indoleacetic acid, the solubilization of phosphate, and the fixation of nitrogen. Additionally, the bacterial genome contained the *nif* and *anf* gene clusters, which, respectively, encode the Fe-Mo and Fe-Fe nitrogenase systems. This was supported by [59], who found that *K. radicincitans* had multiple copies of complex gene clusters that enabled it to colonize and have growth-promoting effects on a variety of plants.

Comment [O.I.24]: please cross check this phrase

Comment [O.I.25]: indoleacetic

Table 4. Yield components, N, P, K and protein content in wheat at harvest.

Treatments	Grain Ardab / fed	Straw Ton/fed	Wt. 100 grains	N%	P%	K%	Protein
Control (full dose NPK)	10.7 ^c	1.7 ^a	3.33 ^b	1.08 ^a	0.560 ^a	0.3 ^a	6.21 ^a
<i>A. chroococcum</i> + half dose of NPK	15.6 ^b	1.7 ^a	3.73 ^{ab}	1.09 ^a	0.590 ^a	0.6 ^a	6.26 ^a
<i>K. radicincitans</i> (AM3) + half dose of NPK	17.6 ^a	1.8 ^a	4.02 ^a	1.56 ^a	0.585 ^a	0.5 ^a	8.97 ^a
F-Value	391.14***	0.23 ^{ns}	12.00**	3.98 ^{ns}	0.01 ^{ns}	1.75 ^{ns}	3.99 ^{ns}

** : P<0.01, ***: P<0.001, ns: non-significant. Means in the same column followed by the same letters are not significantly different (P<0.05) according to Tukey test.

4. CONCLUSION

It can be concluded that wheat plants may be benefited from the use of *Kosakonia radicincitans* as biofertilizers in sustainable farming techniques. Additional research in this area can support our findings.

Comment [O.1.26]: be of

REFERENCES

1. Alom R , Hasan M , Islam M, Wang Q. Germination characters and early seedling growth of wheat (*Triticumaestivum*L.) genotypes under salt stress conditions. J. Crop Sci. Biotechnol. 2016; 19: 383–392.
DOI: 10.1007/s12892-016-0052-1.
2. Yang L , Deng Y, Wang X , Zhang W, Shi X, Chen X. Global direct nitrous oxide emissions from the bio energy crop sugarcane (*Saccharum* spp. inter-specific hybrids). Sci.TotalEnviorn 2021; 752:141795.
DOI:10.1016/j.scitotenv.2020.141795.
3. Kour D , Rana KL, Yadav AN, Yadav N, Kumar M, Kumar V. Microbial biofertilizers: bioresources and eco-friendly technologies for agricultural and environmental sustainability. Biocatal. Agric. Biotechnol .2020; 23:101487.
DOI:10.1016/j.bcab.2019.101487.
4. Ryan RP, Germaine K, Franks A , Ryan DJ, Dowling DN . Bacterial endophytes: recent developments and applications. FEMS MicrobiolLett, 2008;278 (1):1–9.
DOI: 10.1111/j.1574-6968.2007.00918.
5. Rajendran G, Patel MH, Joshi SJ. Isolation and characterization of nodule associated Exiguobacterium sp. from the root nodules of Fenugreek (*Trigonellafoenum-graecum*) and their possible role in plant growth promotion. Int J Microbiol .2011;2012:1–8, 436.
DOI:10.1155/2012/693982.
6. Tariq M, Hameed S, Yasmeen T, Ali A. Non-rhizobial bacteria for improved nodulation and grain yield of mung bean [*Vignaradiata* (L.) Wilczek]. Afr J Biotechnol .2012;11(84):15012–15019.
DOI: 10.5897/AJB11.3438.
7. Pandya M, Kumar GN, Rajkumar S. Invasion of rhizobial infection thread by non-rhizobia for colonization of *Vignaradiata* root nodules. FEMS MicrobiolLett . 2013;348 (1):58-65.
DOI: 10.1111/1574-6968.
8. Mayhood P, Mirza BS (). Soybean Root Nodule and RhizosphereMicrobiome: Distribution of Rhizobial and NonrhizobialEndophytes. Appl. Environ. Microbiol. 2021;87 (10).
DOI: 10.1128/AEM.02884-20.
9. Deng, ZS, Kong ZY, Zhang BC, Zhao LF. Insights into non-symbiotic plant growth promotion bacteria associated with nodules of *Sphaerophysasalsula* growing in northwestern China, Arch. Microbiol. 2020; 202 (2):399-409.
DOI:10.1007/s00203-019-01752-7.
10. De Meyer SE, De Beuf K, Vekeman B, Willems A. A large diversity of non-rhizobialendophytes found in legume root nodules in Flanders (Belgium). Soil BiolBiochem. 2015;83:1–11.
DOI:10.1016/j.soilbio.2015.01.002.
11. Hardoim PR, van Overbeek LS, van Elsas JD . Properties of bacterial endophytes and their proposed role in plant growth Trends Microbiol. 2008;16(10): 463-471.
DOI: 10.1016/j.tim.2008.07.008.
12. Ferrara FIS, Oliveira ZM, Gonzales HHS, Floh EIS, Barbosa HR. Endophytic and rhizosphericenterobacteria isolated from sugar cane have different potentials for producing plant growth-promoting substances. Plant Soil. 2012;353:409-417.
DOI: 10.1007/s11104-011-1042-1.
13. Krey T, Baum C, Ruppel L , Seydel M , EichlerLobermann B. Organic and inorganic P sources interacting with applied rhizosphere bacteria and their effects on growth and P supply of maize. Commun Soil Sci Plan.2013 ;44(22):3205–3215.
<https://doi.org/10.1080/00103624>.
14. Brady C, Cleenwerck I, Venter S, Coutinho T De Vos P. Taxonomic evaluation of the genus *Enterobacter* based on multilocus sequence analysis (MLSA). Syst. Appl. Microbiol. 2013; 36: 309–319.
DOI: 10.1016/j.syapm.2013.03.005.

15. Rosselli R, Romoli O, Vitulo N, Vezzi A, Campanaro S, De Pascale F. Direct 16S rRNA-seq from bacterial communities: a PCR-independent approach to simultaneously assess microbial diversity and functional activity potential of each taxon. *Sci. Rep.* 2016; 6:32165. DOI: 10.1038/srep32165.
16. Berger B, Patz S, Ruppel S, Dietel K, Faetke S, Junge H, Becker M. Successful formulation and application of plant growth-promoting *Kosakonia radicitans* in maize cultivation. *Biomed Res Int.* 2018; 6439481:8. DOI: 10.1155/2018/6439481.
17. Remus R, Ruppel S, Jacob HJ, Hecht-Buchholz C, Merbach W. Colonization behaviour of two enterobacterial strains on cereals. *Biol Fert Soils.* 2000;30(5-6):550–557. DOI: 10.1007/s003740050035
18. Brock AK, Berger B, Mewis I, Ruppel S. Impact of the PGPB *Enterobacter radicitans* DSM 16656 on growth, glucosinolate profile and immune responses of *Arabidopsis thaliana*. *Microb Ecol.* 2013;65:661–670. DOI: 10.1007/s00248-012-0146-3.
19. Schreiner M, Krumbein A, and Ruppel S. Interaction between plants and bacteria: glucosinolates and phyllospheric colonization of cruciferous vegetables by *Enterobacter radicitans* DSM 16656. *J. Mol. Microbiol. Biotechnol.* 2009;17: 124–135. DOI: 10.1159/000226589.
20. Gaballah MS, Mandour MS. Increasing drought resistance of wheat plants during grain filling by using chemical desiccants. *J. Sci. Mansoura Univ.* 2000;25: 833-841. DOI: 10.21608/JPP.2000.258559.
21. El-Sabbagh A, Hossain A, Barutcular C, Iqbal MA, Islam MS, Fahad S. "Consequences of salinity stress on the quality of crops and its mitigation strategies for sustainable crop production: an outlook of arid and semi-arid regions," in *Environment, Climate, Plant and Vegetation Growth*, eds Fahad A, Hasanuzzaman M, Alam M, Ullah H, Saeed M, Khan IA, Adnan M (Cham: Springer). 2021;503–533.
22. Vincent, J.M.. *The cultivation, isolation and maintenance of rhizobia :A Manual for the Practical Study of the Root-Nodule Bacteria* 1970; 1-13.
23. Baldani VLD, Baldani JI, Olivares FL, Döbereiner J. Identification and ecology of *Herbaspirillum seropedicae* and the closely related *Pseudomonas rubrisubalbicans*. *Symbiosis.* 1992;13:65–73.
24. Barrera MC, Schoenwandt DJ, Gomez MI, Becker M, Patel AV, Ruppe S. Salt stress and hydroxyectoine enhance phosphate solubilisation and plant colonisation capacity of *Kosakonia radicitans* *Journal of Advanced Research*, 2019;19: 91-97.
25. Sambrook J, Russell D. *Molecular cloning a laboratory manual*. Vol. 2, 3rd ed. Cold Spring Harbor Laboratory Press, New York. 2001
26. Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA. Common SNPs explain a large proportion of the heritability for human height. *Heath AC, Martin NG, Montgomery GW, Goddard ME, Visscher PM. Nat Genet.* 2010;42 (7):565-9.
27. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution* 2016;33:1870-1874. DOI: 10.1093/molbev/msw054
28. Khamna S, Yokota A, Peberdy JF, Lumyong S. Indole-3-acetic acid production by *Streptomyces* sp. isolated from some Thai medicinal plant rhizosphere soils. *EurAsian J. BioSci.* 2010; 4:23–32. DOI: 10.5053/ejobios.2010.4.0.4
29. ASTM. *American standard for testing and materials*. New York; 2001.

30. Ahmad F, Ahmad I, Khan MS. Screening of free-living rhizospheric bacteria for their multiple plant growth promoting activities. *Microbiol Res.* 2008;163(2): 173– 181.
DOI: 10.1016/j.micres.2006.04.001.
31. Pikovskaya RI. Mobilization of phosphorous in soil connection with the vital activity of some microbial species. *Microbiologiya* 17:362-370. *Plant Soil*, 1948;287:77- 84.
32. Nguyen C, Yan W, Le T Lapeyrie F. Genetic variability of phosphate solubilizing activity by monocaryotic and dicaryotic mycelia of the ectomycorrhizal fungus *Laccaria bicolor* (Maire) P.D. Orton, *Plant Soil*, 1992;143:193– 199.
DOI:10.1007/PF00007873
33. Srivastava S, Yadav KR, Kundu BS. Prospects of using phosphate solubilizing *Pseudomonas* as biofungicide. *Indian J Microbiol.*, 2004;44: 91-94.
34. Sindhu SS, Gupta SK, Dadarwal KR. Antagonistic effect of *Pseudomonas* spp. on pathogenic fungi and enhancement of plant growth in green gram (*Vignaradiata*). *Biology and Fertility of Soils*, 1999;29: 62-68.
35. Sugumaran B, Janarthanam BS. Solubilization of potassium-containing minerals by bacteria and their effect on plant growth *World J. Agric. Sci.* 2007;3: 350–355.
36. Bates LS, Waldren RP, Teare JD. Rapid determination of free proline for water stress studies *Plant Soil.* 1973;39: 205-207.
<http://dx.doi.org/10.1007/BF00018060>
37. Peterson TA, Blackmer TM, Francis DD, Schepers JS. Using a chlorophyll meter to improve N management. *Nebguide G93-1171A. Coop. Ext. Nebraska, Lincoln. Serv., Univ. of Nebraska, Lincoln.* 1993.
38. Cottenie A, Verloo M, Kiekens L, Velghe G, Camerlynck R. Chemical analysis of plant and soils. *Laboratory of Analytical and Agrochemistry, State University of Gent, Belgium.* 1982.
39. AOAC. *Official Methods of Analysis of Association of Official Analytical Chemists.* 17th Ed. Washington, D.C. 2005.
40. Black C.A. "Methods of Soil Analysis". Ser. Agron. No 9 Amer. Soc. Agron., Madison, Wisconsin. Studies on growth characteristics and dry matter accumulation of pearl millet and sorghum hybrids. *Research Report of Rural Development Administration, Livestock.* 1965;32: 45-53.
41. Jackson ML. *Soil chemical analysis Constable and Co L.t.p., London* 1967;371- 387.
42. Soltanpour N, Schwab A. A new soil test for simultaneous extraction of macro and micronutrients in alkaline soils. *Commun. Soil Sci. plant Anal.* 1977;3: 195.
DOI:10.1080/00103627709366714.
43. SPSS. *SPSS Statistics for Windows version 20.0 Chicago, IL, USA, SPSS Inc, 2011.*
44. Singh RK, Singh P, Li HB, Guo DJ, Song Q, Yang L, Malviya MK, Song XP, Li YR. Plant PGPR interaction study of plant growth promoting diazotrophs *Kosakonia radicum* BA1 and *Stenotrophomonas maltophilia* COA2 to enhance growth and stress-related gene expression in *Saccharum* spp., *Journal of plant interactions*, 2020;15(1): 427- 445.
DOI: 10.1080/17429145.2020.1857857.
45. Gao H, Lu C, Wang H, Wang L, Yang Y, Jiang Y, Li S, Xu D, Wu L. Production exopolysaccharide from *Kosakonia cowanii* LT-1 through solid-state fermentation and its application as a plant growth promoter, *Int J Biol Macromol.* 2020;150:955-964.
DOI: 10.1016/j.ijbiomac.2019.10.209.
46. Gao G, Zhang Y, Niu S, Chen Y, Wang S, Anwar N, Chen S, Li G, Ma T. Reclassification of *Enterobacter* sp. FY-07 as *Kosakonia oryzae* FY-07 and Its Potential to Promote Plant Growth, *Microorganisms.* 2022; 10(3): 575.
DOI: 10.3390/microorganisms10030575
47. Lavakusha JY, Verma JP, Jaiswal D, Kumar A. Evaluation of PGPR and different concentration of phosphorus level on plant growth: yield and nutrient content of rice *Oryza sativa*. *Ecol Eng.* 2014;62:123–128.

- 48.Meena VS, Maurya BR, Verma JP, Aeron A, Kumar A, Kim K, Bajpai VK, Potassium solubilizing rhizobacteria (KSR): isolation, identification, and K-release dynamics from waste mica. *Ecol Eng.* 2015;81: 340–347.
DOI: 10.1016/j.ecoleng.2015.04.065
- 49.Fuentes B, Jorquera M, Mora ML. Dynamics of phosphorus and phytate-utilizing bacteria during aerobic degradation of dairy cattle dung. *Chemosphere*, 2009;74: 325-331.
DOI: 10.1016/j.chemosphere.2008.08.045.
- 50.Delauney AJ, Verma DPS. Proline biosynthesis and osmoregulation in plants. *Plant Journal.* 1993;4:215–223.
DOI: 10.1046/j.1365-313X.1993.04020215.x
- 51.Nadeem SM, Zahir ZA, Naveed M, Arshad M. Preliminary investigations on inducing salt tolerance in maize through inoculation with rhizobacteria containing ACC deaminase activity. *Can J Microbiol* 2007;53(10):1141–1149.
DOI: 10.1139/W07-081.
- 52.Kaya C, Tuna AL, Okant AM. Effect of foliar applied kinetin and indole acetic acid on maize plants grown under saline conditions *Turk J Agric For* .2010;34: 529-538.
DOI:10.3906/tar-0906-173
- 53.Tapias DR, Galvan AM, Diaz SP, Obando M, Rivera D, Bonilla R. Effect of inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize (*Zea mays*). *Applied Soil Ecology*.2012 ;61 : 264– 272.
DOI:10.1016/j.apsoil.2012.01.006
- 54.Mehrvarz, S, Chaichi MR, Alikhani, HA. Effects of phosphate solubilizing microorganisms and phosphorus chemical fertilizer on yield and yield components of Barely (*Hordeumvulgare L.*).*Am-Euras. J Agric& Environ Sci*, 2008; 3(6):822-828.
- 55.Kamboj A, Kler DS. Growth analysis and grain yield of wheat as influenced by nitrogen, growth regulator and micronutrients. *Environment and Ecology.* 2007;25: 496-499.
- 56.Berger B, Baldermann S, Ruppel S. The plant growth promoting bacterium *Kosakoniadicincitans* improves fruit yield and quality of *Solanumlycopersicum*. *J Sci Food Agri.* 2017;97(14):4865–4871.
DOI:10.1002/jsfa.8357
- 57.Bergottini VM, Otegui MB, Sosa DA, Zapata PD, Mulot M, Rebord M. Bio-inoculation of yerba mate seedlings (*Ilex paraguariensis* St. Hill.) with native plant growth-promoting rhizobacteria: a sustainable alternative to improve crop yield. *BiolFertil Soils.* 2015;51:749–755.
DOI: 10.1007/s00374-015-1012-5.
- 58.Nunes F Q, Rossi MJ, Nascimento XF. Genomic insights into the plant-associated lifestyle of *Kosakoniadicincitans* MUSA4, a diazotrophic plant-growth-promoting bacterium, *Systematic and Applied Microbiology* 2022;45(2):126303.
DOI: 10.1016/j.syapm.2022.126303.
- 59.Becker M, Patz S, Becker Y, Berger B, Drungowski M, Bunk B, Overmann J, Spröer C, Reetz J, Tchakounte G. Comparative Genomics Reveal a Flagellar System, a Type VI Secretion System and Plant Growth-Promoting Gene Clusters Unique to the Endophytic Bacterium *Kosakoniadicincitans*. *Front. Microbiol* : 2018;9:1997.
DOI: 10.3389/fmicb.2018.01997.