

# Fixed Point Theorems Using Soft Multiplicative Generalized Weak Contractive Mappings

**Abstract:** In the present paper, we establish fixed point theorems for mappings satisfying generalized weak contractive conditions in soft multiplicative metric space.

**Keywords:** Soft metric space, Soft multiplicative metric space, Soft multiplicative weak contractive mapping, Fixed point.

**MSC:** 47H10, 54H25

## 1. Introduction

In 1874, Cantor defined set theory as a branch of mathematics. This theory deals with the problem that contain certain results. But in real life situations, there are various uncertain problems which have imprecise results. To deal with these problems, Molodtsov[4], in 1999, introduced soft set theory and applied this theory in various fields like game theory, operations research etc. In 2002, Maji et al. [11][12] worked on soft set theory and its applications in decision making problems. In 2011, Ali et al. [8] defined various operations in this theory. In 2013, Wardowski [3] worked on soft mappings with the fixed point theorems.

As metric space is one of the prominent branches of mathematics, thus to explore this idea using soft sets, Das and Samanta [15][16] investigated the properties of soft real numbers in 2012 and in 2013, they introduced the concept of soft metric space. After this, several researchers worked on soft metric spaces and their properties. In 2008, Bashirov et al. [1] defined multiplicative metric space. Then various authors worked on this space [6][9][14][19][20][21]. Rathee et al. [17] combined soft metric space and multiplicative metric space and generated a new space called soft multiplicative metric space.

Fixed point theory plays a vital role in various fields of mathematics. In 2016, Wadkar et al. [2] proved fixed point results related to soft sets and in the same year, Yazar et al. [10] proved some fixed point theorems of soft contractive mappings. In 2017, Hosseinzadeh [5] proved fixed point theorems in soft metric space. Then, Abbas et al. [7] introduced various results on fixed point theorems in soft metric spaces. After this, in 2021, Bhardwaj et al. [13] investigated some new fixed point results in soft metric space. Rathee et al. [17] derived some fixed point theorems in soft multiplicative metric space. In 2017, Solankki et al. [18] generalize the concept of soft weak contractive mapping and proved various fixed point theorems in soft metric space. Extending the work of Solankki et al. [18] and Rathee et al. [17], we generate some new fixed point theorems using generalized multiplicative weak contraction mapping in soft multiplicative metric space.

## 2. Preliminaries

This section contains some basic definitions and results which are useful for our research work.

**Definition 2.1[15].** “Let  $I$  be an initial universal set and  $\Omega$  be the non-empty parameter set. Then, a pair  $(T, \Omega)$  is called a soft set over  $I$  if  $T$  is a set valued mapping on  $\Omega$  taking values in  $2^I$  i.e.,  $T : \Omega \rightarrow 2^I$ .”

**Definition 2.2[15].** “A soft set  $(T, \Omega)$  over  $I$  is said to be an absolute soft set if  $T(\alpha) = I \quad \forall \alpha \in \Omega$ . It is denoted by  $\tilde{I}$ .”

**Definition 2.3[15].** “A soft set  $(T, \Omega)$  over  $I$  is said to be a soft point if there is exactly one  $\alpha \in \Omega$  such that  $T(\alpha) = \{i\}$  for some  $i \in I$  and  $T(\beta) = \emptyset$  for all  $\beta \in \Omega \setminus \{\alpha\}$ . Such a soft point is denoted by  $T_\alpha^i$ .”

NOTE. The collection of all soft points of a soft set  $(T, \Omega)$  is denoted by  $SP(T, \Omega)$ .

**Definition 2.4[15].** “Let  $\mathbb{R}$  be the set of real numbers and  $B(\mathbb{R})$  be the collection of all non-empty bounded subset of  $\mathbb{R}$ . Then, the function by  $T : \Omega \rightarrow B(\mathbb{R})$  is called a soft real set and is denoted by  $(T, \Omega)$ . If  $T$  is a single valued function on  $\Omega$  taking values in  $\mathbb{R}$ , then the pair  $(T, \Omega)$  or simply  $T$  is called a soft real number. We denote soft real number and soft constant real number by  $\tilde{r}, \tilde{s}, \tilde{t}$  and  $\bar{r}, \bar{s}, \bar{t}$  respectively where  $\bar{r}$  will denote a particular type of soft real number such that  $\bar{r}(\alpha) = r$  for all  $\alpha \in \Omega$ .”

**Definition 2.5[15].** “For two soft real numbers  $\tilde{p}$  and  $\tilde{q}$ , the following conditions hold for all  $\alpha \in \Omega$ :

- (a)  $\tilde{p} \lesssim \tilde{q}$  if  $\tilde{p}(\alpha) \lesssim \tilde{q}(\alpha)$  ;
- (b)  $\tilde{p} \gtrsim \tilde{q}$  if  $\tilde{p}(\alpha) \gtrsim \tilde{q}(\alpha)$  ;
- (c)  $\tilde{p} < \tilde{q}$  if  $\tilde{p}(\alpha) < \tilde{q}(\alpha)$  ;
- (d)  $\tilde{p} > \tilde{q}$  if  $\tilde{p}(\alpha) > \tilde{q}(\alpha)$  .”

**Definition 2.6[15].** “A mapping  $\tilde{\rho} : SP(\tilde{I}) \times SP(\tilde{I}) \rightarrow R(\Omega)^*$  is a soft metric on the absolute soft set  $\tilde{I}$  if  $\tilde{\rho}$  satisfies the following conditions:

1.  $\tilde{\rho}(T_\alpha^i, T_\beta^j) \gtrsim \bar{0}$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
2.  $\tilde{\rho}(T_\alpha^i, T_\beta^j) = \bar{0}$  if and only if  $\alpha = \beta$  and  $i = j$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
3.  $\tilde{\rho}(T_\alpha^i, T_\beta^j) = \tilde{\rho}(T_\beta^j, T_\alpha^i)$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
4.  $\tilde{\rho}(T_\alpha^i, T_\gamma^k) \lesssim \tilde{\rho}(T_\alpha^i, T_\beta^j) + \tilde{\rho}(T_\beta^j, T_\gamma^k)$  for all  $T_\alpha^i, T_\beta^j, T_\gamma^k \in SP(\tilde{I})$ .

The soft set  $\tilde{I}$  together with soft metric  $\tilde{\rho}$  is called a soft metric space and is denoted by  $(\tilde{I}, \tilde{\rho}, \Omega)$  or simply by  $(\tilde{I}, \tilde{\rho})$ .”

**Definition 2.7[1].** “A mapping  $\rho^*: I \times I \rightarrow \mathbb{R}^*$  is multiplicative metric if  $d^*$  satisfies the following conditions:

1.  $\rho^*(u, v) \geq 1$  for all  $u, v \in I$ ;
2.  $\rho^*(u, v) = 1$  if and only if  $u = v$  for all  $u, v \in I$ ;
3.  $\rho^*(u, v) = \rho^*(v, u)$  for all  $u, v \in I$ ;
4.  $\rho^*(u, w) \leq \rho^*(u, v) \cdot \rho^*(v, w)$  for all  $u, v, w \in I$ .

The pair  $(I, \rho^*)$  is called a multiplicative metric space.”

**Definition 2.8[17].** “A function  $\tilde{\rho}^*: SP(\tilde{I}) \times SP(\tilde{I}) \rightarrow R(\Omega)^*$  is soft multiplicative metric on the absolute soft set  $\tilde{I}$  if  $\tilde{\rho}^*$  meets the following properties:

1.  $\tilde{\rho}^*(T_\alpha^i, T_\beta^j) \geq \bar{1}$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
2.  $\tilde{\rho}^*(T_\alpha^i, T_\beta^j) = \bar{1}$  if and only if  $\alpha = \beta$  and  $i = j$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
3.  $\tilde{\rho}^*(T_\alpha^i, T_\beta^j) = \tilde{\rho}^*(T_\beta^j, T_\alpha^i)$  for all  $T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ;
4.  $\tilde{\rho}^*(T_\alpha^i, T_\gamma^k) \leq \tilde{\rho}^*(T_\alpha^i, T_\beta^j) \tilde{\rho}^*(T_\beta^j, T_\gamma^k)$  for all  $T_\alpha^i, T_\beta^j, T_\gamma^k \in SP(\tilde{I})$ .

The soft set  $\tilde{I}$  together with soft multiplicative metric  $\tilde{\rho}^*$  is called a soft multiplicative metric space and is denoted  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ .”

**Definition 2.9[17].** “Suppose  $(\tilde{I}, \tilde{\rho}^*)$  is a soft multiplicative metric space. Then, a sequence  $\{T_{\alpha_n}^{i_n}\}$  in  $(\tilde{I}, \tilde{\rho}^*)$  is soft multiplicative convergent to a soft point  $T_\beta^j \in \tilde{I}$  if for given  $\tilde{\epsilon} \geq \bar{1}$ , we have a unique positive integer  $n_0$  such that  $\tilde{\rho}^*(T_{\alpha_n}^{i_n}, T_\beta^j) < \tilde{\epsilon}$  for all  $n \geq n_0$  i.e.,  $\tilde{\rho}^*(T_{\alpha_n}^{i_n}, T_\beta^j) \rightarrow \bar{1}$  as  $n \rightarrow \infty$ .”

**Definition 2.10[17].** “Suppose  $(\tilde{I}, \tilde{\rho}^*)$  is a soft multiplicative metric space. Then, a sequence  $\{T_{\alpha_n}^{i_n}\}$  in  $(\tilde{I}, \tilde{\rho}^*)$  is soft multiplicative Cauchy sequence if for given  $\tilde{\epsilon} \geq \bar{1}$ , we have a unique positive integer  $n_0$  such that  $\tilde{\rho}^*(T_{\alpha_m}^{i_m}, T_{\alpha_n}^{i_n}) \leq \tilde{\epsilon}$  for all  $m, n \geq n_0$  i.e.,  $\tilde{\rho}^*(T_{\alpha_m}^{i_m}, T_{\alpha_n}^{i_n}) \rightarrow \bar{1}$  as  $m, n \rightarrow \infty$ .”

**Definition 2.11[17].** “A soft multiplicative metric space  $(\tilde{I}, \tilde{\rho}^*)$  is complete, if every soft multiplicative Cauchy sequence in  $\tilde{I}$  converges to some soft point in  $\tilde{I}$ .”

**Definition 2.12[10].** “Let  $(\tilde{I}, \tilde{\rho}, \Omega)$  and  $(\tilde{I}', \tilde{\rho}', \Omega')$  be two soft metric spaces. Then,  $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}', \tilde{\rho}', \Omega')$  is a soft mapping where  $h: I \rightarrow I'$  and  $\psi: \Omega \rightarrow \Omega'$  are two mappings.”

**Definition 2.13[17].** “Consider a soft multiplicative metric space  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ . A function  $(h, \psi): (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$  is said to be soft multiplicative contraction mapping if for every soft point  $T_\alpha^i, T_\beta^j \in \tilde{I}$ , there exists a soft real number  $\bar{\lambda}, \bar{0} \lesssim \bar{\lambda} \lesssim \bar{1}$  such that  $\tilde{\rho}^*\{(h, \psi)(T_\alpha^i), (h, \psi)(T_\beta^j)\} \lesssim \{\tilde{\rho}^*(T_\alpha^i, T_\beta^j)\}^{\bar{\lambda}}$ .”

**Definition 2.14[18].** “A mapping  $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}, \Omega)$  where  $(\tilde{I}, \tilde{\rho}, \Omega)$  is a soft metric space is said to be soft weakly C-contractive or a soft weak contraction if  $\forall T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ,

$$\tilde{\rho}\{(h, \psi)(T_\alpha^i), (h, \psi)(T_\beta^j)\} \leq \frac{\bar{1}}{2} \left[ \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\beta^j)\} + \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\alpha^i)\} \right] - \xi \left[ \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\beta^j)\}, \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\alpha^i)\} \right],$$

where  $\xi: [\bar{0}, \infty)^2 \rightarrow [\bar{0}, \infty)$  is a continuous mapping such that  $\xi(T_\alpha^i, T_\beta^j) = \bar{0}$  if and only if one of  $T_\alpha^i, T_\beta^j = \bar{0}$ .”

**Definition 2.15[18].** “A mapping  $(h, \psi): (\tilde{I}, \tilde{\rho}, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}, \Omega)$  where  $(\tilde{I}, \tilde{\rho}, \Omega)$  is a soft metric space is said to be soft generalized weakly contractive or a soft generalized weak contraction if  $\forall T_\alpha^i, T_\beta^j \in SP(\tilde{I})$ ,

$$\tilde{\rho}\{(h, \psi)(T_\alpha^i), (h, \psi)(T_\beta^j)\} \leq \bar{\eta} \left[ \max \left\{ \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\alpha^i)\}, \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\beta^j)\}, \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\beta^j)\}, \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\alpha^i)\}, \tilde{\rho}^*(T_\alpha^i, T_\beta^j) \right\} \right] - \xi \left[ \left( \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\alpha^i)\}, \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\beta^j)\} \right), \left( \tilde{\rho}\{T_\alpha^i, (h, \psi)(T_\beta^j)\}, \tilde{\rho}\{T_\beta^j, (h, \psi)(T_\alpha^i)\}, \tilde{\rho}(T_\alpha^i, T_\beta^j) \right) \right],$$

where  $\bar{\eta} \in \left[ \bar{0}, \frac{\bar{1}}{2} \right)$ ,  $\xi: [\bar{0}, \infty)^5 \rightarrow [\bar{0}, \infty)$  is a continuous mapping such that  $\xi(T_\alpha^i, T_\beta^j, T_\gamma^k, T_\delta^l, T_\epsilon^m) = \bar{0}$  if and only if one of  $T_\alpha^i, T_\beta^j, T_\gamma^k, T_\delta^l, T_\epsilon^m = \bar{0}$ .”

### 3. Main Results

In this section, we define soft multiplicative generalized weakly contractive mappings and prove fixed point results using these mappings.

**Theorem 3.1.** Let  $(\tilde{I}, \tilde{\rho}^*, \Omega)$  be a complete soft multiplicative metric space and  $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$  be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping:

$$\tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), (h, \psi)(\tilde{T}_\beta^j) \right\} \leq \frac{\left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\} \right]^{\bar{p}} \left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\beta^j \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\alpha^i \right\} \right]^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\alpha^i \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\beta^j \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\alpha^i), \tilde{T}_\beta^j \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_\beta^j), \tilde{T}_\alpha^i \right\} \right]} \forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I}),$$

where  $\bar{p}, \bar{q}$  are non-negative soft real numbers such that  $\bar{p} + \bar{q} \lesssim \frac{1}{2}$  and  $\xi : [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$  is a continuous function such that  $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l) = \bar{1}$  iff one of  $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l = \bar{1}$ . Then, there exists a unique fixed point of  $(h, \psi)$ .

Proof. Let  $\tilde{T}_{\alpha_0}^{i_0}$  be any soft point in  $SP(\tilde{I})$ . Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi)\tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi)\tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi)\tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) &= \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\} \\ &\leq \frac{\left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n} \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \right]^{\bar{p}} \left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n} \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \tilde{\rho}^* \left\{ (h, \psi)(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]} \end{aligned}$$

$$\begin{aligned}
 & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \\
 & \lesssim \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}}}{\xi \left\{ \begin{array}{l} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \\ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \end{array} \right\}} \\
 & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \\
 & \lesssim \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}}}{\xi \left\{ \begin{array}{l} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \\ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \end{array} \right\}}
 \end{aligned}$$

Since  $\xi$  satisfies the given condition, thus

$$\xi \left\{ \begin{array}{l} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \\ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \end{array} \right\} = \bar{1}$$

and thus

$$\begin{aligned}
 \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}} \\
 & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{q}} \\
 \Rightarrow \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{1}-\bar{p}-\bar{q}} & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}+\bar{q}} \\
 \Rightarrow \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\frac{\bar{p}+\bar{q}}{\bar{1}-\bar{p}-\bar{q}}} \\
 \Rightarrow \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p}+\bar{q}}{\bar{1}-\bar{p}-\bar{q}} \\
 \Rightarrow \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n-2}}^{i_{n-2}} \right) \right\}^{\bar{\eta}^2} \\
 & \vdots \\
 \Rightarrow \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_1}^{i_1}, \tilde{T}_{\alpha_0}^{i_0} \right) \right\}^{\bar{\eta}^n}.
 \end{aligned}$$

For any  $m > n$ , where  $m, n \in \mathbb{N}$

$$\begin{aligned}
 \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) &\lesssim \{\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}})\tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m})\} \\
 &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\
 &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}}) \right\} \\
 &\quad \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\
 &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+2}} \\
 &\quad \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{m-1}} \\
 &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n(\bar{1} + \bar{\eta} + 2\bar{\eta}^2 + \dots + \bar{\eta}^{m-n-1})}} \\
 &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\frac{\bar{\eta}^n}{\bar{1} - \bar{\eta}}}.
 \end{aligned}$$

Since  $\bar{p} + \bar{q} < \frac{\bar{1}}{2}$ , thus  $\bar{\eta} < \bar{1}$  and hence  $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) \rightarrow \bar{1}$  as  $m, n \rightarrow \infty$ . So, the soft sequence  $\{\tilde{T}_{\alpha_n}^{i_n}\}$  is soft multiplicative Cauchy sequence in  $\tilde{I}$ . Being the completeness of  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ , there exists a soft point  $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$  such that  $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$  as  $n \rightarrow \infty$ .

Also,

$$\begin{aligned}
 \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} &\lesssim \left[ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right)\right\} \left\{ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \right] \right. \\
 &\quad \left. \left[ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\} \right]^{\bar{p}} \right. \\
 \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} &\lesssim \frac{\left[ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\} \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \right]^{\bar{q}} \left[ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \right]}{\xi \left[ \begin{array}{l} \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\}, \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \\ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \end{array} \right]} \\
 &\quad \left\{ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\} \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}\right) \right\}^{\bar{p}} \\
 &\lesssim \frac{\left\{ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\} \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}\right) \right\}^{\bar{q}} \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}\right)}{\xi \left[ \begin{array}{l} \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*}\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}\right), \\ \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}\right) \end{array} \right]}
 \end{aligned}$$

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{p}} \\ \left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{p}} & \lesssim \frac{\left\{ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right)}{\xi \left[ \begin{aligned} & \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \right. \\ & \left. \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]} \\ \left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{p}-\bar{q}} & \lesssim \bar{1} \text{ as } n \rightarrow \infty. \end{aligned}$$

Since  $\bar{p} + \bar{q} < \frac{1}{2}$ , thus  $\bar{1} - \bar{p} - \bar{q} > \bar{0}$  and hence  $\tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$ . This signifies that  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ .

Now, if  $\tilde{T}_{\alpha'}^{i'}$  be another “soft fixed point” of  $(h, \psi)$ . Then,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) & = \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\} \\ & \left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right), \tilde{T}_{\alpha'}^{i'} \right\} \right]^{\bar{p}} \\ & \lesssim \frac{\left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha'}^{i'} \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right), \tilde{T}_{\alpha^*}^{i^*} \right\} \right]^{\bar{q}}}{\xi \left[ \begin{aligned} & \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right), \tilde{T}_{\alpha'}^{i'} \right\}, \right. \\ & \left. \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{T}_{\alpha'}^{i'} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right), \tilde{T}_{\alpha^*}^{i^*} \right\} \right]} \\ & = \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{2\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1}-2\bar{q}} & \lesssim \bar{1} \end{aligned}$$

Since  $\bar{p} + \bar{q} < \frac{1}{2}$  and  $\bar{p} > \bar{0}$ , thus  $\bar{1} - 2\bar{q} > \bar{0}$  and hence  $\tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$ .

Hence, there is one and only one soft fixed point of  $(h, \psi)$ .

**Theorem 3.2.** Let  $(\tilde{I}, \tilde{\rho}^*, \Omega)$  be a complete soft multiplicative metric space and  $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$  be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping  $\forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I})$ ,

$$\tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\alpha^i \right), (h, \psi) \left( \tilde{T}_\beta^j \right) \right\} \leq \frac{\left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\alpha^i \right), \tilde{T}_\alpha^i \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\beta^j \right), \tilde{T}_\beta^j \right\} \right]^{\bar{q}} \left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\alpha^i \right), \tilde{T}_\beta^j \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\beta^j \right), \tilde{T}_\alpha^i \right\} \right]^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_\alpha^i, \tilde{T}_\beta^j \right), \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\alpha^i \right), \tilde{T}_\alpha^i \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\beta^j \right), \tilde{T}_\beta^j \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\alpha^i \right), \tilde{T}_\beta^j \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_\beta^j \right), \tilde{T}_\alpha^i \right\} \right]}$$

where  $\bar{p}$ ,  $\bar{q}$  and  $\bar{r}$  are non-negative soft real number such that  $\bar{p} + 2\bar{q} + 2\bar{r} \lesssim \bar{1}$  and  $\xi: [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$  is a continuous function such that  $\xi(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m) = \bar{1}$  iff one of  $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m = \bar{1}$ . Then, there exists a unique fixed point of  $(h, \psi)$ .

Proof. Let  $\tilde{T}_{\alpha_0}^{i_0}$  be any soft point in  $SP(\tilde{I})$ . Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi) \tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi) \tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi) \tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_n}^{i_n} \right), (h, \psi) \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{T}_{\alpha_n}^{i_n} \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \right]^{\bar{q}} \\ &\leq \frac{\left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\} \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{T}_{\alpha_n}^{i_n} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{T}_{\alpha_n}^{i_n} \right\} \right]} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{q}} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n} \right) \right]} \end{aligned}$$

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{q}} \\ & \leq \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{r}}}{\xi \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \right.} \\ & \quad \left. \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \right\}} \end{aligned}$$

Since  $\xi$  satisfies the given condition, thus

$$\xi \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \right. \\ \left. \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right), \bar{1} \right\} = \bar{1}$$

and thus

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{1}-\bar{q}-\bar{r}} \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{p}+\bar{q}+\bar{r}} \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\frac{\bar{p}+\bar{q}+\bar{r}}{\bar{1}-\bar{q}-\bar{r}}} \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p}+\bar{q}+\bar{r}}{\bar{1}-\bar{q}-\bar{r}} \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n-2}}^{i_{n-2}} \right) \right\}^{\bar{\eta}^2} \\ & \quad \vdots \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_1}^{i_1}, \tilde{T}_{\alpha_0}^{i_0} \right) \right\}^{\bar{\eta}^n}. \end{aligned}$$

For any  $m > n$ , where  $m, n \in \mathbb{N}$

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}} \right) \right\} \\ & \quad \dots \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+2}} \\ & \quad \dots \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{m-1}} \end{aligned}$$

$$\begin{aligned} &\lesssim \{\tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1})\}^{\bar{\eta}^{n(\bar{1}+\bar{\eta}+2\bar{\eta}^2+\dots+\bar{\eta}^{m-n-1})}} \\ &\lesssim \{\tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1})\}^{\frac{\bar{\eta}^n}{1-\bar{\eta}}}. \end{aligned}$$

Since  $\bar{p} + 2\bar{q} + 2\bar{r} < \bar{1}$ , thus  $\bar{\eta} < \bar{1}$ . Therefore,  $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) \rightarrow \bar{1}$  as  $m, n \rightarrow \infty$ . So, the soft sequence  $\{\tilde{T}_{\alpha_n}^{i_n}\}$  is soft multiplicative Cauchy sequence in  $\tilde{I}$ . Being the completeness of  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ , there exists a soft point  $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$  such that  $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$  as  $n \rightarrow \infty$ .

Also,

$$\begin{aligned} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} &\lesssim \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), (h, \psi)(\tilde{T}_{\alpha_n}^{i_n})\} \left\{ \tilde{\rho}^*(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n} \right\} \right] \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{p}} \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n}\} \right]^{\bar{q}} \\ &\quad \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\} \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha^*}^{i^*}\} \right]^{\bar{r}} \\ &\lesssim \frac{\left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha^*}^{i^*}\} \right]}{\xi \left[ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}), \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha_n}^{i_n}\}, \right. \\ &\quad \left. \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\}, \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha_n}^{i_n}), \tilde{T}_{\alpha^*}^{i^*}\} \right]} \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{p}} \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} \right]^{\bar{q}} \left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \right]^{\bar{q}} \\ &\quad \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\} \right]^{\bar{r}} \left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]^{\bar{r}} \\ &\lesssim \frac{\left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]}{\xi \left[ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}), \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\}, \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}), \right. \\ &\quad \left. \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\}, \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]} \\ &\quad \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{p}} \left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}) \right]^{\bar{q}} \\ &\quad \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\} \right]^{\bar{r}} \left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]^{\bar{r}} \\ &\quad \left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]} \\ \left[ \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\} \right]^{\bar{1}-\bar{q}} &\lesssim \frac{\left[ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]}{\xi \left[ \tilde{\rho}^*(\tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_n}^{i_n}), \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha^*}^{i^*}\}, \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_n}^{i_n}), \right. \\ &\quad \left. \tilde{\rho}^*\{(h, \psi)(\tilde{T}_{\alpha^*}^{i^*}), \tilde{T}_{\alpha_n}^{i_n}\}, \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha^*}^{i^*}) \right]} \end{aligned}$$

$$\left[ \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1} - \bar{q} - \bar{r}} \lesssim \bar{1} \text{ as } n \rightarrow \infty.$$

Since  $\bar{p} + 2\bar{q} + 2\bar{r} \lesssim \bar{1}$  and  $\bar{p} \gtrsim \bar{0}$ , therefore  $\bar{1} - \bar{q} - \bar{r} \gtrsim \bar{0}$  and hence  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ . Now, if  $\tilde{T}_{\alpha'}^{i'}$  be another “soft fixed point” of  $(h, \psi)$ . Then,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left[ \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right\} \right]^{\bar{q}} \\ &\leq \frac{\left[ \left\{ \tilde{\rho}^* \left( (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \right]^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right\}, \right. \\ &\quad \left. \left\{ \tilde{\rho}^* \left( (h, \psi) \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left\{ (h, \psi) \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \right]} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{q}} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{2\bar{r}} \\ &= \frac{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \right. \\ &\quad \left. \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right\} \right]}{\xi} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1} - \bar{p} - 2\bar{r}} \lesssim \bar{1} \end{aligned}$$

Since  $\bar{p} + 2\bar{q} + 2\bar{r} \lesssim \bar{1}$  and  $\bar{q} \gtrsim \bar{0}$ , therefore  $\bar{1} - \bar{p} - \bar{r} \gtrsim \bar{0}$ . Thus,

$$\tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}.$$

Hence, there is one and only one soft fixed point of  $(h, \psi)$ .

**Theorem 3.3.** Let  $(\tilde{I}, \tilde{\rho}^*, \Omega)$  be a complete soft multiplicative metric space and  $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$  be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping:

$$\tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_\alpha^i\right), (h, \psi)\left(\tilde{T}_\beta^j\right)\right\} \leq \frac{\left[\max \left[\tilde{\rho}^*\left\{\tilde{T}_\alpha^i, (h, \psi)\left(\tilde{T}_\alpha^i\right)\right\}, \tilde{\rho}^*\left\{\tilde{T}_\beta^j, (h, \psi)\left(\tilde{T}_\beta^j\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_\alpha^i, \tilde{T}_\beta^j\right)\right]^{\bar{p}}}{\left[\tilde{\rho}^*\left\{\tilde{T}_\alpha^i, (h, \psi)\left(\tilde{T}_\beta^j\right)\right\} \tilde{\rho}^*\left\{\left(\tilde{T}_\beta^j\right), (h, \psi)\left(\tilde{T}_\alpha^i\right)\right\}\right]^{\bar{q}}}$$

$$\xi \left[ \begin{array}{l} \tilde{\rho}^*\left\{\tilde{T}_\alpha^i, (h, \psi)\left(\tilde{T}_\alpha^i\right)\right\}, \tilde{\rho}^*\left\{\tilde{T}_\beta^j, (h, \psi)\left(\tilde{T}_\beta^j\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_\alpha^i, \tilde{T}_\beta^j\right), \\ \tilde{\rho}^*\left\{\tilde{T}_\alpha^i, (h, \psi)\left(\tilde{T}_\beta^j\right)\right\}, \tilde{\rho}^*\left\{\left(\tilde{T}_\beta^j\right), (h, \psi)\left(\tilde{T}_\alpha^i\right)\right\} \end{array} \right]$$

$$\forall \tilde{T}_\alpha^i, \tilde{T}_\beta^j \in SP(\tilde{I}),$$

where  $\bar{p}, \bar{q}$  are non-negative soft real numbers such that  $\bar{p} + 2\bar{q} \lesssim \bar{1}$  and  $\xi: [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$  is a continuous function such that  $\xi\left(\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m\right) = \bar{1}$  iff one of  $\tilde{T}_\alpha^i, \tilde{T}_\beta^j, \tilde{T}_\gamma^k, \tilde{T}_\delta^l, \tilde{T}_\kappa^m = \bar{1}$ . Then, there exists a unique fixed point of  $(h, \psi)$ .

Proof. Let  $\tilde{T}_{\alpha_0}^{i_0}$  be any soft point in  $SP(\tilde{I})$ . Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi)\tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi)\tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi)\tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$

Now,

$$\tilde{\rho}^*\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}\right) = \tilde{\rho}^*\left\{(h, \psi)\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right), (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right\}$$

$$\left[\max \left\{\tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right), (h, \psi)\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right)\right\}, \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_n}^{i_n}\right), (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}\right]^{\bar{p}}$$

$$\lesssim \frac{\left[\tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right), (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right\} \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_n}^{i_n}\right), (h, \psi)\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right)\right\}\right]^{\bar{q}}}{\xi \left[ \begin{array}{l} \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right), (h, \psi)\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right)\right\}, \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_n}^{i_n}\right), (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right), \\ \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right), (h, \psi)\left(\tilde{T}_{\alpha_n}^{i_n}\right)\right\}, \tilde{\rho}^*\left\{\left(\tilde{T}_{\alpha_n}^{i_n}\right), (h, \psi)\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}\right)\right\} \end{array} \right]}$$

$$\left[\max \left\{\tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right), \tilde{\rho}^*\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}\right), \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}\right]^{\bar{p}}$$

$$\lesssim \frac{\left\{\tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}\right) \tilde{\rho}^*\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}\right)\right\}^{\bar{q}}}{\xi \left[ \begin{array}{l} \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right), \tilde{\rho}^*\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}\right), \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}\right), \\ \tilde{\rho}^*\left(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}\right), \tilde{\rho}^*\left(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_n}^{i_n}\right) \end{array} \right]}$$

$$\begin{aligned} & \left[ \max \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \right]^{\bar{p}} \\ & \leq \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}}}{\xi \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1} \right\}} \end{aligned}$$

Since  $\xi$  satisfies the given condition, thus

$$\xi \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1} \right\} = \bar{1}$$

and thus

$$\tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq M^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}}$$

where  $M = \max \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}$

Now, two cases will be arised:

CASE 1. If  $M = \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right)$ , then

$$\begin{aligned} & \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{q}} \\ & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1} - \bar{p} - \bar{q}} \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{q}} \\ & \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\bar{q}}{\bar{1} - \bar{p} - \bar{q}}} \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{q}}{\bar{1} - \bar{p} - \bar{q}} \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\bar{\eta}^2} \\ & \quad \vdots \\ \Rightarrow & \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n}. \end{aligned}$$

CASE 2. If  $M = \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n})$ , then

$$\begin{aligned} \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{p}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\}^{\bar{q}} \\ \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\}^{1-\bar{q}} &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{p}+\bar{q}} \\ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\frac{\bar{p}+\bar{q}}{1-\bar{q}}} \\ \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n}) \right\}^{\bar{\eta}}, \quad \text{where } \bar{\eta} = \frac{\bar{p}+\bar{q}}{1-\bar{q}} \\ \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}) \right\}^{\bar{\eta}^2} \\ &\vdots \\ \Rightarrow \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n}. \end{aligned}$$

Using both the cases, for any  $m > n$ , where  $m, n \in \mathbb{N}$ , we have

$$\begin{aligned} \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}) \right\} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_{n+3}}^{i_{n+3}}) \right\} \\ &\quad \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m}) \right\} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+1}} \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n+2}} \\ &\quad \dots \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{m-1}} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\bar{\eta}^{n(\bar{1}+\bar{\eta}+2\bar{\eta}^2+\dots+\bar{\eta}^{m-n-1})}} \\ &\lesssim \left\{ \tilde{\rho}^*(\tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1}) \right\}^{\frac{\bar{\eta}^n}{1-\bar{\eta}}}. \end{aligned}$$

Since  $\bar{p} + 2\bar{q} \lesssim \bar{1}$ , therefore  $\bar{\eta} \lesssim \bar{1}$  and hence  $\tilde{\rho}^*(\tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m}) \rightarrow \bar{1}$  as  $m, n \rightarrow \infty$ . So, the soft sequence  $\{\tilde{T}_{\alpha_n}^{i_n}\}$  is soft multiplicative Cauchy sequence in  $\tilde{I}$ . Being the completeness of  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ , there exists a soft point  $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$  such that  $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$  as  $n \rightarrow \infty$ .



$$\begin{aligned}
 \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}} &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i^*} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{p}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{p}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}+\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\rightarrow \bar{1} \quad \text{as } n \rightarrow \infty.
 \end{aligned}$$

Since  $\bar{p} + \bar{2}\bar{q} \lesssim \bar{1}$  and  $\bar{p} \gtrsim \bar{0}$ , therefore  $\bar{1} - \bar{q} \gtrsim \bar{0}$  which indicates that  $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$  and hence  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ .

Now, assume  $M_2 = \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right)$ . Then, from (3.1), we have

$$\begin{aligned}
 \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}} &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i^*} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\stackrel{\leq}{\approx} \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{p}+\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\
 &\rightarrow \bar{1} \quad \text{as } n \rightarrow \infty.
 \end{aligned}$$

Since  $\bar{p} + \bar{2}\bar{q} \lesssim \bar{1}$  and  $\bar{p} \gtrsim \bar{0}$ , therefore  $\bar{1} - \bar{q} \gtrsim \bar{0}$  which indicates that  $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$  and hence  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ .

Now, if  $M_2 = \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}$ . Then, from (3.1), we have

$$\begin{aligned} \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}} &\lesssim \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{1}-\bar{q}-\bar{p}} &\lesssim \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{\bar{1}+\bar{q}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha^*}^{i^*} \right) \right]} \\ &\left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha_n}^{i_n}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right] \end{aligned}$$

$\rightarrow \bar{1}$  as  $n \rightarrow \infty$ .

Since  $\bar{p} + 2\bar{q} < \bar{1}$ , therefore  $\bar{1} - \bar{q} - \bar{p} > \bar{0}$  which indicates that  $\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$  and hence  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ .

Thus, in all the cases, we get  $\tilde{T}_{\alpha^*}^{i^*}$  as a “soft fixed point” of  $(h, \psi)$ . Now, if  $\tilde{T}_{\alpha'}^{i'}$  be another “soft fixed point” of  $(h, \psi)$ . Then,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\} \\ &\left[ \max \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right] \right]^{\bar{p}} \\ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &\leq \frac{\left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\bar{q}}}{\xi \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho} \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \right.} \\ &\left. \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]} \\ &\left[ \max \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho} \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\} \right]^{\bar{p}} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho} \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \right.} \\ &\left. \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}} \\ &\leq \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}}}{\xi \left\{ \bar{1}, \bar{1} \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right), \right.} \\ &\left. \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}} \\ &\leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}+2\bar{q}} \\ \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1}-\bar{p}-2\bar{q}} &\leq \bar{1} \end{aligned}$$

Since  $\bar{p} + 2\bar{q} \lesssim \bar{1}$ , therefore  $\bar{1} - \bar{p} - 2\bar{q} \gtrsim \bar{0}$  Thus,  $\tilde{\rho}^* (\tilde{T}_{\alpha^*}^{i^*}, T_{\alpha^*}^{i'}) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^* = \tilde{T}_{\alpha^*}^{i'}$ .

Hence, there is one and only one soft fixed point of  $(h, \psi)$ .

**Theorem 3.4.** Let  $(\tilde{I}, \tilde{\rho}^*, \Omega)$  be a complete soft multiplicative metric space and  $(h, \psi) : (\tilde{I}, \tilde{\rho}^*, \Omega) \rightarrow (\tilde{I}, \tilde{\rho}^*, \Omega)$  be a mapping, which satisfies the soft multiplicative generalized weak contractive mapping

$$\tilde{\rho}^* \left\{ (h, \psi) (\tilde{T}_{\alpha}^i), (h, \psi) (\tilde{T}_{\beta}^j) \right\} \leq \frac{\left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\beta}^j) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\alpha}^i) \right\} \right]^{\frac{\bar{p}}{2}} \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\beta}^j) \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\alpha}^i) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* (\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j) \right\}^{\bar{r}}}{\xi \left[ \begin{array}{l} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\alpha}^i) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\beta}^j) \right\}, \\ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha}^i, (h, \psi) (\tilde{T}_{\beta}^j) \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\beta}^j, (h, \psi) (\tilde{T}_{\alpha}^i) \right\}, \tilde{\rho}^* (\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j) \end{array} \right]} \quad \forall \tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j \in SP(\tilde{I}),$$

where  $\bar{p}, \bar{q}$  and  $\bar{r}$  are non-negative soft real numbers such that  $\bar{p} + \bar{q} + \bar{r} \lesssim \bar{1}$  and  $\xi : [\bar{1}, \infty)^2 \rightarrow [\bar{1}, \infty)$  is a continuous function such that  $\xi (\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l) = \bar{1}$  iff one of  $\tilde{T}_{\alpha}^i, \tilde{T}_{\beta}^j, \tilde{T}_{\gamma}^k, \tilde{T}_{\delta}^l = \bar{1}$ . Then, there exists a unique fixed point of  $(h, \psi)$ .

Proof. Let  $\tilde{T}_{\alpha_0}^{i_0}$  be any soft point in  $SP(\tilde{I})$ . Fix

$$\begin{aligned} \tilde{T}_{\alpha_1}^{i_1} &= (h, \psi) \tilde{T}_{\alpha_0}^{i_0} \\ \tilde{T}_{\alpha_2}^{i_2} &= (h, \psi) \tilde{T}_{\alpha_1}^{i_1} \\ &\vdots \\ \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} &= (h, \psi) \tilde{T}_{\alpha_n}^{i_n} \end{aligned}$$



$$\xi \left[ \begin{array}{l} \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \\ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right), \bar{1}, \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \end{array} \right] = \bar{1}.$$

Thus, we have

$$\begin{aligned} & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\}^{1 - \frac{\bar{p}}{2} - \frac{\bar{q}}{2}} \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{r}} \\ \Rightarrow & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^{\frac{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{r}}{1 - \frac{\bar{p}}{2} - \frac{\bar{q}}{2}}} \\ \Rightarrow & \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-1}}^{i_{n-1}}, \tilde{T}_{\alpha_n}^{i_n} \right) \right\}^\eta, \quad \eta = \frac{\frac{\bar{p}}{2} + \frac{\bar{q}}{2} + \bar{r}}{1 - \frac{\bar{p}}{2} - \frac{\bar{q}}{2}} \\ & \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n-2}}^{i_{n-2}}, \tilde{T}_{\alpha_{n-1}}^{i_{n-1}} \right) \right\}^{\eta^2} \\ & \vdots \\ & \leq \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\eta^n}. \end{aligned}$$

For any  $m > n$ , where  $m, n \in \mathbb{N}$

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+2}}^{i_{n+2}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_{n+1}}^{i_{n+1}} \right) \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{n+1}}^{i_{n+1}}, \tilde{T}_{\alpha_{n+2}}^{i_{n+2}} \right) \right\} \dots \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_{m-1}}^{i_{m-1}}, \tilde{T}_{\alpha_m}^{i_m} \right) \right\} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^n} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n+1}} \dots \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{m-1}} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\bar{\eta}^{n \left( \bar{1} + \bar{\eta} + \bar{\eta}^2 + \dots + \bar{\eta}^{m-n-1} \right)}} \\ & \lesssim \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha_0}^{i_0}, \tilde{T}_{\alpha_1}^{i_1} \right) \right\}^{\frac{\bar{\eta}^n}{\bar{1} - \bar{\eta}}}. \end{aligned}$$

Since  $\bar{p} + \bar{q} + \bar{r} < \bar{1}$ , therefore  $\bar{\eta} < \bar{1}$  which indicates that  $\tilde{\rho}^* \left( \tilde{T}_{\alpha_n}^{i_n}, \tilde{T}_{\alpha_m}^{i_m} \right) \rightarrow \bar{1}$  as  $m, n \rightarrow \infty$ . So, the soft sequence  $\{\tilde{T}_{\alpha_n}^{i_n}\}$  is soft multiplicative Cauchy sequence in  $\tilde{I}$ . Being the completeness of  $(\tilde{I}, \tilde{\rho}^*, \Omega)$ , there exists a soft point  $\tilde{T}_{\alpha^*}^{i^*} \in \tilde{I}$  such that  $\tilde{T}_{\alpha_n}^{i_n} \rightarrow \tilde{T}_{\alpha^*}^{i^*}$  as  $n \rightarrow \infty$ .



$\tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) \right\} = \bar{1}$  as  $n \rightarrow \infty$ . This shows that  $(h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right) = \tilde{T}_{\alpha^*}^{i^*}$  and hence  $\tilde{T}_{\alpha^*}^{i^*}$  is a “soft fixed point” of  $(h, \psi)$ .

Now, if  $\tilde{T}_{\alpha'}^{i'}$  be another “soft fixed point” of  $(h, \psi)$ . Then,

$$\begin{aligned} \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) &= \tilde{\rho}^* \left\{ (h, \psi) \left( \tilde{T}_{\alpha^*}^{i^*} \right), (h, \psi) \left( \tilde{T}_{\alpha'}^{i'} \right) \right\} \\ &= \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right\} \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\} \right]^{\frac{\bar{p}}{2}} \\ &\leq \frac{\left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right) \right\} \right]^{\frac{\bar{q}}{2}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right\}, \tilde{\rho}^* \left\{ \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\}, \right. \\ &\quad \left. \tilde{\rho}^* \left\{ \tilde{T}_{\alpha^*}^{i^*}, (h, \psi) \tilde{T}_{\alpha'}^{i'} \right\}, \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, (h, \psi) \tilde{T}_{\alpha^*}^{i^*} \right) \right\}, \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right]} \\ &\quad \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{p}} \left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right) \right\}^{\bar{q}} \\ &= \frac{\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{r}}}{\xi \left[ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha'}^{i'} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \tilde{\rho}^* \left( \tilde{T}_{\alpha'}^{i'}, \tilde{T}_{\alpha^*}^{i^*} \right), \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right]} \\ &\left\{ \tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) \right\}^{\bar{1} - \bar{q} - \bar{r}} \lesssim \bar{1} \end{aligned}$$

Since  $\bar{p} + \bar{q} + \bar{r} \lesssim \bar{1}$  and  $\bar{p} > \bar{0}$ , therefore  $\bar{1} - \bar{q} - \bar{r} \gtrsim \bar{0}$  and hence

$$\tilde{\rho}^* \left( \tilde{T}_{\alpha^*}^{i^*}, \tilde{T}_{\alpha'}^{i'} \right) = \bar{1} \Rightarrow \tilde{T}_{\alpha^*}^{i^*} = \tilde{T}_{\alpha'}^{i'}$$

Hence, there is one and only one soft fixed point of  $(h, \psi)$ .

## REFERENCES

- [1] A.E. Bashirov, E.M. Kurpirnar and A. Ozgapiri (2008). Multiplicative calculus and its applications. *Journal of Mathematical Analysis and Applications*, 337, 36-48.
- [2] B.R. Wadkar, L.N. Mishra, R. Bhardwaj and B. Singh (2016), Fixed Point Results Related to Soft Sets, *Australian Journal of Basic and Applied Sciences*, 10(16), 128-137.
- [3] D. Wardowski(2013), On a soft mapping and its fixed points, *Fixed Point Theory and Applications*, 2013:182, 1-11.
- [4] D.A. Molodtsov(1999), Soft set theory-first result, *Computers and Mathematics with Applications*,37, 19-31.
- [5] H. Hosseinzadeh(2017), Fixed point theorem on soft metric spaces, *Journal of Fixed Point Theory and Applications*, 18(4), 1-22.
- [6] K. Abodayeh, A. Pitea, W. Shatanavi and T. Abdegawad(2015). Remarks on multiplicative metric spaces and related fixed points. arXiv:1512.03771v1, 1-7.
- [7] M. Abbas, G. Murtaza(2016), On the fixed point theory of soft metric spaces, *Fixed Point Theory and Applications* ,17, 1-11.
- [8] M.I. Ali, F. Feng, X. Liu, W.K. Min and M. Shabir(2011), On some new operation in soft set theory, *Computer and Mathematics with Applications* ,62, 351-358.
- [9] M. Ozavsar and A.C. Cevikel (2012). Fixed Point of multiplicative contraction mappings on multiplicative metric space. arXiv:1205.5131v1,1-14.
- [10]M.I. Yazar, C. Gunduz and S. Bayramov(2016), Fixed point theorems of soft contractive mappings, *Filomat*, 30, 269-279.
- [11] P.K. Maji, A.R. Roy and R. Biswas(2002), An application of soft real number in a decision making problem, *Computers and Mathematics with Applications*, 42,1077-1083.
- [12]P.K. Maji, R. Biswas and A.R. Roy(2003), Soft set theory, *Computers and Mathematics with Applications* ,45,555-562.
- [13]R. Bhardwaj, S. Singh, S. Pandey and B.R. Wadkar(2021), Fixed point results in soft metric space, *International Journal of Modern Agriculture*, 10(2), 1792-1803.
- [14]R. P. Agrawal, E. Karapinar, B. Samet(2016), An essential remark on fixed point results on multiplicative metric spaces, *Fixed Point Theory and Applications*,21, 1-3.
- [15] S. Das and S.K. Samanta(2013), On soft metric spaces, *Journal of Fuzzy Mathematics*, 21, 207-213.

- [16] S. Das and S.K. Samanta(2012), Soft real sets, soft real numbers and their properties, *Journal of Fuzzy Mathematics*, 20, 551-576.
- [17] S. Rathee, R. Girdhar and K. Dhingra(2020), Fixed point theorems in soft multiplicative metric spaces, *Communications in Mathematics and Applications*, 11, 425-442.
- [18] S. Solanki, R. Bhardwaj, B.K. Singh and M. Sharma(2017), Fixed point theory of soft weak contraction on a complete soft metric space, *International Journal of Advanced Engineering Science and Technological Research*, 41(1), 425-432.
- [19] T. Dosenovic, M. Pastolache and S. Radenovic(2016). On multiplicative metric spaces: survey, *Fixed Point Theory and Applications*, 92, 1-17.
- [20] T. Dosenovic, M. Pastolache and S. Radenovic(2017). Some essential remark on fixed point results on multiplicative metric space. *Journal of Advanced Mathematical Studies*, 10(1), 20-24.
- [21] X. He, M. Song and D. Chen (2014). Common fixed points for weak commutative mappings on a multiplicative metric space. *Fixed Point Theory and Applications*, 48, 1-9.