

## Original Research Article

# **Musculoskeletal Ultrasound in Rheumatological Diseases**

### **Abstract:**

Rheumatology has seen a sonographic revolution in recent years. Ultrasonography (US) has been demonstrated to be capable of significantly altering the traditional approach to most clinical problems in daily rheumatologic practice due to its ability to provide rapid, safe, and inexpensive access to anatomical information on the early targets of the majority of rheumatic diseases. Ultrasonography completes the physical examination with speed and accuracy and enables the detection of a broad range of abnormal findings involving various musculoskeletal (MSK) anatomical components.

Ultrasonography is mostly used for interventional guidance (joint aspiration, synovial or soft tissue biopsy, joint or tendon sheath injection). When fluid collections are very tiny or when the inflammatory process is next to anatomical structures that might be severely harmed by the injection, sonographic guiding is very effective.

MSK US is a significant imaging modality for practicing rheumatologists and provides an effective diagnostic tool for assessing patients with MSK problems. In the United States, the usage of MSK ultrasonography has evolved due to the availability of less priced, portable US equipment that give high-quality grayscale and power Doppler signals. The American College of Rheumatology (ACR) has given standardized training courses and a certification program for practicing

rheumatologists, and similar training is also offered through a number of fellowship programs.

In certain clinical circumstances, rheumatoid arthritis (RA) might be difficult to diagnose since crystalline arthropathies can closely resemble RA. For instance, palindromic RA might appear with articular symptoms that, based on history and clinical characteristics, mimic a crystalline disease. This study sought to investigate the relevance of MSK ultrasonography in rheumatologic illness diagnosis.

**Keywords:** Musculoskeletal Ultrasound, Rheumatological Diseases, Ultrasonography, Rheumatoid Arthritis

## Introduction

Rheumatology is an intriguing field of study. The diagnosis relies heavily on the medical history and physical examination. For the majority of rheumatologic disorders, viable therapeutic options are available once a diagnosis has been made. Can ultrasonography (US) enhance our clinical abilities while examining a patient with rheumatic illness suspicion? Indeed, it can! Combining a patient's medical history and physical examination with ultrasonography helps the rheumatologist to characterize rheumatic disease entities with precision. The rheumatologist can perform and interpret an US while questioning and evaluating the patient at the point of service. The additional benefit of US with clinical correlation expedites the diagnosis, management, and targeted therapy (e.g., US-guided injections) of rheumatologic disorders, with the potential for improved clinical outcomes and lower healthcare costs (1).

As rheumatology encompasses a broad spectrum of disorders, ultrasonography can be utilized for a growing variety of indications. The American College of Rheumatology has produced a study on the acceptable use of MSKUS in rheumatology clinical practice that gives an overview of clinical situations with favorable evidence-based recommendations based on current literature (2).

US of the MSK system has become an established imaging method for the diagnosis and follow-up of patients with rheumatic disorders during the past decade.

1-5 This is a result of technology advancements that have led to faster processors and higher frequency transducers. US is most frequently used to evaluate soft tissue disorders or identify fluid collection, but it may also be used to examine other tissues, including cartilage and bone surfaces (3).

Due to the superior axial and lateral resolution of US, even the smallest bone surface anomalies may be seen. Consequently, damaging and/or reparative/hypertrophic alterations on the bone surface might be detected prior to their visibility on x-rays or even MRI. However, US frequencies cannot penetrate bone, therefore intra-articular disease imaging is typically not achievable. The "real time" capabilities of US allow for the dynamic evaluation of joint and tendon motions, which can frequently help in the discovery of structural problems. The non-invasive nature, mobility, relative affordability, absence of ionizing radiation, and repeatability of US make it particularly valuable for monitoring therapy. US can also be utilized for aspiration, biopsy, and injection therapy guidance (4).

The majority of MSK work is accomplished using "grey scale," which produces black-and-white pictures; each white dot represents a reflected sound wave. Sound waves travel similarly to light waves, thus the denser a substance, such as bone cortex, the more reflecting it is and the whiter it will seem on the screen. Water is the least reflecting body substance; therefore, sound waves go directly through it and look black (5).

Color and power Doppler imaging, which offer color maps of tissue, are among the newest US methods now being researched. Here, the quantity of color is proportional to the amount of blood flow, which may be useful for assessing vascular tissues, such as those affected by soft tissue inflammation. To boost the sensitivity of power Doppler even further, intravenous bubble contrast agents are currently being developed (6).

## **Characteristics of MSK US**

Joint US is an accessible, simple, fast, and economical imaging method used to check MSK tissues, such as the tendons, ligaments, bursae, and articular surface. It does provide accurate images of structural abnormalities and a pathology caused by inflammation or degeneration conditions in a variety of rheumatologic diseases, including RA, osteoarthritis, SpA, and septic arthritis, and connective tissue diseases, including systemic lupus erythematosus (SLE), systemic sclerosis, inflammatory myositis, Sjogren's syndrome, and systemic vasculitis (7).

MSK US and MRI are among the new imaging modalities for rheumatic disorders with distinct properties. The benefits of MSK US over other imaging techniques include visibility of the joint cavity and soft tissue involvement, cost-effectiveness, availability, lack of radiation, multiplanar imaging capacity, and real-time dynamic evaluation of joints. MSK US is also a valuable instrument for directing invasive surgeries. It allows for accurate and exact needle placement and is a safe technique for patients requiring synovial fluid aspiration, injectable treatment, and/or biopsy (8). [12–14, 16]

MRI is a more sensitive imaging method for viewing inflammation and damage and is especially beneficial for identifying bone inflammation; nevertheless, it is costly, not easily accessible, time-consuming, and restricted to a small anatomical area each examination (9).

## **Limitations of MSK US**

The drawbacks of the US are its narrow field of view compared with MRI, and the fact that the US beam does not penetrate under the bone. Moreover, the US is highly operator dependent.

The sonographer should be aware of a variety of artifacts (Power Doppler artifacts and anisotropy), so as not to falsely recognize pathology where there is none. In addition, one must be aware not to over interpret slight US abnormalities that are observed by “sensitive” US machines and may not reflect disease (10).

## **Examination technique**

Continuous technological advancements in the United States have resulted in the availability of exceptionally high-quality sonographic equipment, enabling the histological examination of superficial tissues (tendons, tiny joints, skin, peripheral nerves, and blood vessels) (11).

The term 'acoustic microscopy' is appropriate for characterizing sonographic exams using probes with a very high frequency (0.16 MHz) and a resolution power of less than 0.1 mm. If having a high-quality US equipment is a must for MSK US, then the driver's expertise is even more crucial. US is a 'technical art' and is without a doubt the most 'user dependent' diagnostic imaging modality. In investigations of individuals with rheumatoid arthritis, a high level of interobserver concordance was

documented for the evaluation of synovitis and bone erosions in the minor joints of the hands and feet (12).

Even if the selection of the acoustic window is determined by the unique diagnostic necessity, a standardized method of examining the various anatomical locations is recommended. Each anatomical region should be investigated on many scan planes in order to collect all relevant data. An in-depth understanding of anatomy is necessary for MSKUS (13).

Sonographic follow-up requires the acquisition of similar images of the same anatomical region at different time intervals. Even minute differences in the control parameters of the sonographic beam, such as location, applied pressure, and beam tilt, might impact the sonographic picture, hence influencing its interpretation (14).

Sonographic palpation may be of tremendous use in the investigation of regional pain syndromes if it can elicit tenderness in particular anatomical regions. Three-dimensional MSKUS is an interesting new subject in US. The capture of the whole volumes of echoes generated by the probe permits a three-dimensional reconstruction of the studied area on an infinite number of scan planes, therefore enabling virtual MSK endoscopy. This approach is highly promising for evaluating entheses, tendons, and tiny joints(15).

### **Technical equipment**

MSK work need high-quality, high-resolution instruments. The choice of transducer will rely on the exams that will likely be conducted. High frequency linear transducers (7.5–20 MHz) are optimal for displaying surface structures such as tendons, ligaments, and tiny joints, but low frequency linear transducers (3.5–5 MHz) are occasionally more appropriate for bigger or deeper-sited joints such as the shoulder or hip (16).

In the United States, there is a continual trade-off between picture resolution and sound wave penetration depth. Higher frequency transducers offer superior spatial resolution, but their depth of penetration is less than that of lower frequency transducers. The size of the footprint (the transducer's contact area with the skin) is also a significant examination procedure component (17).

For instance, transducers with a big footprint cannot be maneuvered well enough to visualize fully tiny joints such as the metacarpophalangeal joints. However, these are merely basic factors; the most important factor is the total image resolution, which must be thoroughly analyzed and compared prior to purchase. For practical reasons, it is recommended to test whether a specific piece of sonographic equipment is capable of displaying the fine definition of small structures, such as the insertion of a small extensor tendon of the finger or the minute amount of fluid normally detectable in the pre-Achilles bursa (18).

The practical utility of color Doppler/power Doppler capabilities is currently being evaluated, particularly in light of the added expense. The purpose of color/power Doppler is to identify enhanced perfusion of soft tissue. Additionally, the prospective applications of three-dimensional US are also being evaluated. Finally, emphasis must be paid to picture documenting techniques. Every inspection should be meticulously documented. Images can be stored on paper, film, video cassettes, laser-printed x-ray acetates, optical discs, and digital storage systems. To increase the repeatability of these discoveries, all shown structures should be recorded in a standardized manner. In two perpendicular planes, pathological findings should be reported (4).

## **The development of rheumatological ultrasound**

The 1978 demonstration of synovitis of the knee in rheumatoid arthritis (RA) likely marked the beginning of modern 'rheumatologicUS' Advances in US technology have enabled high-resolution imaging of MSK structures, and MSK US is now a commonly utilized imaging technique for the MSK system. MSK US is a great tool for rheumatologists due to its improved awareness of rheumatic disorders, multiplanar imaging of articular structures in real time, cheap operating costs, lack of radiation, and high patient acceptance. Whether MSK US should be conducted by radiologists, rheumatologists, or other MSK specialists, or by all three, remains a source of debate (19).

The increasing inclusion of US as a common extension of clinical examination in other clinical specialties, such as obstetrics, gynecology, and cardiology, suggested that MSK US would become a natural extension of MSK assessment conducted by rheumatologists. In Germany, rheumatologists pioneered the introduction of MSK US into clinical practice, and MSK US is now a routine component of the rheumatology training curriculum. Indeed, the use of MSK US by rheumatologists has made a significant contribution to the knowledge of the natural history of rheumatic disorders and to the improvement of clinical diagnostic and therapeutic abilities (20).

However, the use of MSK US by rheumatologists abroad varies significantly. MSK US is a standard element of rheumatology training in Italy and Germany, but it remains mostly the domain of radiologists in the United States and the United Kingdom, despite a worldwide interest among rheumatologists. The complicated reasons for this include money, equipment, standardization, training, and the conventional views of rheumatologists toward MSK US (21).

### **Three- and four-dimensional ultrasonography**

USlagged behind CT and MRI in terms of 3D imaging until recently. Now, breakthroughs in computation have enabled 3D US imaging (termed 4D as it has the advantage of being performed in real time). All of these cutting-edge technologies are now being studied and have the potential to remove MSK US from the shadow of MRI in the evaluation of MSK disorders. In a matter of seconds, 3D sonographic images of a target region may be obtained, enabling an off-line, exhilarating virtual anatomy tour in longitudinal, transverse, and coronal planes, as well as simultaneous 3D reconstruction. After the workstation has been appropriately configured, an expert sonographer and a beginner will capture identical images. In addition, 3D US has the potential to significantly shorten the length of MSK US examinations while simultaneously improving the quality and consistency of the approach (22).

Three-dimensional images give a magnificent and in-depth view of the tiny joints, are easier to comprehend, and vividly highlight a number of anatomical aspects that cannot be seen by conventional US. In rheumatology, the primary clinical indications for 3D MSK US may include the early diagnosis of bone erosions in tiny joints and a more thorough evaluation of enthesitis and partial tendon tears. Although the influence of 3D US on final diagnosis or therapeutic monitoring has not yet been determined, it appears fair to anticipate that as technology developments continue to enhance picture quality, the use of 3D sonography in the diagnosis of MSK problems will increase rapidly (23).

The proper selection of MSK US equipment is crucial for allowing the incorporation of MSK US into rheumatology practice and for facilitating MSK US training. When selecting sonographic equipment for MSK US,

cost, picture resolution and quality, transducer design, equipment size/portability, and color/power Doppler options are crucial factors (24).

The cost of US equipment is directly proportional to picture quality and resolution. A high-resolution system with enhanced imaging definition generates high-quality, interpretable pictures. A high-quality, high-resolution system might cost anywhere from £30,000 to £130,000. By configuring the system exclusively for MSK imaging, significant cost savings can be realized. Small, transportable systems are less costly (£15,000), provide pictures of adequate quality, and are simple to operate at the bedside. However, pictures on a smaller screen are more difficult to interpret for less experienced ultrasonographers. The ideal combination, in the opinion of the authors, is to have access to a larger, high-resolution system for training and research purposes, as well as for "difficult cases," while a smaller, portable system is sufficient for the majority of routine clinical practice performed by the experienced sonographer (25).

Thirteen out of fifteen studies in our comprehensive assessment of the literature found that MSK US is useful for diagnosing RA when combined with clinical examination and laboratory evaluation. One research found greater sensitivity in identifying arthritis at the joint level, but no meaningful value was identified at the patient level (26).

Another research demonstrated no predictive value for RA when MSK US was included as a standard component of early arthritis evaluation. In terms of population size, these two studies are the greatest; the latter contained 379 participants. There are, however, limits to the MSK US performance in this investigation. The MSK US process lasted ten minutes, and only sixteen joints were checked. Additionally, wrists and MTP 3–5 were excluded. This might be a reason for the bad results, as RA frequently

causes wrist inflammation. MSK US is deemed beneficial in each of the remaining 13 trials included in this review, and wrist examinations are performed in each (27).

## Conclusions

MUSU is quick, simple to apply, and readily available, no ionizing radiation, it is a low-cost procedure, well tolerated by patients, of better sensitivity than physical examination, allows evaluation of many structures over a relatively short period of time, enables dynamic examination during movement of joints and tendons which can be helpful to assess impingement syndrome of the shoulder for example and detects subclinical synovitis, enthesitis, early erosions.

## References

1. Brühlmann P, Michel BA. [History and clinical examination in rheumatology--key to diagnosis and prognosis]. *Ther Umsch.* 2006;63(7):485-90.
2. McAlindon T, Kissin E, Nazarian L, Ranganath V, Prakash S, Taylor M, et al. American College of Rheumatology report on reasonable use of musculoskeletal ultrasonography in rheumatology clinical practice. *Arthritis Care Res (Hoboken).* 2012;64(11):1625-40.
3. Grassi W, Lamanna G, Farina A, Cervini C. Sonographic imaging of normal and osteoarthritic cartilage. *Semin Arthritis Rheum.* 1999;28(6):398-403.
4. Backhaus M, Burmester G-R, Gerber T, Grassi W, Machold KP, Swen WA, et al. Guidelines for musculoskeletal ultrasound in rheumatology. *Annals of the Rheumatic Diseases.* 2001;60(7):641-9.

5. Tok F, Demirkaya E, Özçakar L. Musculoskeletal ultrasound in pediatric rheumatology. *Pediatric Rheumatology*. 2011;9(1):25.
6. Cutolo M, Smith V. Detection of microvascular changes in systemic sclerosis and other rheumatic diseases. *Nature Reviews Rheumatology*. 2021;17(11):665-77.
7. Kaeley GS, Bakewell C, Deodhar A. The importance of ultrasound in identifying and differentiating patients with early inflammatory arthritis: a narrative review. *Arthritis Res Ther*. 2020;22(1):1.
8. Kondo Y, Suzuki K, Inoue Y, Sakata K, Takahashi C, Kaneko Y, et al. Safety and tolerability of ultrasound-guided synovial needle biopsy in Japanese arthritis patients. *Mod Rheumatol*. 2021;31(5):960-5.
9. MacRitchie N, Frleta-Gilchrist M, Sugiyama A, Lawton T, McInnes IB, Maffia P. Molecular imaging of inflammation - Current and emerging technologies for diagnosis and treatment. *Pharmacol Ther*. 2020;211:107550.
10. D'Agostino MA, Terslev L, Wakefield R, Østergaard M, Balint P, Naredo E, et al. Novel algorithms for the pragmatic use of ultrasound in the management of patients with rheumatoid arthritis: from diagnosis to remission. *Ann Rheum Dis*. 2016;75(11):1902-8.
11. Baldini M, Giannoni S, Cioffi E, Grifoni E, Bertini E. Ultrasound-guided diagnosis of an uncommon stroke: A case of giant cell arteritis. *J Clin Images Med Case Rep*. 2021;2(3):1173.
12. Sambri A, Spinnato P, Tedeschi S, Zamparini E, Fiore M, Zucchini R, et al. Bone and joint infections: The role of imaging in tailoring diagnosis to improve patients' care. *Journal of Personalized Medicine*. 2021;11(12):1317.
13. Enache L, Mogosan C, Popescu C, Gutoiu G, Codreanu C. SAT0644 Is ankle involvement important in rheumatoid arthritis? : BMJ Publishing Group Ltd; 2017.

14. Grassi W, Filippucci E, Busilacchi P. Musculoskeletal ultrasound. *Best Practice & Research Clinical Rheumatology*. 2004;18(6):813-26.
15. Lowe C, Ghulam Q, Bredahl K, Rogers S, Ghosh J, Sillesen H, et al. Three-dimensional Ultrasound in the Management of abdominal aortic aneurysms: a topical review. *European Journal of Vascular and Endovascular Surgery*. 2016;52(4):466-74.
16. McPhillips R. *Fabrication of Ultrasound Transducers and Arrays Integrated within Needles for Imaging Guidance and Diagnosis*: University of Dundee; 2017.
17. Shung K, Cannata J, Qifa Zhou M, Lee J. High frequency ultrasound: a new frontier for ultrasound. *Annu Int Conf IEEE Eng Med Biol Soc*. 2009;2009:1953-5.
18. Allen GM, Wilson DJ. *Ultrasound Guided Musculoskeletal Injections E-Book*: Elsevier Health Sciences; 2017.
19. Kane D, Balint PV, Sturrock R, Grassi W. Musculoskeletal ultrasound—a state of the art review in rheumatology. Part 1: Current controversies and issues in the development of musculoskeletal ultrasound in rheumatology. *Rheumatology*. 2004;43(7):823-8.
20. Grassi W. Clinical evaluation versus ultrasonography: who is the winner? *J Rheumatol*. 2003;30(5):908-9.
21. Wakefield RJ, Goh E, Conaghan PG, Karim Z, Emery P. Musculoskeletal ultrasonography in Europe: results of a rheumatologist-based survey at a EULAR meeting. *Rheumatology (Oxford)*. 2003;42(10):1251-3.
22. Lees W. Ultrasound imaging in three and four dimensions. *Semin Ultrasound CT MR*. 2001;22(1):85-105.

23. Jockenhövel J. A Three-Dimensional Checkerboard: The Long Take in 3D Films. *Die ästhetisch-narrativen Dimensionen des 3D-Films*: Springer; 2016. p. 55-69.
24. Iagnocco A, Porta F, Cuomo G, Delle Sedie A, Filippucci E, Grassi W, et al. The Italian MSK US Study Group recommendations for the format and content of the report and documentation in musculoskeletal ultrasonography in rheumatology. *Rheumatology (Oxford)*. 2014;53(2):367-73.
25. Lucas T, Quidu I, Bridal SL, Gateau J. High-contrast and-resolution 3-D ultrasonography with a clinical linear transducer array scanned in a rotate-translate geometry. *Applied Sciences*. 2021;11(2):493.
26. Lage-Hansen PR, Lindegaard H, Chrysidis S, Terslev L. The role of ultrasound in diagnosing rheumatoid arthritis, what do we know? An updated review. *Rheumatology International*. 2017;37(2):179-87.
27. Pratt AG, Lorenzi AR, Wilson G, Platt PN, Isaacs JD. Predicting persistent inflammatory arthritis amongst early arthritis clinic patients in the UK: is musculoskeletal ultrasound required? *Arthritis Research & Therapy*. 2013;15:R118 - R.