

## Short Research Article

### An association between CT SCORE and the severity of COVID-19

#### **ABSTRACT**

**Introduction-** COVID-19 has changed the world fabric. While many still haven't recovered from the grief of losing their loved ones; the virus continues to mutate leaving the world to deal with a social and economic pandemic. In this study we aim at determining an association between CT SCORE and the severity of COVID-19 thereby understanding the prognosis/severity of disease in patients.

**Method-** A prevalidated google questionnaire was used to obtain data. The study population consisted of people who tested RT-PCR COVID-19 positive.

**Results-** A statistically significant correlation was seen between CT score and outcome of disease.

**Conclusion-** Routine CT scan of suspected COVID-19 cases gives early diagnosis; it could also guide in hastening the treatment received by the patient and improve overall prognosis be used as a guide for the management of the patients and hence improve overall prognosis.

#### **INTRODUCTION**

The Coronavirus disease 2019 (COVID-19), was declared a global pandemic by World Health Organization on March 11, 2020. Since then, the virus has known no bounds as it swept rapidly across major continents causing an alarming rise in the number of cases and overwhelming the health care system of many countries.

Chest CT has proven to be a useful supplement to RT-PCR and has been shown to have high sensitivity to diagnose the disease. Apart from being a diagnostic tool, CT can also potentially help in prognostication and evaluation of disease progression.

The CT severity score index is a scoring system used to assess the lung changes and involvement by COVID-19 based on approximate estimation of pulmonary involved areas. Each of the five lung lobes has been visually scored from 1 to 5:

Score 1: <5% lobar involvement.

Score 2: 5–25% lobar involvement.

Score 3: 26–50% lobar involvement.

Score 4: 51–75% lobar involvement.

Score 5: > 75% lobar involvement.

The final score is calculated by the summation of the individual lobar scores. The maximum (total) lobar score is 25. The total lung involvement is obtained by multiplying the total score by 4.<sup>1</sup>

Mounting evidence from across the globe seems to suggest that CT score could play a role in determining the severity of disease in an individual due to COVID-19.<sup>1</sup>

CTs conveniently provide diagnosis in Emergency Medicine because the results are obtained immediately, and alternate diagnoses can be excluded. In addition, using a CT severity score has shown to correlate with disease severity, and it can be used as a prognostic marker.<sup>2, 3, 4, 5</sup>

Rapid and accurate diagnosis is essential to profile patients and utilize scarce resources adequately. However, this is hampered by limited sensitivity of RT-PCR.<sup>6, 7</sup> Although real time RT-PCR is considered the gold standard for diagnosis of COVID-19 infection, time required to get the report of RT-PCR is a huge problem. CT has an advantage of immediate reporting. It has also shown to be of diagnostic importance in cases of false negative results of RT-PCR. Besides being a diagnostic tool, it also has an immense role in monitoring disease progression and evaluating therapeutic efficacy.<sup>8</sup>

The precise patients' categorization and radiological severity scoring are essential for appropriate management, especially in mild cases before patient deterioration. Chest X-ray is seen to have very low sensitivity in early-stage disease, CT is therefore the primary imaging tool.<sup>9</sup>

Additionally, there is evidence that sensitivity of combined nasal and pharyngeal swabs may be inadequate if obtained at a single point of time, and also depends on the technical characteristics of the test and method of specimen collection.<sup>10, 11</sup> The relatively long turnaround time (TAT) for viral testing together with the low sensitivity of a single real-time reverse-transcriptase polymerase-chain reaction (RT-PCR) assay of nasal and pharyngeal swab specimens also implies that a large number of SARS-CoV-2 patients would not be quickly identified and may not be appropriately managed.

In retrospective studies, sensitivity of chest CT for COVID-19 is excellent, and it may even be greater than that of PCR.<sup>12</sup>

Whereas some studies also suggest that there is no correlation of CT score with severity of disease.<sup>13</sup>

Therefore, this study aims at determining, if any relationship exists between CT score and severity of disease.

## **METHODOLOGY**

### **Study design and data collection**

This is a Cross sectional, questionnaire-based study. The study was conducted between 30th April 2021 to 30th June 2021 after obtaining Ethics committee approval. A prevalidated google questionnaire was prepared and circulated to people through various social media platforms. Relatives of deceased individuals were asked to fill the questionnaire on behalf of them. Data collected from google questionnaire was assessed and correlation was established.

The study population consisted of people who tested RT-PCR COVID-19 positive. A sample size of 105 was obtained, of which 71 people tested positive for RT-PCR test and 42 of them underwent CT scan. Data was analyzed using Chi-square test and Fischer's test.

#### Inclusion criteria

- All individuals above 18 years, both genders.
- People who tested RT-PCR positive for COVID-19.

#### Exclusion criteria

- All individuals who did not give consent to participate in the study.

After eligibility to the inclusion and exclusion criteria, 71 patients were included in the study.

## RESULTS

**TABLE 1: DEMOGRAPHIC DATA**

The study comprised of 105 individuals of which 71 tested positive for COVID-19 and 42 of whom underwent CT scan. There was a total of 36 males (50.7%) and 35 females (49.3%) with varied ages. The age groups were categorised into 4 groups.i.e. <21 years, 21-40 years, 41-60 years and >60 years respectively.

	CT SCORE				Total CT Scan patients (N)	Total Covid-19 patients N(%)	Statistical Analysis	Recovered	Deceased
	1-10 n(%)		>10 n(%)						
<b>PATIENTS</b>	24 (57.1)		18 (42.8)		42	71 (100%)	Chi square=1, df=1, p=1, OR=1	65	6
<b>Age(Years)</b>									
<40	10 (66.6)		5 (33.3)		15	32			
>40	14 (51.8)		13 (48.1)		27	39 (43.7%)		34	5
	CT SCORE				Total CT Scan patients (N)	Total Covid-19 patients N(%)	Statistical Analysis	Recovered	Deceased
	1-5 n(%)	6-10 n(%)	11-15 n(%)	>15 n(%)					
<b>Gender</b>							df=3, fisher's exact test, p=0.732		
Male	7 (50%)	7 (70%)	8 (66.6%)	3 (50%)	25	36 (50.7%)		33	3
Female	7(50%)	3 (30%)	4 (33.3%)	3 (50%)	17	35 (49.3%)		32	3

**TABLE 2: CORRELATION BETWEEN CT SEVERITY SCORE AND TYPE OF TREATMENT REQUIRED**

CT SCORE	Home n(%)	Required hospitalisation without oxygen n(%)	Required hospitalisation with oxygen n(%)	Needed Ventilator/ICU n(%)	Statistical Analysis
<10	17	4	1	2	Chi square=6.763, p=0.009
>10	6	3	4	5	

Data of 42 patients who have undergone CT scan

Significant correlation can be seen between CT score and need for hospitalisation,  $p=0.009$ .

Out of 24 individuals with CT score less than 10, only 7 needed hospitalisation (29.16%) while 12 out of 18 individuals (66.67%) with CT score more than 10 required hospitalisation.

**TABLE 3: CORRELATION BETWEEN CT SEVERITY SCORE WITH OUTCOME OF PATIENT**

CT severity score n(%)	Patients n(%)	Recovered n(%)	Deceased n(%)	Statistical Analysis
<10	24	24	Nil	Fisher's exact test p=0.0035
>10	18	12	6	

Significant correlation can be seen between CT score and outcome of patient,  $p=0.0035$ . All patients recovered with CT score less than 10, whereas 6 out of 18 patients succumbed to the disease (Mortality 33.33%) with CT score of 10 and above.

**TABLE 4: CORRELATION BETWEEN TREATMENT REQUIRED WITH OUTCOME OF PATIENT**

Type of treatment received	Total patients N=71 n(%)	Recovered n(%)	Deceased n(%)	Statistical Analysis
Treated at Home	50	50	0	Fisher's exact test p=0.0001
Required hospitalisation without oxygen	9	8	1	
Required hospitalisation with oxygen	5	4	1	
Needed Ventilator/ICU	7	3	4	

Data of all 71 patients with RT-PCR COVID-19 positive test

Significant correlation can be seen between treatment required and outcome of the patient,  $p=0.0001$ . Recovery rate was 100% for patients treated at home. Whereas mortality rate was 66.66% for patients with ICU admissions.

**TABLE 5: CORRELATION BETWEEN CO-MORBIDITIES AND TYPE OF TREATMENT REQUIRED**

Comorbidity	Type of Treatment received				CT Scan performed	Recovered	Deceased	Total
	Home	Hospitalisation without oxygen	Hospitalisation with oxygen	Ventilator/ICU				
HTN+DM	5	0	0	5	9	7	3	10
HTN only	4	1	0	1	1	6	0	6
DM only	0	2	1	0	3	2	1	3
Obesity	3	2	1	0	6	6	0	6
Cancer	0	2	0	0	1	1	1	2
Asthma	2	0	0	0	0	2	0	2
No comorbidity	16	2	3	1	22	21	1	22
<b>Total</b>	30	9	5	7	42	45	6	51

Data of 29 patients with one or more co-morbid conditions of which 20 underwent CT Scan.

Significant correlation can be seen between presence of co-morbidity and mortality rate.

Mortality rate is 83.33% (5/6) in patients with one or more co-morbid condition.

Significant correlation can also be seen between presence of co-morbidity and ICU admission. Of the 7 ICU admissions, 6 patients were having one or more co-morbid conditions.

UNDER PEER REVIEW

## DISCUSSION

The WHO recommended the use of chest imaging as a part of diagnostic tool for COVID-19 disease in case RT-PCR testing is not available, test results are delayed or in patients suspected to have COVID-19 but test negative with RT-PCR.<sup>20</sup> Mounting evidence from across the globe seems to suggest that CT score could play a role in determining the severity of disease in an individual due to COVID-19.<sup>21, 22</sup>

Several existing literatures suggest that the more severe CT pulmonary changes (i.e., higher CT severity score) was significantly associated with Males.<sup>14,15</sup> But in our study, we did not find a statically significant correlation with gender. This could be particularly due to small sample size. The possible reason for more males being affected than females could be possibly due to higher circulating ACE2 receptor levels in men than women.<sup>19</sup> Hence, men are more susceptible to acquire the virus than women.

As seen in many existing literatures, elderly people experienced the brunt of the disease and succumbed to the virus more than any other age group.<sup>16</sup> Results of our study show that 48.1% (13/18) patients above 40 years have CT score more than 10. Whereas 33.3% (5/18) patients below 40 years CT score more than 10. The possible reason for increased susceptibility to COVID-19 could be due to increased prevalence of comorbidities with increasing age which weakens one's immune system, as well as due to upregulation of ACE-2 receptors in specific organs particularly in individuals with comorbidities.<sup>19</sup>

Results from our study show that the need for hospitalisation increased with higher CT scores. Table 2 shows that oxygen requirements also increased with the increasing CT severity score, consistent with past literature. The increased need for oxygen could be due to severe damage to multiple alveoli by the virus. In our study 22.22% (4/18) individuals with CT score more than 10 required oxygen therapy while only 4.54% (1/22) individuals with CT

score less than 10 required oxygen therapy. A statistically significant correlation was seen between presence of co-morbidity and ICU admission. As shown in Table 5, 85.71% (6/7) ICU admissions had one or more comorbidity. Also, individuals with comorbidities were seen to maximally constitute ICU admissions. It can be seen in table 5, that 5 deceased patients were from the ICU admissions and each one of them had one or the other comorbidity.

Presence of co-morbid condition is positively correlated with increasing CT severity score. 61.1% (11/18) patients having CT severity score more than 10, had one or more co-morbid condition.

Additionally, studies have shown that the presence of comorbidities particularly hypertension, diabetes, obesity, and cancer carry a poor prognosis in disease outcome, with even worse outcome when multiple comorbidities are present.<sup>17,18</sup> We can find a similar relation in our study, 5 out of 6 deceased patients were having one or more co-morbid condition.i.e., 3 patients had both HTN and DM (Table 5). As the CT severity score increases, mortality rate also increases, all 6 deceased patients had CT severity score more than 10 (Table 3)

## **CONCLUSION**

In conclusion, CT scans can play a pivotal role in assisting physicians in the management plan and is a valuable indicator for disease severity and outcome. In a pandemic, one needs a diagnostic test that provides immediate results to confirm the diagnosis. RT-PCR, though considered the gold standard has long waiting hours for testing and often, test results are delayed or take longer than usual. This delay in obtaining test results could possibly mean that an infected person would continue to infect others in the community and the disease would progress further unchecked. CT Scan if implemented, would provide timely results,

and hasten the process of treatment thereby halt the disease progression much before extensive damage to the immune system is caused by the virus. Also, CT score would provide information about severity of the disease and help in deciding further course of action. Findings of our study show a statistically significant correlation between CT score and severity of disease. Individuals with one or more comorbid conditions are more likely to have a higher CT Score and thereby carry poor prognosis. However, more research is needed to further clarify the value of chest CT for prognostication in COVID-19 disease, including correlation with patient outcome.

## **LIMITATIONS**

There are several limitations in this study: First, the need for a larger sample size from larger multicentre cohort to increase the accuracy of the findings and second, information about the day of infection when CT scan was done could not be obtained. Finally, the other factors that might contribute to the disease outcome such as lifestyle and relying on self-reporting/underreporting of the comorbidities should be considered.

## **REFERENCES**

1. Francone M, Iafrate F, Masci GM, Coco S, Cilia F, Manganaro L, Panebianco V, Andreoli C, Colaiacomo MC, Zingaropoli MA, Ciardi MR (2020) Chest CT score in COVID-19 patients: correlation with disease severity and short-term prognosis. *Eur Radiol* 30(12):6808–17. [[PMC free article](#)] [[PubMed](#)]
2. Pan F., Ye T., Sun P. Time course of lung changes at chest CT during recovery from coronavirus disease 2019 (COVID-19) *Radiology*. 2020;295(3):715–721. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
3. Li K., Wu J., Wu F. The clinical and chest CT features associated with severe and critical COVID-19 pneumonia. *Invest Radiol*. 2020;55(6):327–331. [[PMC free](#)]

[article](#)] [[PubMed](#)] [[Google Scholar](#)]

4. Li K., Fang Y., Li W. CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19) *Eur Radiol.* 2020;30(8):4407–4416. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

5. Wong H.Y.F., Lam H.Y.S., Fong A.H.-T. Frequency and distribution of chest radiographic findings in COVID-19 positive patients. *Radiology.* 2020;296(2):E72–E78. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

6. Sethuraman N, Jeremiah SS, Ryo A. Interpreting diagnostic tests for SARS-CoV-2. *JAMA.* 323(22):2249-2251. [[PubMed](#)]

7. Abbasi J. The promise and peril of antibody testing for COVID-19. *JAMA.* 2020;323(19):1881–1883. [[PubMed](#)] [[Google Scholar](#)]

8. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet.* 2020;395(10223):470–473. doi: 10.1016/S0140-6736(20)30185-9. [[PMC free article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

9. Chou R., Pappas M., Buckley D. Use of chest imaging in COVID-19: a rapid advice guide. [https://apps.who.int/iris/bitstream/handle/10665/332326/WHO-2019-nCoV-Clinical- Radiology\\_imaging-Web\\_Annex\\_A-2020.1-eng.pdf](https://apps.who.int/iris/bitstream/handle/10665/332326/WHO-2019-nCoV-Clinical-Radiology_imaging-Web_Annex_A-2020.1-eng.pdf) Accessed June 13, 2020.

10. Wang W, Xu Y, Gao R et al (2020) Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA.* 10.1001/jama.2020.3786 [[PMC free article](#)] [[PubMed](#)]

11. Han H, Luo Q, Mo F, Long L, Zheng W (2020) SARS-CoV-2 RNA more readily detected in induced sputum than in throat swabs of convalescent COVID-19 patients.

Lancet Infect Dis 0. 10.1016/S1473-3099(20)30174-2 [[PMC free article](#)] [[PubMed](#)]

12. Ai T., Yang Z., Hou H. Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology*.

2020;296(2):E32– E40. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

13. Sweta Shah, Tanu Singhal, Namita Davar, and Pooja Thakkar. No correlation between Ct values and severity of disease or mortality in patients with COVID 19 disease. *Indian J Med Microbiol*. 2021 Jan; 39(1): 116–117. [[PMC free article](#)]

[[PubMed](#)]

14. Omar S, Motawea AM, Yasin R. High-resolution CT features of COVID-19 pneumonia in confirmed cases. *Egyptian J Radiol Nucl Med*. 2020;51(1):1–9. doi: 10.1186/s43055-020-

00236-9. [[CrossRef](#)] [[Google Scholar](#)]

15. Borghesi A, Zigliani A, Masciullo R, et al. Radiographic severity index in COVID-19 pneumonia: relationship to age and sex in 783 Italian patients. *La radiologia*

*medica*. 2020;125:461–464. doi: 10.1007/s11547-020-01202-1. [[PMC free](#)

[article](#)] [[PubMed](#)] [[CrossRef](#)] [[Google Scholar](#)]

16. S. Mallapaty, “The coronavirus is most deadly if you are older and male—new data reveal the risks,” *Nature*, vol. 585, no. 7823, pp. 16-17, 2020. View at: [Publisher Site](#) | [Google Scholar](#)

17. W.-j. Guan, W.-h. Liang, Y. Zhao et al., “Comorbidity and its impact on 1590 patients with covid-19 in China: a nationwide analysis,” *European Respiratory Journal*, vol. 55, no.

5, p. 2000547, 2020. View at: [Publisher Site](#) | [Google Scholar](#)

18. Japi. Clinico-radiological evaluation and correlation of CT chest images with progress of disease in COVID-19 patients. <http://www.japi.org>. <https://www.japi.org/w2f4d464/clinico->

radiological-evaluation-and-correlation-of-ct-chest-images-with-progress-of-disease-in-covid-19-patients.

19. Patel SK, Velkoska E, Burrell LM. Emerging markers in cardiovascular disease: where does angiotensin-converting enzyme 2 fit in? *Clin Exp Pharmacol Physiol.* (2013) 40:551–9. doi: 10.1111/1440-1681.12069

20. <http://www.who.int>.. Use of Chest Imaging in COVID-19.

<https://www.who.int/publications/i/item/use-of-chest-imaging-in-covid-19>.

21. Bunawan NC, Mokoagow MI, Djojo AY, Wistriany N, Nugroho HM, Harlivasari AD, Diponegoro S. Delayed RT-PCR Time-To-Positivity in an adult with SARS-CoV-2 Infection. *J Infect Dev Ctries.* 2021 Jul 31;15(7):913-917. doi: 10.3855/jidc.14766. PMID: 34343115

22. Wikramaratna PS, Paton RS, Ghafari M, Lourenço J. Estimating the false-negative test probability of SARS-CoV-2 by RT-PCR. *Euro Surveill.* 2020 Dec;25(50):2000568. doi: 10.2807/1560-7917.ES.2020.25.50.2000568. PMID: 33334398; PMCID: PMC7812420.