

## Original Research Article

# EMULSION PROPERTIES OF SYNBIOTIC YOGHURT SKIN EXTRACT OF RED DRAGON FRUIT (*Hylocereus polyrhizus*) EVAPORATION WITH HONEY

---

### ABSTRACT

Synbiotic yogurt is a combination of probiotics and prebiotics. Red dragon fruit peel extract contains 10.79% pectin, which can be used as a prebiotic. Dragon fruit skin synbiotic yogurt produces an unpleasant aftertaste. The addition of honey can improve the functional properties of dragon fruit peel synbiotic yogurt. The purpose of this study was to determine the emulsion properties and antibacterial activity of synbiotic yogurt with evaporated red dragon fruit peel extract and honey sweetener. This research was conducted at the Laboratory of Animal Products Technology, Faculty of Animal Husbandry, Universitas Brawijaya. The material used is synbiotic yogurt made from skim milk, yogurt starter, red dragon fruit peel extract, and honey. This study used an experimental method in a laboratory using a completely randomized design (CRD) with 3 treatments and 3 replications. Treatment uses yogurt, 20% red dragon fruit peel extract synbiotic yogurt, and synbiotic yogurt with 2% honey sweetener. The data obtained were analyzed using analysis of variance (ANOVA), if the test results showed a difference, then the Duncan test (DMRT) was carried out. The results showed that adding evaporated red dragon fruit peel extract and honey sweetener to synbiotic yogurt did not result in a significant difference ( $P>0.05$ ) in emulsion stability or yogurt turbidity, but did result in a significant difference ( $P>0.05$ ) in emulsion activity. Antibacterial activity on the average number of bacteria, molds and yeasts, and *Escherichia coli* increased, while coliform bacteria and *Enterococcus* spp. decreased. This indicates that the evaporated red dragon fruit peel extract in synbiotic yogurt with honey sweetener contains antibacterial properties. It was concluded that the evaporation of red dragon fruit peel extract with honey sweetener affected the emulsion properties and antibacterial activity.

*Keywords: emulsion stability, emulsion activity, turbidity, red dragon fruit peel, synbiotic yogurt, honey*

### 1. INTRODUCTION

Yogurt is a product obtained from fermented milk using suitable lactic acid bacteria, with/or without the addition of other food ingredients. The physical properties of yogurt play an important role in determining the texture of yogurt products [1]. One of the efforts to improve the functional properties of probiotic yogurt is the addition of prebiotic ingredients made from dietary fiber (prebiotics). The addition of prebiotics to probiotic yogurt is called synbiotic yogurt [2]. Yogurt with the addition of prebiotics is expected to increase the viability of bacteria and store food for microbes in the digestive tract [3]. One of the ingredients that can be used as a prebiotic is the peel of red dragon fruit (*Hylocereus polyrhizus*).

Red dragon fruit peel becomes waste even though the content contained in red dragon fruit peel can be used as a prebiotic and increases the functional value of yogurt. Red dragon fruit peel contains 46.7% fiber, 10.79% pectin, 150.46/100g betacyanin, alkaloid compounds, steroids, saponins, and tannins [4]. The use of additional ingredients such as red dragon fruit peel extract in synbiotic yogurt can stabilize the milk colloid system to prevent phase separation, strengthen the casein network, and reduce syneresis [5]. Pectin in red dragon fruit skin extract is +10.8% [6]. Pectin can function as a gelling agent, a thickening agent, and a stabilizer in food. The addition of red dragon fruit peel extract also affects the taste of the yogurt. Making synbiotic yogurt with red dragon fruit peel extract produces an aftertaste that consumers don't really like [7].

The addition of natural sweeteners such as honey is an attempt to enhance the sweet taste of yogurt. The addition of honey to the yogurt set improves the quality and nutritional value of yogurt and improves its organoleptic properties [8]. The content contained in honey is 41% fructose, 35% glucose, and 1.9% sucrose [9]. Honey additions of up to 3% can improve yogurt quality without affecting the characteristics of lactic acid bacteria (LAB) or yogurt [10]. The polysaccharides contained in honey will wrap oil droplets with steric repulsion because they have larger hydrophilic groups [11]. Sucrose in honey increases the viscosity and lowers the flow index of the aqueous solution [12].

Yogurt with the addition of dietary fiber from red dragon fruit peel extract (*Hylocereus polyrhizus*) and the natural sweetener honey can be studied to determine the properties of yogurt emulsions. The purpose of this study was to review the physical characteristics of synbiotic yogurt, including emulsion activity, emulsion stability, and turbidity.

## 2. MATERIAL AND METHODS

### 2.1 Material

The materials for this research are yoghurt made of skim milk, starter cultures standard yoghurt which contains *Lactobacillus bulgaricus* and *Streptococcus thermophilus* 1:1 (v/v). 20% of red dragon fruit peel extract (*Hylocereus polyrhizus*) was added into synbiotic yoghurt red dragon fruit (*Hylocereus polyrhizus*) was obtained in Malang City, East Java, and the peel was taken. Extraction of red dragon fruit peel (*Hylocereus polyrhizus*) using Microwave-assisted extraction (MAE). The red dragon fruit peel (*Hylocereus polyrhizus*) then be cut into pieces, 50g of red dragon fruit peel added to 50 ml of distilled water. Extraction was carried out in a microwave at 90°C for 5 minutes. Furthermore, 50 ml of the extraction solution was put into a 1 L glass beaker then put into the microwave, evaporated at 70°C for 10 minutes. Probiotic yoghurt (P1) was made from skim milk (10%) with the addition of aquades, as for synbiotic yoghurt (P2 and P3) was added the red dragon fruit peel extract (*Hylocereus polyrhizus*) 20%. All samples were pasteurized at 72°C for 15 minutes, then the temperature was lowered to 42°C, and the mixture was well stirred after inoculating starter yoghurt (3%). Incubation at room temperature (25-28°C) for 24 hours. In treatment P3, the honey 2% was added.

### 2.2 Emulsion Stability and Activity

Emulsion stability and activity was measured according to Arioui *et al.* with modifications. Soybean oil was added to the sample and then homogenized using a hand mixer for 1 minute. As plenty as 0.1 mL of the sample and 0.1% of SDS as plenty as 100 mL and stirred the use of vortex for 1 minutes. Afterward, about 3 mL of sample became taken and put in a cuvette and examined the use of a spectrophotometer with a wavelength of 500 nm and the absorbance value (A0) changed into recorded. After that, the emulsion process waited for 10 min and a comparable test (A10) is performed. The ensuing dispersion is measure through the formula:

Description:

$$\text{Emulsion stability (\%)} = \frac{A_0 \times \Delta t}{A_{10} - A_0}$$

A0 = A<sub>500</sub> at time of 0 minutes A10 = A<sub>500</sub> at time of 10 minutes.

$$\text{Emulsion activity (m}^2\text{/g)} = \frac{2 \times 2,303 \times A_0 \times DF}{1 \times \phi \times C}$$

### 2.3 Turbidity

Turbidity measurement analyzed with 1 ml of sample was taken and dissolved in 100 ml of distilled water. Take 3 ml sample to cuvette. Prepare a spectrophotometer. Determined absorbance of 600 nm. Record the result [13].

### 3. RESULTS AND DISCUSSION

#### 3.1 Emulsion Stability

**Table 1. Emulsion Stability**

Treatment	Stability Emulsion (min)
P1	21.96±2.12
P2	25.45±4.79
P3	25.61±2.80

Based on Table 1, the average stability of the yogurt emulsion shows an increase in value. Emulsion stability indicates material stability. The emulsion contained in the material does not tend to form a separate layer. The protein in milk that is used for yogurt has properties as an emulsifier. Proteins have emulsification properties because of their amphipathic nature (having hydrophobic and hydrophilic groups) and their ability to form layers at the interface with oil and water [14]. The layer formed can reduce the rate of aggregation due to electrostatic repulsion.

The P2 treatment increased due to the addition of extra red dragon fruit peels to the yogurt. Red dragon fruit peel extract contains pectin, which is a polysaccharide. Natural polysaccharides, such as pectin contained in red dragon fruit peel extract (*Hylocereus polyrhizus*) can be good emulsifiers because they have proteins bound to their hydrophilic carbohydrate chains [15]. In addition to polysaccharides, red dragon fruit peel extract contains saponins. Saponins in dragon fruit peel extract can also be used as emulsifiers because they dissolve in water and have hydrophilic carbohydrates that have bonds with non-polar aglycone groups [16]. *Lactobacillus bulgaricus* and *Streptococcus thermophilus* in synbiotic yogurt can synthesize polysaccharides and produce a stabilizer, namely exopolysaccharide. Exopolysaccharides in large quantities can improve the texture better in yogurt [17]. Probiotics that produce exopolysaccharides are known to improve texture and become gel-thickening agents and stabilizers in yogurt [18].

The P3 treatment experienced an increase in value due to the addition of red dragon fruit peel extract and honey. Honey has a sugar content of 41% fructose and 35% glucose [19]. The addition of sugar to yogurt can stabilize the oil-water emulsion [20]. The addition of honey with a sugar content of 41% can prevent the formation of oil droplets into larger droplets so that they will remain separate [21]. The polysaccharides contained in honey will wrap the oil droplets with steric repulsion because they have larger hydrophilic groups [22]. The mechanism of the stabilizer is to form a thin layer that will envelop the emulsified particles, thereby reducing the surface tension and preventing these particles from joining with similar particles [23].

#### 3.2 Emulsion Activity

**Table 2. Emulsion Activity**

Treatment	Emulsion Activity (m <sup>2</sup> /g)
P1	7.68±0.70 <sup>a</sup>
P2	8.84±0.49 <sup>b</sup>
P3	8.90±0.28 <sup>b</sup>

Based on Table 2, the average activity of the yogurt emulsion shows an increase in value. Emulsion activity can be measured by determining the ability of oil to bind to proteins. Emulsion activity indicates the interfacial area that can be obtained between oil and water per unit weight of protein or product [24]. Protein is one of the ingredients that can be used as an emulsifier and is often used in the food industry because it has the characteristic that the surface contains a mixture of hydrophilic and hydrophobic amino acids in the polypeptide chain. The protein layer will surround the oil droplet and result in a decrease in the aggregation rate with electrostatic repulsion if the pH is far from the isoelectric point of the protein [25]. Inhibition of aggression is carried out by electrostatic repulsion, namely binding carbohydrates and forming a thick surface layer [26].

In treatment P2 increased compared to P1. This is due to the addition of red dragon fruit skin extract. The pectin contained in red dragon fruit peel extract is a polysaccharide compound which dissolves in water and is 27 pectinic acids containing methoxyl groups [27]. Pectin is a negatively charged hydrophilic colloid. The casein molecule has a positive and negative charge at neutral pH. Pectin will not bind to pectin at the same charge at a neutral pH, but at a pH below the protein isoelectric (4.6) pectin will be absorbed into the casein micelles through electrostatic interactions causing steric

stability [28]. Steric stability will achieve emulsion stability by pectin [29]. In treatment P3 increased compared to P2. This is due to the addition of red dragon fruit skin extract and honey. Honey has osmolarity properties due to the presence of sugar. The osmolarity of honey as an additive in yogurt will attract water to the casein micelles and reduce the release of water into the environment.

### 3.3 Turbidity

**Table 3. Turbidity**

Treatment	Turbidity (OD)
P1	0.43±0.33
P2	0.63±0.40
P3	0.57±0.33

Based on Table 3, the average turbidity of yogurt shows an increase in value. Turbidity is an index to show the presence of protein aggregation [30]. The average yogurt turbidity shows an increase when compared to P1. This was influenced by the addition of red dragon fruit peel extract (*Hylocereus polyrhizus*). Red dragon fruit peel extract will be utilized by lactic acid bacteria to produce turbidity exopolysaccharide caused by the presence of suspended and dissolved organic and inorganic materials as well as inorganic and organic materials in the form of microorganisms. Exopolysaccharides from Lactic Acid Bacteria (LAB) will inhibit protein-protein interactions, reduce protein aggregation, and cause a decrease in particle size and turbidity [31]. The molecules contained in yogurt affect the level of absorbance by scattering light. The higher the absorbance value, the more dispersed the metabolites in the yogurt are. Turbidity is related to the size and concentration of dissolved particles [32]. Metabolites released by lactic acid LAB can increase the absorbance value of yogurt [33].

The P2 treatment increased due to the presence of red dragon fruit skin extract. Dragon fruit skin contains 20.1% pectin [34]. Dragon fruit peel extract will affect turbidity because pectin will interact with casein. The pectin and casein bonds will form an enlargement of the particle size in yogurt. Pectin will form several layers in casein so that the particle size increases, absorbance increases, and turbidity also increases [35]. The electrostatic interaction that occurs between pectin and casein's negative charge prevents the aggregation of casein [36].

P3 treatment decreased when compared to P2 due to the addition of honey to synbiotic yogurt. The addition of honey to synbiotic yogurt lowers the turbidity value. The decrease in turbidity indicates the uniformity of the particle size in the yogurt. Small and uniform particle size will increase the surface area. The ability to absorb color or the level of clarity is determined by the surface area of the % transmittance, where the greater the surface area of the transmittance, the higher the level of clarity of a solution (clear). The surface area of % transmittance will vary for each type of raw material and concentration used [37].

## 4. CONCLUSION

The addition of red dragon fruit peel extract and honey affected the emulsion stability, emulsion activity and turbidity of yogurt. This is due to the addition of 20% red dragon fruit skin extract (*Hylocereus polyrhizus*) and 2% honey. Honey can affect emulsion stability, emulsion activity and turbidity due to the presence of fructose.

## REFERENCES

- [1] Shaker RR, Jumah RY, Abu-Jdayil B. Rheological properties of plain yogurt during coagulation process: impact of fat content and preheat treatment of milk. *Journal of Food Engineering*. 2000;44(3):175-80.
- [2] Yaumi Y, Hadju R, Yelnetty A, Lontaan NN. Kualitas sensoris yoghurt sinbiotik menggunakan pati termodifikasi dari umbi uwi ungu (*Dioscorea alata*). *Zootec*. 2020;40(1):196-206.
- [3] Tari AI, Hartati S. Performa kesehatan tikus sprague dawley akibat pemberian yoghurt prebiotik sebagai anti-diare. *Jurnal Ilmiah Teknosains*. 2018;4(2):108-113.
- [4] Jamilah B, Shu CE, Kharidah M, Dzulkily MA, Noranizan A. Physico-chemical characteristics of red pitaya (*Hylocereus polyrhizus*) peel. *International Food Research Journal*. 2011;18(1).

- [5] Hematyar N, Samarin AM, Poorazarang H, Elhamirad AH. Effect of gums on yogurt characteristics. *World Applied Sciences Journal*. 2012;20(5):661-5.
- [6] Yati K, Ladeska V, Wirman AP. Isolasi pektin dari kulit buah naga (*Hylocereus polyrhizus*) dan pemanfaatannya sebagai pengikat pada sediaan pasta gigi. *Media Farmasi*. 2017;14(1):1-6.
- [7] Nirmalawaty A, Mahayani AA. Analisa kimia bakpia kering substitusi tepung kulit buah naga. *STIGMA: Jurnal Matematika dan Ilmu Pengetahuan Alam Unipa*. 2020;13(01):15-23.
- [8] Sert D, Akin N, Dertli E. Effects of sunflower honey on the physicochemical, microbiological and sensory characteristics in set type yoghurt during refrigerated storage. *International journal of dairy technology*. 2011;64(1):99-107.
- [9] Yurliasni Y, Hanum Z, Hikmawan R. Potensi madu dalam meningkatkan kualitas minuman kefir. *Jurnal Ilmu dan Teknologi Hasil Ternak (JITEK)*. 2019;14(1):50-9.
- [10] Varga L. Effect of acacia (*Robinia pseudo-acacia* L.) honey on the characteristic microflora of yogurt during refrigerated storage. *International journal of food microbiology*. 2006;108(2):272-5.
- [11] Ozturk B, McClements DJ. Progress in natural emulsifiers for utilization in food emulsions. *Current Opinion in Food Science*. 2016;7:1-6.
- [12] Sintasari RA, Kurnadi J, Ningtyas DW. Pengaruh penambahan konsentrasi susu skim dan sukrosa terhadap karakteristik minuman probiotik sari beras merah [in press juli 2014]. *Jurnal pangan dan Agroindustri*. 2014;2(3):65-75.
- [13] Liu YF, Oey I, Bremer P, Carne A, Silcock P. Effects of pH, temperature and pulsed electric fields on the turbidity and protein aggregation of ovomucin-depleted egg white. *Food research international*. 2017;91:161-170.
- [14] McClements DJ. Theoretical analysis of factors affecting the formation and stability of multilayered colloidal dispersions. *Langmuir*. 2005;21(21):9777-85.
- [15] Dickinson E. Hydrocolloids at interfaces and the influence on the properties of dispersed systems. *Food hydrocolloids*. 2003;17(1):25-39.
- [16] Osbourn A, Goss RJ, Field RA. The saponins–polar isoprenoids with important and diverse biological activities. *Natural product reports*. 2011;28(7):1261-8.
- [17] Han X, Yang Z, Jing X, Yu P, Zhang Y, Yi H, Zhang L. Improvement of the texture of yogurt by use of exopolysaccharide producing lactic acid bacteria. *BioMed research international*. 2016;1-6.
- [18] Jurášková D, Ribeiro SC, Silva CC. Exopolysaccharides produced by lactic acid bacteria: From biosynthesis to health-promoting properties. *Foods*. 2022;11(2):156.
- [19] Yurliasni Y, Hanum Z, Hikmawan R. Potensi madu dalam meningkatkan kualitas minuman kefir. *Jurnal Ilmu dan Teknologi Hasil Ternak (JITEK)*. 2019;14(1):50-9.
- [20] Maskan M, Göğüş F. Effect of sugar on the rheological properties of sunflower oil–water emulsions. *Journal of food engineering*. 2000;43(3):173-7.
- [21] Alpaslan M, Hayta ME. Rheological and sensory properties of pekmez (grape molasses)/tahin (sesame paste) blends. *Journal of Food Engineering*. 2002;54(1):89-93.
- [22] Ozturk B, McClements DJ. Progress in natural emulsifiers for utilization in food emulsions. *Current Opinion in Food Science*. 2016;7(1):1-6.
- [23] Hambali E, Suryani A, Rivai M, Permadi P. *Teknologi surfaktan dan aplikasinya (edisi revisi)*. Bogor : PT Penerbit IPB Press; 2019.

- [24] Pearce KN, Kinsella JE. Emulsifying properties of proteins: evaluation of a turbidimetric technique. *Journal of agricultural and food chemistry*. 1978;26(3):716-23.
- [25] McClements DJ. Theoretical analysis of factors affecting the formation and stability of multilayered colloidal dispersions. *Langmuir*. 2005;21(21):9777-85.
- [26] Wooster TJ, Augustin MA. The emulsion flocculation stability of protein-carbohydrate diblock copolymers. *Journal of colloid and interface science*. 2007;313(2):665-75.
- [27] Husni P, Ikhrom UK, Hasanah U. Uji dan Karakterisasi Serbuk Pektin dari Albedo Durian sebagai Kandidat Eksiipien Farmasi. *Majalah Farmasetika*. 2021;6(3):202-212
- [28] Wusigale, Liang LI, Luo Y. Casein and pectin: Structures, interactions, and applications. *Trends in Food Science & Technology*. 2020;97:391-403.
- [29] Nguémazong ED, Christiaens S, Shpigelman A, Van Loey A, Hendrickx M. The emulsifying and emulsion-stabilizing properties of pectin: A review. *Comprehensive Reviews in Food Science and Food Safety*. 2015;14(6):705-18.
- [30] Yang X, Ke C, Li L. Physicochemical, rheological and digestive characteristics of soy protein isolate gel induced by lactic acid bacteria. *Journal of Food Engineering*. 2021;292(1):1-11
- [31] Abid Y, Joulak I, Amara CB, Casillo A, Attia H, Gharsallaoui A, Azabou S. Study of interactions between anionic exopolysaccharides produced by newly isolated probiotic bacteria and sodium caseinate. *Colloids and Surfaces B: Biointerfaces*. 2018;167:516-23.
- [32] He J, Giusti MM. Anthocyanins: natural colorants with health-promoting properties. *Annu. Rev. Food Sci. Technol.* 2010;1(1):163-87.
- [33] Ryu JK, Lee HS, Koo BK, Kim HK. Comparison of the biochemical activities of commercial yogurts and lactobacillus acidophilus-containing yogurt. *Korean Journal of Clinical Laboratory Science*. 2015;47(2):59-64.
- [34] Nazaruddin R, Norazelina SM, Norziah MH, Zainudin M. Pectins from dragon fruit (*Hylocereus polyrhizus*) peel. *Malaysian Applied Biology*. 2011;40(1):19-23.
- [35] Tuinier R, Rolin C, De Kruif CG. Electrosorption of pectin onto casein micelles. *Biomacromolecules*. 2002;3(3):632-8.
- [36] Jensen S, Rolin C, Ipsen R. Stabilisation of acidified skimmed milk with HM pectin. *Food hydrocolloids*. 2010;24(4):291-9.
- [37] Jaya F. Produk-produk lebah madu dan hasil olahannya. Malang: Universitas Brawijaya Press; 2017.