

# A Study on Generalized Blaise Numbers

**Abstract.** In this paper, we introduce and investigate the generalized Blaise sequences and we deal with, in detail, two special cases, namely, Blaise and Blaise-Lucas sequences. We present Binet's formulas, generating functions, Simson formulas, and the summation formulas for these sequences. Furthermore, we show that there are close relations between Blaise, Blaise-Lucas and Jacobsthal-Padovan, Jacobsthal-Perrin, adjusted Jacobsthal-Padovan, modified Jacobsthal-Padovan numbers. Moreover, we give some identities and matrices related with these sequences.

**2020 Mathematics Subject Classification.** 11B37, 11B39, 11B83.

**Keywords.** Blaise numbers, Blaise-Lucas numbers, Jacobsthal-Padovan numbers, Jacobsthal-Perrin numbers, adjusted Jacobsthal-Padovan numbers, modified Jacobsthal-Padovan numbers.

## 1. Introduction

Adjusted Jacobsthal-Padovan sequence  $\{K_n\}_{n \geq 0}$  (OEIS: A159287, [31]), Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas) sequence  $\{L_n\}_{n \geq 0}$  (OEIS: A072328, [31]), Jacobsthal-Padovan sequence  $\{Q_n\}_{n \geq 0}$  (OEIS: A159284, [31]), and modified Jacobsthal-Padovan sequence  $\{M_n\}_{n \geq 0}$  are defined, respectively, by the third-order recurrence relations

$$K_{n+3} = K_{n+1} + 2K_n, \quad K_0 = 0, K_1 = 1, K_2 = 0, \quad (1.1)$$

$$L_{n+3} = L_{n+1} + 2L_n, \quad L_0 = 3, L_1 = 0, L_2 = 2, \quad (1.2)$$

$$Q_{n+3} = Q_{n+1} + 2Q_n, \quad Q_0 = 1, Q_1 = 1, Q_2 = 1, \quad (1.3)$$

$$M_{n+3} = M_{n+1} + 2M_n, \quad M_0 = 3, M_1 = 1, M_2 = 3. \quad (1.4)$$

The sequences  $\{Q_n\}_{n \geq 0}$ ,  $\{L_n\}_{n \geq 0}$ ,  $\{K_n\}_{n \geq 0}$  and  $\{M_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$\begin{aligned} K_{-n} &= -\frac{1}{2}K_{-(n-1)} + \frac{1}{2}K_{-(n-3)}, \\ L_{-n} &= -\frac{1}{2}L_{-(n-1)} + \frac{1}{2}L_{-(n-3)}, \\ Q_{-n} &= -\frac{1}{2}Q_{-(n-1)} + \frac{1}{2}Q_{-(n-3)}, \\ M_{-n} &= -\frac{1}{2}M_{-(n-1)} + \frac{1}{2}M_{-(n-3)}, \end{aligned}$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrences (1.1)-(1.4) hold for all integer  $n$ . For more information on Jacobsthal-Padovan sequence, see [38] and [8].

Now, we define two sequences related to Adjusted Jacobsthal-Padovan, Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas), Jacobsthal-Padovan, and modified Jacobsthal-Padovan numbers. Blaise and Blaise-Lucas numbers are defined as

$$B_n = B_{n-2} + 2B_{n-3} + 1, \quad \text{with } B_0 = 0, B_1 = 1, B_2 = 1, \quad n \geq 3,$$

and

$$C_n = C_{n-2} + 2C_{n-3} - 2, \quad \text{with } C_0 = 4, C_1 = 1, C_2 = 3, \quad n \geq 3,$$

respectively. The first few values of Blaise and Blaise-Lucas numbers are

$$0, 1, 1, 2, 4, 5, 9, 14, 20, 33, 49, 74, 116, 173, \dots$$

and

$$4, 1, 3, 7, 3, 11, 15, 15, 35, 43, 63, 111, 147, 235, \dots$$

respectively. The sequences  $\{B_n\}$  and  $\{C_n\}$  satisfy the following fourth order linear recurrences:

$$\begin{aligned} B_n &= B_{n-1} + B_{n-2} + B_{n-3} - 2B_{n-4}, & B_0 = 0, B_1 = 1, B_2 = 1, B_3 = 2, & n \geq 4, \\ C_n &= C_{n-1} + C_{n-2} + C_{n-3} - 2C_{n-4}, & C_0 = 4, C_1 = 1, C_2 = 3, C_3 = 7, & n \geq 4. \end{aligned}$$

There are close relations between Blaise, Blaise-Lucas and Adjusted Jacobsthal-Padovan, Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas), Jacobsthal-Padovan, and modified Jacobsthal-Padovan numbers. For example,

they satisfy the following interrelations:

$$\begin{aligned}
 2B_n &= K_{n+2} + K_{n+1} + 2K_n - 1, \\
 2C_n &= -K_{n+2} + 6K_{n+1} + K_n + 2, \\
 52B_n &= 2L_n + 9L_{n+1} + 10L_{n+2} - 26, \\
 C_n &= L_n + 1, \\
 2B_n &= Q_{n+2} - 1, \\
 2C_n &= -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2, \\
 46B_n &= 8M_{n+2} + 5M_{n+1} - 2M_n - 23, \\
 46C_n &= -5M_{n+2} - 6M_{n+1} + 53M_n + 46,
 \end{aligned}$$

and

$$\begin{aligned}
 2K_n &= -B_{n+2} + 3B_n + 1, \\
 52K_n &= 9C_{n+2} - C_{n+1} - 6C_n - 2, \\
 4L_n &= -3B_{n+2} + 14B_{n+1} - 9B_n + 1, \\
 2L_n &= C_{n+3} - C_{n+1}, \\
 2Q_n &= -B_{n+2} + 2B_{n+1} + B_n + 1, \\
 26Q_n &= 4C_{n+2} + C_{n+1} + 6C_n - 11, \\
 2M_n &= -B_{n+2} + 6B_{n+1} - 3B_n + 1, \\
 26M_n &= 3C_{n+2} + 4C_{n+1} + 24C_n - 31.
 \end{aligned}$$

The purpose of this article is to generalize and investigate these interesting sequence of numbers (i.e., Blaise, Blaise-Lucas numbers). First, we recall some properties of the generalized Tetranacci numbers.

The generalized  $(r, s, t, u)$  sequence (or generalized Tetranacci sequence or generalized 4-step Fibonacci sequence)  $\{W_n(W_0, W_1, W_2, W_3; r, s, t, u)\}_{n \geq 0}$  (or shortly  $\{W_n\}_{n \geq 0}$ ) is defined as follows:

$$W_n = rW_{n-1} + sW_{n-2} + tW_{n-3} + uW_{n-4}, \quad W_0 = c_0, W_1 = c_1, W_2 = c_2, W_3 = c_3, \quad n \geq 4 \quad (1.5)$$

where  $W_0, W_1, W_2, W_3$  are arbitrary complex (or real) numbers and  $r, s, t, u$  are real numbers.

This sequence has been studied by many authors and more detail can be found in the extensive literature dedicated to these sequences, see for example [15,22,23,29,33,35,36,59,60]. The sequence  $\{W_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$W_{-n} = -\frac{t}{u}W_{-(n-1)} - \frac{s}{u}W_{-(n-2)} - \frac{r}{u}W_{-(n-3)} + \frac{1}{u}W_{-(n-4)}$$

for  $n = 1, 2, 3, \dots$  when  $u \neq 0$ . Therefore, recurrence (1.5) holds for all integers  $n$ .

As  $\{W_n\}$  is a fourth-order recurrence sequence (difference equation), its characteristic equation is

$$z^4 - rz^3 - sz^2 - tz - u = 0 \quad (1.6)$$

whose roots are  $\alpha, \beta, \gamma, \delta$ . Note that we have the following identities

$$\begin{aligned} \alpha + \beta + \gamma + \delta &= r, \\ \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta &= -s, \\ \alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta &= t, \\ \alpha\beta\gamma\delta &= -u. \end{aligned}$$

Using these roots and the recurrence relation, Binet's formula can be given as follows:

**THEOREM 1.** *(Four Distinct Roots Case:  $\alpha \neq \beta \neq \gamma \neq \delta$ ) For all integers  $n$ , Binet's formula of generalized Tetranacci numbers is*

$$W_n = \frac{p_1 \alpha^n}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{p_2 \beta^n}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{p_3 \gamma^n}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{p_4 \delta^n}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)} \quad (1.7)$$

where

$$\begin{aligned} p_1 &= W_3 - (\beta + \gamma + \delta)W_2 + (\beta\gamma + \beta\delta + \gamma\delta)W_1 - \beta\gamma\delta W_0, \\ p_2 &= W_3 - (\alpha + \gamma + \delta)W_2 + (\alpha\gamma + \alpha\delta + \gamma\delta)W_1 - \alpha\gamma\delta W_0, \\ p_3 &= W_3 - (\alpha + \beta + \delta)W_2 + (\alpha\beta + \alpha\delta + \beta\delta)W_1 - \alpha\beta\delta W_0, \\ p_4 &= W_3 - (\alpha + \beta + \gamma)W_2 + (\alpha\beta + \alpha\gamma + \beta\gamma)W_1 - \alpha\beta\gamma W_0. \end{aligned}$$

Usually, it is customary to choose  $\alpha, \beta, \gamma, \delta$  so that the Equ. (1.6) has at least one real (say  $\alpha$ ) solutions. Note that the Binet form of a sequence satisfying (1.6) for non-negative integers is valid for all integers  $n$  (see [16]).

Next, we consider two special cases of the generalized  $(r, s, t, u)$  sequence  $\{W_n\}$  which we call them  $(r, s, t, u)$ -Fibonacci and  $(r, s, t, u)$ -Lucas sequences.  $(r, s, t, u)$ -Fibonacci sequence  $\{G_n\}_{n \geq 0}$  and  $(r, s, t, u)$ -Lucas sequence  $\{H_n\}_{n \geq 0}$  are defined, respectively, by the fourth-order recurrence relations

$$G_{n+4} = rG_{n+3} + sG_{n+2} + tG_{n+1} + uG_n, \quad (1.8)$$

$$G_0 = 0, G_1 = 1, G_2 = r, G_3 = r^2 + s,$$

$$H_{n+4} = rH_{n+3} + sH_{n+2} + tH_{n+1} + uH_n, \quad (1.9)$$

$$H_0 = 4, H_1 = r, H_2 = 2s + r^2, H_3 = r^3 + 3sr + 3t.$$

The sequences  $\{G_n\}_{n \geq 0}$  and  $\{H_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$\begin{aligned} G_{-n} &= -\frac{t}{u}G_{-(n-1)} - \frac{s}{u}G_{-(n-2)} - \frac{r}{u}G_{-(n-3)} + \frac{1}{u}G_{-(n-4)}, \\ H_{-n} &= -\frac{t}{u}H_{-(n-1)} - \frac{s}{u}H_{-(n-2)} - \frac{r}{u}H_{-(n-3)} + \frac{1}{u}H_{-(n-4)}, \end{aligned}$$

for  $n = 1, 2, 3, \dots$  respectively. Therefore, recurrences (1.8) and (1.9) hold for all integers  $n$ .

For all integers  $n$ ,  $(r, s, t, u)$ -Fibonacci and  $(r, s, t, u)$ -Lucas numbers (using initial conditions in (1.8) or (1.9)) can be expressed using Binet's formulas as in the following corollary.

**COROLLARY 2.** *(Four Distinct Roots Case:  $\alpha \neq \beta \neq \gamma \neq \delta$ ) Binet's formula of  $(r, s, t, u)$ -Fibonacci and  $(r, s, t, u)$ -Lucas numbers are*

$$G_n = \frac{\alpha^{n+2}}{(\alpha - \beta)(\alpha - \gamma)(\alpha - \delta)} + \frac{\beta^{n+2}}{(\beta - \alpha)(\beta - \gamma)(\beta - \delta)} + \frac{\gamma^{n+2}}{(\gamma - \alpha)(\gamma - \beta)(\gamma - \delta)} + \frac{\delta^{n+2}}{(\delta - \alpha)(\delta - \beta)(\delta - \gamma)}$$

and

$$H_n = \alpha^n + \beta^n + \gamma^n + \delta^n,$$

respectively.

*Proof.* Take  $W_n = G_n$  and  $W_n = H_n$  in Theorem 1, respectively.  $\square$

Next, we give the ordinary generating function  $\sum_{n=0}^{\infty} W_n z^n$  of the sequence  $W_n$ .

**LEMMA 3.** *Suppose that  $f_{W_n}(z) = \sum_{n=0}^{\infty} W_n z^n$  is the ordinary generating function of the generalized  $(r, s, t, u)$  sequence  $\{W_n\}_{n \geq 0}$ . Then,  $\sum_{n=0}^{\infty} W_n z^n$  is given by*

$$\sum_{n=0}^{\infty} W_n z^n = \frac{W_0 + (W_1 - rW_0)z + (W_2 - rW_1 - sW_0)z^2 + (W_3 - rW_2 - sW_1 - tW_0)z^3}{1 - rz - sz^2 - tz^3 - uz^4}. \quad (1.10)$$

*Proof.* For a proof, see Soykan [33, Lemma 1].  $\square$

The following theorem presents Simson's formula of generalized  $(r, s, t, u)$  sequence (generalized Tetranacci sequence)  $\{W_n\}$ .

**THEOREM 4** (Simson's Formula of Generalized  $(r, s, t, u)$  Numbers). *For all integers  $n$ , we have*

$$\begin{vmatrix} W_{n+3} & W_{n+2} & W_{n+1} & W_n \\ W_{n+2} & W_{n+1} & W_n & W_{n-1} \\ W_{n+1} & W_n & W_{n-1} & W_{n-2} \\ W_n & W_{n-1} & W_{n-2} & W_{n-3} \end{vmatrix} = (-1)^n u^n \begin{vmatrix} W_3 & W_2 & W_1 & W_0 \\ W_2 & W_1 & W_0 & W_{-1} \\ W_1 & W_0 & W_{-1} & W_{-2} \\ W_0 & W_{-1} & W_{-2} & W_{-3} \end{vmatrix}. \quad (1.11)$$

*Proof.* (1.11) is given in Soykan [32].  $\square$

The following theorem shows that the generalized Tetranacci sequence  $W_n$  at negative indices can be expressed by the sequence itself at positive indices.

THEOREM 5. For  $n \in \mathbb{Z}$ , for the generalized Tetranacci sequence (or generalized  $(r, s, t, u)$ -sequence or 4-step Fibonacci sequence) we have the following:

$$\begin{aligned} W_{-n} &= \frac{1}{6}(-u)^{-n}(-6W_{3n} + 6H_nW_{2n} - 3H_n^2W_n + 3H_{2n}W_n + W_0H_n^3 + 2W_0H_{3n} - 3W_0H_nH_{2n}) \\ &= (-1)^{-n-1}u^{-n}(W_{3n} - H_nW_{2n} + \frac{1}{2}(H_n^2 - H_{2n})W_n - \frac{1}{6}(H_n^3 + 2H_{3n} - 3H_{2n}H_n)W_0). \end{aligned}$$

Proof. For the proof, see Soykan [34, Theorem 1.].  $\square$

Using Theorem 5, we have the following corollary, see Soykan [34, Corollary 4].

COROLLARY 6. For  $n \in \mathbb{Z}$ , we have

$$\begin{aligned} \text{(a): } 2(-u)^{n+4}G_{-n} &= -(3ru^2 + t^3 - 3stu)^2G_n^3 - (2su - t^2)^2G_{n+3}^2G_n - (-rt^2 - tu + 2rsu)^2G_{n+2}^2G_n \\ &\quad - (-st^2 + 2s^2u + 4u^2 + rtu)^2G_{n+1}^2G_n + 2(3ru^2 + t^3 - 3stu)((-2su + t^2)G_{n+3} + (-rt^2 - tu + 2rsu)G_{n+2} + \\ &\quad (-st^2 + 2s^2u + 4u^2 + rtu)G_{n+1})G_n^2 + 2(2su - t^2)(-rt^2 - tu + 2rsu)G_{n+3}G_{n+2}G_n + 2(2su - t^2)(-st^2 + \\ &\quad 2s^2u + 4u^2 + rtu)G_{n+3}G_{n+1}G_n - 2(-st^2 + 2s^2u + 4u^2 + rtu)(-rt^2 - tu + 2rsu)G_{n+2}G_{n+1}G_n - 2G_{3n}u^4 + \\ &\quad u^2(-2su + t^2)G_{2n+3}G_n + u^2(-rt^2 - tu + 2rsu)G_{2n+2}G_n + u^2(-st^2 + 2s^2u + 4u^2 + rtu)G_{2n+1}G_n - \\ &\quad 2u^2(2su - t^2)G_{2n}G_{n+3} + 2u^2(-rt^2 - tu + 2rsu)G_{2n}G_{n+2} + 2u^2(-st^2 + 2s^2u + 4u^2 + rtu)G_{2n}G_{n+1} - \\ &\quad 3u^2(3ru^2 + t^3 - 3stu)G_{2n}G_n. \\ \text{(b): } H_{-n} &= \frac{1}{6}(-u)^{-n}(H_n^3 + 2H_{3n} - 3H_{2n}H_n). \end{aligned}$$

Note that  $G_{-n}$  and  $H_{-n}$  can be given as follows by using  $G_0 = 0$  and  $H_0 = 4$  in Theorem 5,

$$G_{-n} = \frac{1}{6}(-u)^{-n}(-6G_{3n} + 6H_nG_{2n} - 3H_n^2G_n + 3H_{2n}G_n), \quad (1.12)$$

$$H_{-n} = \frac{1}{6}(-u)^{-n}(H_n^3 + 2H_{3n} - 3H_{2n}H_n), \quad (1.13)$$

respectively.

If we define the square matrix  $A$  of order 4 as

$$A = A_{rstu} = \begin{pmatrix} r & s & t & u \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

and also define

$$B_n = \begin{pmatrix} G_{n+1} & sG_n + tG_{n-1} + uG_{n-2} & tG_n + uG_{n-1} & uG_n \\ G_n & sG_{n-1} + tG_{n-2} + uG_{n-3} & tG_{n-1} + uG_{n-2} & uG_{n-1} \\ G_{n-1} & sG_{n-2} + tG_{n-3} + uG_{n-4} & tG_{n-2} + uG_{n-3} & uG_{n-2} \\ G_{n-2} & sG_{n-3} + tG_{n-4} + uG_{n-5} & tG_{n-3} + uG_{n-4} & uG_{n-3} \end{pmatrix}$$

and

$$U_n = \begin{pmatrix} W_{n+1} & sW_n + tW_{n-1} + uW_{n-2} & tW_n + uW_{n-1} & uW_n \\ W_n & sW_{n-1} + tW_{n-2} + uW_{n-3} & tW_{n-1} + uW_{n-2} & uW_{n-1} \\ W_{n-1} & sW_{n-2} + tW_{n-3} + uW_{n-4} & tW_{n-2} + uW_{n-3} & uW_{n-2} \\ W_{n-2} & sW_{n-3} + tW_{n-4} + uW_{n-5} & tW_{n-3} + uW_{n-4} & uW_{n-3} \end{pmatrix}$$

then we get the following Theorem.

**THEOREM 7.** *For all integers  $m, n$ , we have*

**(a):**  $B_n = A^n$ , i.e.,

$$\begin{pmatrix} r & s & t & u \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n = \begin{pmatrix} G_{n+1} & sG_n + tG_{n-1} + uG_{n-2} & tG_n + uG_{n-1} & uG_n \\ G_n & sG_{n-1} + tG_{n-2} + uG_{n-3} & tG_{n-1} + uG_{n-2} & uG_{n-1} \\ G_{n-1} & sG_{n-2} + tG_{n-3} + uG_{n-4} & tG_{n-2} + uG_{n-3} & uG_{n-2} \\ G_{n-2} & sG_{n-3} + tG_{n-4} + uG_{n-5} & tG_{n-3} + uG_{n-4} & uG_{n-3} \end{pmatrix}.$$

**(b):**  $U_1 A^n = A^n U_1$ .

**(c):**  $U_{n+m} = U_n B_m = B_m U_n$ .

**Proof.** For the proof, see Soykan [33, Theorem 19].  $\square$

**THEOREM 8.** *For all integers  $m, n$ , we have*

$$W_{n+m} = W_n G_{m+1} + W_{n-1}(sG_m + tG_{m-1} + uG_{m-2}) + W_{n-2}(tG_m + uG_{m-1}) + uW_{n-3}G_m. \quad (1.14)$$

**Proof.** For the proof, see Soykan [33, Theorem 20].  $\square$

In the next sections, we present new results.

## 2. Generalized Blaise Sequence

In this paper, we consider the case  $r = 1, s = 1, t = 1, u = -2$ . A generalized Blaise sequence  $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2, W_3)\}_{n \geq 0}$  is defined by the fourth-order recurrence relation

$$W_n = W_{n-1} + W_{n-2} + W_{n-3} - 2W_{n-4} \quad (2.1)$$

with the initial values  $W_0 = c_0, W_1 = c_1, W_2 = c_2, W_3 = c_3$  not all being zero.

The sequence  $\{W_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$W_{-n} = \frac{1}{2}W_{-(n-1)} + \frac{1}{2}W_{-(n-2)} + \frac{1}{2}W_{-(n-3)} - \frac{1}{2}W_{-(n-4)}$$

for  $n = 1, 2, 3, \dots$ . Therefore, recurrence (2.1) holds for all integers  $n$ .

Characteristic equation of  $\{W_n\}$  is

$$z^4 - z^3 - z^2 - z + 2 = (z^3 - z - 2)(z - 1) = 0$$

whose roots are

$$\begin{aligned}\alpha &= \sqrt[3]{1 + \frac{\sqrt{78}}{9}} + \sqrt[3]{1 - \frac{\sqrt{78}}{9}} \simeq 1.521379706804568, \\ \beta &= \omega \sqrt[3]{1 + \frac{\sqrt{78}}{9}} + \omega^2 \sqrt[3]{1 - \frac{\sqrt{78}}{9}}, \\ \gamma &= \omega^2 \sqrt[3]{1 + \frac{\sqrt{78}}{9}} + \omega \sqrt[3]{1 - \frac{\sqrt{78}}{9}}, \\ \delta &= 1,\end{aligned}$$

where

$$\omega = \frac{-1 + i\sqrt{3}}{2} = \exp(2\pi i/3).$$

Note that

$$\begin{aligned}\alpha + \beta + \gamma + \delta &= 1, \\ \alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta &= -1, \\ \alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta &= 1, \\ \alpha\beta\gamma\delta &= 2.\end{aligned}$$

Note also that

$$\begin{aligned}\alpha + \beta + \gamma &= 0, \\ \alpha\beta + \alpha\gamma + \beta\gamma &= -1, \\ \alpha\beta\gamma &= 2.\end{aligned}$$

The first few generalized Blaise numbers with positive subscript and negative subscript are given in the following Table 1.

Table 1. A few generalized Blaise numbers

$n$	$W_n$	$W_{-n}$
0	$W_0$	$W_0$
1	$W_1$	$\frac{1}{2}(W_0 + W_1 + W_2 - W_3)$
2	$W_2$	$\frac{1}{4}(3W_0 + 3W_1 - W_2 - W_3)$
3	$W_3$	$\frac{1}{8}(9W_0 + W_1 + W_2 - 3W_3)$
4	$W_1 - 2W_0 + W_2 + W_3$	$\frac{1}{16}(11W_0 + 11W_1 + 3W_2 - 9W_3)$
5	$2W_2 - W_1 - 2W_0 + 2W_3$	$\frac{1}{32}(33W_0 + 17W_1 - 7W_2 - 11W_3)$
6	$W_2 - 4W_0 + 4W_3$	$\frac{1}{64}(67W_0 + 19W_1 + 11W_2 - 33W_3)$
7	$4W_2 - 8W_0 + 5W_3$	$\frac{1}{128}(105W_0 + 89W_1 + W_2 - 67W_3)$
8	$5W_2 - 3W_1 - 10W_0 + 9W_3$	$\frac{1}{256}(283W_0 + 107W_1 - 29W_2 - 105W_3)$
9	$6W_2 - W_1 - 18W_0 + 14W_3$	$\frac{1}{512}(497W_0 + 225W_1 + 73W_2 - 283W_3)$
10	$13W_2 - 4W_1 - 28W_0 + 20W_3$	$\frac{1}{1024}(947W_0 + 643W_1 - 69W_2 - 497W_3)$
11	$16W_2 - 8W_1 - 40W_0 + 33W_3$	$\frac{1}{2048}(2233W_0 + 809W_1 - 47W_2 - 947W_3)$
12	$25W_2 - 7W_1 - 66W_0 + 49W_3$	$\frac{1}{4096}(3851W_0 + 2139W_1 + 339W_2 - 2233W_3)$
13	$42W_2 - 17W_1 - 98W_0 + 74W_3$	$\frac{1}{8192}(8129W_0 + 4529W_1 - 615W_2 - 3851W_3)$

Note that the sequences  $\{B_n\}$  and  $\{C_n\}$  which are defined in the section Introduction, are the special cases of the generalized Blaise sequence  $\{W_n\}$ . For convenience, we can give the definition of these two special cases of the sequence  $\{W_n\}$  in this section as well. Blaise sequence  $\{B_n\}_{n \geq 0}$  and Blaise-Lucas sequence  $\{C_n\}_{n \geq 0}$  are defined, respectively, by the fourth-order recurrence relations

$$\begin{aligned}
 B_n &= B_{n-1} + B_{n-2} + B_{n-3} - 2B_{n-4}, & B_0 = 0, B_1 = 1, B_2 = 1, B_3 = 2, & n \geq 4, \\
 C_n &= C_{n-1} + C_{n-2} + C_{n-3} - 2C_{n-4}, & C_0 = 4, C_1 = 1, C_2 = 3, C_3 = 7, & n \geq 4.
 \end{aligned}$$

The sequences  $\{B_n\}_{n \geq 0}$  and  $\{C_n\}_{n \geq 0}$  can be extended to negative subscripts by defining

$$\begin{aligned}
 B_{-n} &= \frac{1}{2}B_{-(n-1)} + \frac{1}{2}B_{-(n-2)} + \frac{1}{2}B_{-(n-3)} - \frac{1}{2}B_{-(n-4)} \\
 C_{-n} &= \frac{1}{2}C_{-(n-1)} + \frac{1}{2}C_{-(n-2)} + \frac{1}{2}C_{-(n-3)} - \frac{1}{2}C_{-(n-4)}
 \end{aligned}$$

for  $n = 1, 2, 3, \dots$  respectively.

Next, we present the first few values of the Blaise and Blaise-Lucas numbers with positive and negative subscripts:

Table 2. The first few values of the special third-order numbers with positive and negative subscripts.

$n$	0	1	2	3	4	5	6	7	8	9	10	11	12	13
$B_n$	0	1	1	2	4	5	9	14	20	33	49	74	116	173
$B_{-n}$	0	0	0	$-\frac{1}{2}$	$-\frac{1}{4}$	$-\frac{3}{8}$	$-\frac{9}{16}$	$-\frac{11}{32}$	$-\frac{33}{64}$	$-\frac{67}{128}$	$-\frac{105}{256}$	$-\frac{283}{512}$	$-\frac{497}{1024}$	$-\frac{947}{2048}$
$C_n$	4	1	3	7	3	11	15	15	35	43	63	111	147	235
$C_{-n}$	4	$\frac{1}{2}$	$\frac{5}{4}$	$\frac{19}{8}$	$\frac{1}{16}$	$\frac{51}{32}$	$\frac{89}{64}$	$\frac{43}{128}$	$\frac{417}{256}$	$\frac{451}{512}$	$\frac{745}{1024}$	$\frac{2971}{2048}$	$\frac{2929}{4096}$	$\frac{8243}{8192}$

Theorem 1 can be used to obtain the Binet formula of generalized Blaise numbers. Using these (the above) roots and the recurrence relation, Binet's formula of generalized Blaise numbers can be given as follows:

**THEOREM 9.** *(Four Distinct Roots Case:  $\alpha \neq \beta \neq \gamma \neq \delta = 1$ ) For all integers  $n$ , Binet's formula of generalized Blaise numbers is*

$$W_n = \frac{(\alpha W_3 - \alpha(1 - \alpha)W_2 + (-\alpha^2 + 2)W_1 - 2W_0)\alpha^n}{2\alpha^2 + 4\alpha - 6} + \frac{(\beta W_3 - \beta(1 - \beta)W_2 + (-\beta^2 + 2)W_1 - 2W_0)\beta^n}{2\beta^2 + 4\beta - 6} \\ + \frac{(\gamma W_3 - \gamma(1 - \gamma)W_2 + (-\gamma^2 + 2)W_1 - 2W_0)\gamma^n}{2\gamma^2 + 4\gamma - 6} - \frac{W_3 - W_1 - 2W_0}{2}$$

Blaise and Blaise-Lucas numbers can be expressed using Binet's formulas as follows:

**COROLLARY 10.** *(Four Distinct Roots Case:  $\alpha \neq \beta \neq \gamma \neq \delta = 1$ ) For all integers  $n$ , Binet's formulas of Blaise and Blaise-Lucas numbers are*

$$B_n = \frac{\alpha^{n+3}}{2\alpha^2 + 4\alpha - 6} + \frac{\beta^{n+3}}{2\beta^2 + 4\beta - 6} + \frac{\gamma^{n+3}}{2\gamma^2 + 4\gamma - 6} - \frac{1}{2}$$

and

$$C_n = \alpha^n + \beta^n + \gamma^n + 1$$

respectively.

Note that for all integers  $n$ , adjusted Jacobsthal-Padovan, Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas), Jacobsthal-Padovan, and modified Jacobsthal-Padovan numbers can be expressed using Binet's formulas as

$$K_n = \frac{1}{(\alpha - \beta)(\alpha - \gamma)}\alpha^{n+1} + \frac{1}{(\beta - \alpha)(\beta - \gamma)}\beta^{n+1} + \frac{1}{(\gamma - \alpha)(\gamma - \beta)}\gamma^{n+1}, \\ L_n = \alpha^n + \beta^n + \gamma^n, \\ M_n = \frac{(3\alpha + 1)}{(\alpha - \beta)(\alpha - \gamma)}\alpha^{n+1} + \frac{(3\beta + 1)}{(\beta - \alpha)(\beta - \gamma)}\beta^{n+1} + \frac{(3\gamma + 1)}{(\gamma - \alpha)(\gamma - \beta)}\gamma^{n+1}, \\ Q_n = \frac{(\alpha + 1)}{(\alpha - \beta)(\alpha - \gamma)}\alpha^{n+1} + \frac{(\beta + 1)}{(\beta - \alpha)(\beta - \gamma)}\beta^{n+1} + \frac{(\gamma + 1)}{(\gamma - \alpha)(\gamma - \beta)}\gamma^{n+1},$$

respectively, see Soykan [38] for more details. So, by using Binet's formulas of Blaise, Blaise-Lucas and Adjusted Jacobsthal-Padovan, Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas), Jacobsthal-Padovan, and modified Jacobsthal-Padovan numbers, (or by using mathematical induction), we get the following Lemma which contains many identities:

**LEMMA 11.** *For all integers  $n$ , the following equalities (identities) are true:*

**(a):**

- $K_{n+1} = B_{n+1} - B_n.$
- $2K_n = B_{n+3} - B_{n+2} - B_{n+1} + B_n.$
- $2B_{n+4} = 5K_{n+2} + 9K_{n+1} + 6K_n - 1.$
- $2B_n = K_{n+2} + K_{n+1} + 2K_n - 1.$

- $2K_n = -B_{n+2} + 3B_n + 1.$

(b):

- $52K_{n+3} = 12C_{n+3} + 17C_{n+2} - 11C_{n+1} - 18C_n.$

- $52K_n = C_{n+3} - 2C_{n+1} + 9C_{n+2} - 8C_n.$

- $C_{n+4} = 6K_{n+2} + 2K_{n+1} + 4K_n + 1.$

- $2C_n = -K_{n+2} + 6K_{n+1} + K_n + 2.$

- $52K_n = 9C_{n+2} - C_{n+1} - 6C_n - 2.$

- $C_{n+1} + 6C_n = 18K_{n+1} + 2K_n + 7.$

(c):

- $L_{n+3} = 2B_{n+2} + 4B_{n+1} - 6B_n.$

- $4L_n = B_{n+3} - 3B_{n+2} + 13B_{n+1} - 11B_n.$

- $52B_{n+4} = 30L_{n+2} + 53L_{n+1} + 58L_n - 26.$

- $52B_n = 2L_n + 9L_{n+1} + 10L_{n+2} - 26.$

- $4L_n = -3B_{n+2} + 14B_{n+1} - 9B_n + 1.$

- $4(10B_{n+1} - 9B_n) = 14L_n + 3L_{n+1} - 2.$

(d):

- $2L_{n+3} = 3C_{n+3} - C_{n+1} - 2C_n.$

- $2L_n = C_{n+3} - C_{n+1}.$

- $C_{n+4} = L_{n+2} + 2L_{n+1} + 1.$

- $C_n = L_n + 1.$

- $L_n = C_n - 1.$

(e):

- $Q_{n+3} = B_{n+3} + B_{n+1} - 2B_n.$

- $2Q_n = B_{n+3} - B_{n+2} + B_{n+1} - B_n.$

- $2B_{n+4} = Q_{n+2} + 4Q_{n+1} + 4Q_n - 1.$

- $2B_n = Q_{n+2} - 1.$

- $2Q_n = -B_{n+2} + 2B_{n+1} + B_n + 1.$

- $2B_{n+1} = Q_{n+1} + 2Q_n - 1.$

(f):

- $52Q_{n+3} = 41C_{n+3} + 18C_{n+2} - 17C_{n+1} - 42C_n.$

- $52Q_n = 11C_{n+3} + 8C_{n+2} - 9C_{n+1} - 10C_n.$

- $C_{n+4} = 4Q_{n+2} + 2Q_{n+1} - 4Q_n + 1.$

- $2C_n = -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2.$

- $26Q_n = 4C_{n+2} + C_{n+1} + 6C_n - 11.$

- $3C_{n+1} + 2C_n = 8Q_{n+1} - 2Q_n + 5.$

(g):

- $M_{n+3} = B_{n+3} + 2B_{n+2} + 3B_{n+1} - 6B_n$ .
- $2M_n = B_{n+3} - B_{n+2} + 5B_{n+1} - 5B_n$ .
- $46B_{n+4} = 16M_{n+2} + 33M_{n+1} + 42M_n - 23$ .
- $46B_n = 8M_{n+2} + 5M_{n+1} - 2M_n - 23$ .
- $2M_n = -B_{n+2} + 6B_{n+1} - 3B_n + 1$ .
- $2(8B_{n+1} - 5B_n) = M_{n+1} + 6M_n - 3$ .

(h):

- $52M_{n+3} = 99C_{n+3} + 20C_{n+2} - 29C_{n+1} - 90C_n$ .
- $52M_n = 31C_{n+3} + 6C_{n+2} - 23C_{n+1} - 14C_n$ .
- $23C_{n+4} = 18M_{n+2} + 40M_{n+1} - 16M_n + 23$ .
- $46C_n = -5M_{n+2} - 6M_{n+1} + 53M_n + 46$ .
- $26M_n = 3C_{n+2} + 4C_{n+1} + 24C_n - 31$ .
- $6C_n - 5C_{n+1} = 8M_n - 6M_{n+1} + 1$ .

Proof. We only prove  $K_{n+1} = B_{n+1} - B_n$  by using Binet's formulas of  $K_n$  and  $B_n$  as the others can be proved similarly. By using Binet's formulas, we get

$$\begin{aligned}
B_{n+1} - B_n &= \frac{\alpha^{n+4}}{2\alpha^2 + 4\alpha - 6} - \frac{\alpha^{n+3}}{2\alpha^2 + 4\alpha - 6} + \frac{\beta^{n+4}}{2\beta^2 + 4\beta - 6} - \frac{\beta^{n+3}}{2\beta^2 + 4\beta - 6} + \frac{\gamma^{n+4}}{2\gamma^2 + 4\gamma - 6} - \frac{\gamma^{n+3}}{2\gamma^2 + 4\gamma - 6} \\
&= \frac{1}{2} \frac{\alpha^{n+3}}{\alpha + 3} + \frac{1}{2} \frac{\beta^{n+3}}{\beta + 3} + \frac{1}{2} \frac{\gamma^{n+3}}{\gamma + 3} \\
&= \frac{1}{(\alpha - \beta)(\alpha - \gamma)} \alpha^{n+2} + \frac{1}{(\beta - \alpha)(\beta - \gamma)} \beta^{n+2} + \frac{1}{(\gamma - \alpha)(\gamma - \beta)} \gamma^{n+2} \\
&= K_{n+1}
\end{aligned}$$

where

$$\begin{aligned}
\frac{1}{2} \frac{\alpha}{\alpha + 3} &= \frac{1}{(\alpha - \beta)(\alpha - \gamma)}, \\
\frac{1}{2} \frac{\beta}{\beta + 3} &= \frac{1}{(\beta - \alpha)(\beta - \gamma)}, \\
\frac{1}{2} \frac{\gamma}{\gamma + 3} &= \frac{1}{(\gamma - \alpha)(\gamma - \beta)}. \quad \square
\end{aligned}$$

Next, we give the ordinary generating function  $\sum_{n=0}^{\infty} W_n z^n$  of the sequence  $W_n$ .

LEMMA 12. Suppose that  $f_{W_n}(z) = \sum_{n=0}^{\infty} W_n z^n$  is the ordinary generating function of the generalized Blaise sequence  $\{W_n\}$ . Then,  $\sum_{n=0}^{\infty} W_n z^n$  is given by

$$\sum_{n=0}^{\infty} W_n z^n = \frac{W_0 + (W_1 - W_0)z + (W_2 - W_1 - W_0)z^2 + (W_3 - W_2 - W_1 - W_0)z^3}{1 - z - z^2 - z^3 + 2z^4}.$$

Proof. Take  $r = 1, s = 1, t = 1, u = -2$  in Lemma 3.

The previous lemma gives the following results as particular examples.

COROLLARY 13. *Generating functions of Blaise and Blaise-Lucas numbers are*

$$\sum_{n=0}^{\infty} B_n z^n = \frac{z}{1 - z - z^2 - z^3 + 2z^4},$$

$$\sum_{n=0}^{\infty} C_n z^n = \frac{4 - 3z - 2z^2 - z^3}{1 - z - z^2 - z^3 + 2z^4},$$

respectively.

Proof. Set  $W_n = B_n, B_0 = 0, B_1 = 1, B_2 = 1, B_3 = 2$  and  $W_n = C_n, C_0 = 4, C_1 = 1, C_2 = 3, C_3 = 7$  in the previous Lemma, respectively.  $\square$

### 3. Simson Formulas

Now, we present Simson's formula of generalized Blaise numbers.

THEOREM 14 (Simson's Formula of Generalized Blaise Numbers). *For all integers  $n$ , we have*

$$\begin{vmatrix} W_{n+3} & W_{n+2} & W_{n+1} & W_n \\ W_{n+2} & W_{n+1} & W_n & W_{n-1} \\ W_{n+1} & W_n & W_{n-1} & W_{n-2} \\ W_n & W_{n-1} & W_{n-2} & W_{n-3} \end{vmatrix} = 2^{n-3}(W_3 - W_1 - 2W_0)(W_3^3 + 2W_2^3 - W_1^3 - 4W_0^3 + (-3W_2 - W_1 + W_0)W_3^2 + 2(W_3 - W_1 - 3W_0)W_2^2 + (5W_3 + 3W_2 - 5W_0)W_1^2 + 4(W_2 + 2W_1)W_0^2 - 2W_1W_2W_3 + 4W_0W_2W_3 - 6W_0W_1W_3).$$

Proof. Take  $r = 1, s = 1, t = 1, u = -2$  in Theorem 4.  $\square$

The previous theorem gives the following results as particular examples.

COROLLARY 15. *For all integers  $n$ , the Simson's formulas of Blaise and Blaise-Lucas numbers are given*

as

$$\begin{vmatrix} B_{n+3} & B_{n+2} & B_{n+1} & B_n \\ B_{n+2} & B_{n+1} & B_n & B_{n-1} \\ B_{n+1} & B_n & B_{n-1} & B_{n-2} \\ B_n & B_{n-1} & B_{n-2} & B_{n-3} \end{vmatrix} = 2^{n-1},$$

$$\begin{vmatrix} C_{n+3} & C_{n+2} & C_{n+1} & C_n \\ C_{n+2} & C_{n+1} & C_n & C_{n-1} \\ C_{n+1} & C_n & C_{n-1} & C_{n-2} \\ C_n & C_{n-1} & C_{n-2} & C_{n-3} \end{vmatrix} = -13 \times 2^{n+2},$$

respectively.

Proof. Set  $W_n = B_n$ ,  $B_0 = 0, B_1 = 1, B_2 = 1, B_3 = 2$  and  $W_n = C_n$ ,  $C_0 = 4, C_1 = 1, C_2 = 3, C_3 = 7$  in the previous Theorem, respectively.  $\square$

#### 4. Some Identities

In this section, we obtain some identities of Blaise and Blaise-Lucas numbers. First, we can give a few basic relations between  $\{W_n\}$  and  $\{B_n\}$ .

LEMMA 16. *The following equalities are true:*

- (a):  $16W_n = (11W_0 + 11W_1 + 3W_2 - 9W_3)B_{n+5} + (7W_0 - 9W_1 - W_2 + 3W_3)B_{n+4} - (17W_0 + W_1 + 9W_2 - 11W_3)B_{n+3} - (33W_0 + 17W_1 - 7W_2 - 11W_3)B_{n+2}$ .
- (b):  $8W_n = (9W_0 + W_1 + W_2 - 3W_3)B_{n+4} - (3W_0 - 5W_1 + 3W_2 - W_3)B_{n+3} - (11W_0 + 3W_1 - 5W_2 - W_3)B_{n+2} - (11W_0 + 11W_1 + 3W_2 - 9W_3)B_{n+1}$ .
- (c):  $4W_n = (3W_0 + 3W_1 - W_2 - W_3)B_{n+3} - (W_0 + W_1 - 3W_2 + W_3)B_{n+2} - (W_0 + 5W_1 + W_2 - 3W_3)B_{n+1} - (9W_0 + W_1 + W_2 - 3W_3)B_n$ .
- (d):  $2W_n = (W_0 + W_1 + W_2 - W_3)B_{n+2} + (W_0 - W_1 - W_2 + W_3)B_{n+1} - (3W_0 - W_1 + W_2 - W_3)B_n - (3W_0 + 3W_1 - W_2 - W_3)B_{n-1}$ .
- (e):  $W_n = W_0B_{n+1} + (W_1 - W_0)B_n + (W_2 - W_1 - W_0)B_{n-1} + (W_3 - W_1 - W_2 - W_0)B_{n-2}$ .

Proof. Note that all the identities hold for all integers  $n$ . We prove (a). To show (a), writing

$$W_n = a \times B_{n+5} + b \times B_{n+4} + c \times B_{n+3} + d \times B_{n+2}$$

and solving the system of equations

$$W_0 = a \times B_5 + b \times B_4 + c \times B_3 + d \times B_2$$

$$W_1 = a \times B_6 + b \times B_5 + c \times B_4 + d \times B_3$$

$$W_2 = a \times B_7 + b \times B_6 + c \times B_5 + d \times B_4$$

$$W_3 = a \times B_8 + b \times B_7 + c \times B_6 + d \times B_5$$

we find that  $16a = 11W_0 + 11W_1 + 3W_2 - 9W_3$ ,  $16b = 7W_0 - 9W_1 - W_2 + 3W_3$ ,  $16c = 11W_3 - W_1 - 9W_2 - 17W_0$ ,  $16d = 7W_2 - 17W_1 - 33W_0 + 11W_3$ . The other equalities can be proved similarly.  $\square$

Note that all the identities in the above Lemma can be proved by induction as well.

Next, we present a few basic relations between  $\{W_n\}$  and  $\{C_n\}$ .

LEMMA 17. *The following equalities are true:*

- (a):  $104W_n = -(56W_0 + 27W_1 + 2W_2 - 33W_3)C_{n+5} - 2(5W_0 - 2W_1 - 4W_2 + W_3)C_{n+4} + (60W_0 + 15W_1 + 4W_2 - 27W_3)C_{n+3} + 2(55W_0 + 30W_1 - 5W_2 - 28W_3)C_{n+2}$ .
- (b):  $104W_n = -(66W_0 + 23W_1 - 6W_2 - 31W_3)C_{n+4} + 2(2W_0 - 6W_1 + W_2 + 3W_3)C_{n+3} + (54W_0 + 33W_1 - 12W_2 - 23W_3)C_{n+2} + 2(56W_0 + 27W_1 + 2W_2 - 33W_3)C_{n+1}$ .

$$\text{(c): } 104W_n = -(62W_0 + 35W_1 - 8W_2 - 37W_3)C_{n+3} - 2(6W_0 - 5W_1 + 3W_2 - 4W_3)C_{n+2} + (46W_0 + 31W_1 + 10W_2 - 35W_3)C_{n+1} + 2(66W_0 + 23W_1 - 6W_2 - 31W_3)C_n.$$

$$\text{(d): } 104W_n = -(74W_0 + 25W_1 - 2W_2 - 45W_3)C_{n+2} - 2(8W_0 + 2W_1 - 9W_2 - W_3)C_{n+1} + (70W_0 + 11W_1 - 4W_2 - 25W_3)C_n + 2(62W_0 + 35W_1 - 8W_2 - 37W_3)C_{n-1}.$$

$$\text{(e): } 104W_n = -(90W_0 + 29W_1 - 20W_2 - 47W_3)C_{n+1} - 2(2W_0 + 7W_1 + W_2 - 10W_3)C_n + (50W_0 + 45W_1 - 14W_2 - 29W_3)C_{n-1} + 2(74W_0 + 25W_1 - 2W_2 - 45W_3)C_{n-2}.$$

Now, we give a few basic relations between  $\{B_n\}$  and  $\{C_n\}$ .

LEMMA 18. *The following equalities are true:*

$$104B_n = 37C_{n+5} + 8C_{n+4} - 35C_{n+3} - 62C_{n+2},$$

$$104B_n = 45C_{n+4} + 2C_{n+3} - 25C_{n+2} - 74C_{n+1},$$

$$104B_n = 47C_{n+3} + 20C_{n+2} - 29C_{n+1} - 90C_n,$$

$$104B_n = 67C_{n+2} + 18C_{n+1} - 43C_n - 94C_{n-1},$$

$$104B_n = 85C_{n+1} + 24C_n - 27C_{n-1} - 134C_{n-2},$$

and

$$16C_n = B_{n+5} + 37B_{n+4} - 19B_{n+3} - 51B_{n+2},$$

$$8C_n = 19B_{n+4} - 9B_{n+3} - 25B_{n+2} - B_{n+1},$$

$$4C_n = 5B_{n+3} - 3B_{n+2} + 9B_{n+1} - 19B_n,$$

$$2C_n = B_{n+2} + 7B_{n+1} - 7B_n - 5B_{n-1},$$

$$C_n = 4B_{n+1} - 3B_n - 2B_{n-1} - B_{n-2}.$$

## 5. Relations Between Special Numbers

In this section, we present identities on Blaise, Blaise-Lucas numbers and adjusted Jacobsthal-Padovan, Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas), Jacobsthal-Padovan, and modified Jacobsthal-Padovan numbers. We know from Lemma 11 that

$$52B_n = 2L_n + 9L_{n+1} + 10L_{n+2} - 26,$$

$$2C_n = -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2.$$

Note also that from Lemma 16 and Lemma 17, we have the formulas of  $W_n$  as

$$\begin{aligned} 4W_n &= (3W_0 + 3W_1 - W_2 - W_3)B_{n+3} - (W_0 + W_1 - 3W_2 + W_3)B_{n+2} \\ &\quad - (W_0 + 5W_1 + W_2 - 3W_3)B_{n+1} - (9W_0 + W_1 + W_2 - 3W_3)B_n, \\ 104W_n &= -(62W_0 + 35W_1 - 8W_2 - 37W_3)C_{n+3} - 2(6W_0 - 5W_1 + 3W_2 - 4W_3)C_{n+2} \\ &\quad + (46W_0 + 31W_1 + 10W_2 - 35W_3)C_{n+1} + 2(66W_0 + 23W_1 - 6W_2 - 31W_3)C_n. \end{aligned}$$

Using the above identities, we obtain relation of generalized Blaise numbers in the following forms (in terms of Jacobsthal-Perrin (Jacobsthal-Perrin-Lucas) and Jacobsthal-Padovan numbers):

LEMMA 19. *For all integers  $n$ , we have the following identities:*

$$\begin{aligned} \text{(a): } 52W_n &= (4W_3 - 3W_2 + 5W_1 - 6W_0)L_{n+2} + (W_3 + 9W_2 - 2W_1 - 8W_0)L_{n+1} + 2(3W_3 + W_2 - 6W_1 + \\ &\quad 2W_0)L_n - 26W_3 + 26W_1 + 52W_0. \\ \text{(b): } 2W_n &= (W_1 - W_0)Q_{n+2} + (W_2 - W_1)Q_{n+1} + (W_3 - W_2 - W_1 + W_0)Q_n - W_3 + W_1 + 2W_0. \end{aligned}$$

## 6. On the Recurrence Properties of Generalized Blaise Sequence

Taking  $r = 1, s = 1, t = 1, u = -2$  in Theorem 5, we obtain the following Proposition.

PROPOSITION 20. *For  $n \in \mathbb{Z}$ , generalized Blaise numbers (the case  $r = 1, s = 1, t = 1, u = -2$ ) have the following identity:*

$$W_{-n} = \frac{2^{-n-1}}{3}(-6W_{3n} + 6C_n W_{2n} - 3C_n^2 W_n + 3C_{2n} W_n + W_0 C_n^3 + 2W_0 C_{3n} - 3W_0 C_n C_{2n}).$$

From the above Proposition 20 (or by taking  $G_n = B_n$  and  $H_n = C_n$  in (1.12) and (1.13) respectively), we have the following corollary which gives the connection between the special cases of generalized Blaise sequence at the positive index and the negative index: for Blaise and Blaise-Lucas numbers: take  $W_n = B_n$  with  $B_0 = 0, B_1 = 1, B_2 = 1, B_3 = 2$  and take  $W_n = C_n$  with  $C_0 = 4, C_1 = 1, C_2 = 3, C_3 = 7$ , respectively. Note that in this case  $H_n = C_n$ .

COROLLARY 21. *For  $n \in \mathbb{Z}$ , we have the following recurrence relations:*

(a): *Blaise sequence:*

$$B_{-n} = \frac{2^{-n-1}}{3}(-6B_{3n} + 6C_n B_{2n} - 3C_n^2 B_n + 3C_{2n} B_n).$$

(b): *Blaise-Lucas sequence:*

$$C_{-n} = \frac{2^{-n-1}}{3}(C_n^3 + 2C_{3n} - 3C_{2n} C_n).$$

We can also present the formulas of  $B_{-n}$  and  $C_{-n}$  in the following forms.

COROLLARY 22. *For  $n \in \mathbb{Z}$ , we have the following recurrence relations:*

$$(a): B_{-n} = \frac{2^{-n-5}}{3}(-96B_{3n} + 24(5B_{n+3} - 3B_{n+2} + 9B_{n+1} - 19B_n)B_{2n} - 3(5B_{n+3} - 3B_{n+2} + 9B_{n+1} - 19B_n)^2B_n + 12(5B_{2n+3} - 3B_{2n+2} + 9B_{2n+1} - 19B_{2n})B_n).$$

(b):

$$(i): 2^{n+1}B_{-n} = -K_n^2 - K_{n-1}^2 - 2K_{n-2}^2 + (K_{n+2} - 6K_{n+1} - 6K_{n-1} + 2K_{n-2})K_n + (K_{n+1} - 12K_{n-2})K_{n-1} + 2K_{2n} + 2K_{2n-2} + 4K_{2n-4} - 2^n.$$

$$(ii): 2^{n+2}C_{-n} = -K_n^2 - 12K_{n-1}^2 + 4K_{n-2}^2 + (K_{n+2} - 6K_{n+1} - 72K_{n-1} - 4K_{n-2})K_n + 12(K_{n+1} + 2K_{n-2})K_{n-1} + 2K_{2n} + 24K_{2n-2} - 8K_{2n-4} + 2^{n+2}.$$

(c):

$$(i): 13B_{-n} = \frac{1}{2^{n+2}}(L_n^2 + 9L_{n-1}^2 + 20L_{n-2}^2 - L_{2n} - 9L_{2n-2} - 20L_{2n-4} - 13 \times 2^{n+1}).$$

$$(ii): 2^{n+1}C_{-n} = L_n^2 - L_{2n} + 2^{n+1}.$$

(d):

$$(i): 2^{n+2}B_{-n} = 9Q_n^2 + 4Q_{n-1}^2 + 21Q_{n-2}^2 + 6Q_{2n-2} - 4Q_{2n-3} - 6Q_{2n-4} - 12Q_nQ_{n-1} - 30Q_nQ_{n-2} + 20Q_{n-1}Q_{n-2} - 2^{n+1}.$$

$$(ii): 2^{n+4}C_{-n} = 63Q_{n+2}^2 + 64Q_{n+1}^2 + 55Q_n^2 + 36Q_{n-1}^2 - 252Q_{n-2}^2 + 2(-105Q_{n+2} + 46Q_{n+1} + 112Q_{n-1} + 180Q_{n-2})Q_n - 120(Q_{n+1} + 2Q_{n-2})Q_{n-1} - 84Q_{n+1}Q_{n+2} + 42Q_{2n+2} - 28Q_{2n+1} - 18Q_{2n} - 16Q_{2n-1} - 96Q_{2n-2} + 48Q_{2n-3} + 72Q_{2n-4} + 2^{n+4}.$$

(e):

$$(i): 12167B_{-n} = \frac{1}{2^{n+3}}(-75M_{n+2}^2 + 267M_{n+1}^2 - 1811M_n^2 + 19483M_{n-1}^2 + 56816M_{n-2}^2 + 2(+565M_{n+2} + 1128M_{n+1} - 1950M_{n-1} - 9040M_{n-2})M_n - 226(25M_{n+1} + 96M_{n-2})M_{n-1} - 180M_{n+1}M_{n+2} + 6532M_{2n} + 4140M_{2n-1} - 828M_{2n+1} - 4370M_{2n-2} - 690M_{2n+2} + 13248M_{2n-3} - 49312M_{2n-4} - 12167 \times 2^{n+2}).$$

$$(ii): 12167C_{-n} = \frac{1}{2^{n+4}}(3975M_{n+2}^2 + 4824M_{n+1}^2 + 185407M_n^2 - 44772M_{n-1}^2 - 71020M_{n-2}^2 + 2(-29945M_{n+2} - 37014M_{n+1} + 6336M_{n-1} + 11300M_{n-2})M_n + 13560(M_{n+1} + 2M_{n-2})M_{n-1} + 9540M_{n+1}M_{n+2} + 36570M_{2n+2} + 43884M_{2n+1} - 171626M_{2n} - 9936M_{2n-1} + 23184M_{2n-2} - 16560M_{2n-3} + 61640M_{2n-4} + 12167 \times 2^{n+4}).$$

Proof. We use the identities, see Soykan [37],

$$K_{-n} = \frac{1}{2^{n+1}}(-K_n^2 + 2K_{2n} + K_{n+2}K_n - 6K_{n+1}K_n),$$

$$L_{-n} = \frac{1}{2^{n+1}}(L_n^2 - L_{2n}),$$

and

$$Q_{-n} = \frac{1}{2^{n+3}}(9Q_{n+2}^2 + 4Q_{n+1}^2 + 21Q_n^2 + 6Q_{2n+2} - 4Q_{2n+1} - 6Q_{2n} - 12Q_{n+2}Q_{n+1} - 30Q_{n+2}Q_n + 20Q_{n+1}Q_n),$$

$$M_{-n} = \frac{1}{529 \times 2^{n+3}}(108M_{n+1}^2 + 75M_{n+2}^2 + 3551M_n^2 + 690M_{2n+2} + 828M_{2n+1} - 3082M_{2n} + 180M_{n+2}M_{n+1} - 1130M_{n+2}M_n - 1356M_{n+1}M_n).$$

We also use the identities

$$\begin{aligned}
2B_n &= K_{n+2} + K_{n+1} + 2K_n - 1, \\
2C_n &= -K_{n+2} + 6K_{n+1} + K_n + 2, \\
52B_n &= 2L_n + 9L_{n+1} + 10L_{n+2} - 26, \\
C_n &= L_n + 1, \\
2B_n &= Q_{n+2} - 1, \\
2C_n &= -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2, \\
46B_n &= 8M_{n+2} + 5M_{n+1} - 2M_n - 23, \\
46C_n &= -5M_{n+2} - 6M_{n+1} + 53M_n + 46.
\end{aligned}$$

(a): By using the identity  $4C_n = 5B_{n+3} - 3B_{n+2} + 9B_{n+1} - 19B_n$  and Corollary 18, (or by using Corollary 6 (a)), we obtain (a).

(b): Since  $2B_n = K_{n+2} + K_{n+1} + 2K_n - 1$ ,  $2C_n = -K_{n+2} + 6K_{n+1} + K_n + 2$  and  $K_{-n} = \frac{1}{2^{n+1}}(-K_n^2 + 2K_{2n} + K_{n+2}K_n - 6K_{n+1}K_n)$ , we get (b)

(c): Since  $52B_n = 2L_n + 9L_{n+1} + 10L_{n+2} - 26$ ,  $C_n = L_n + 1$  and  $L_{-n} = \frac{1}{2^{n+1}}(L_n^2 - L_{2n})$ , we obtain (c).

(d): Since  $2B_n = Q_{n+2} - 1$ ,  $2C_n = -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2$  and  $Q_{-n} = \frac{1}{2^{n+3}}(9Q_{n+2}^2 + 4Q_{n+1}^2 + 21Q_n^2 + 6Q_{2n+2} - 4Q_{2n+1} - 6Q_{2n} - 12Q_{n+2}Q_{n+1} - 30Q_{n+2}Q_n + 20Q_{n+1}Q_n)$ , we get (d).

(e): Since  $46B_n = 8M_{n+2} + 5M_{n+1} - 2M_n - 23$ ,  $46C_n = -5M_{n+2} - 6M_{n+1} + 53M_n + 46$  and  $M_{-n} = \frac{1}{529 \times 2^{n+3}}(108M_{n+1}^2 + 75M_{n+2}^2 + 3551M_n^2 + 690M_{2n+2} + 828M_{2n+1} - 3082M_{2n} + 180M_{n+2}M_{n+1} - 1130M_{n+2}M_n - 1356M_{n+1}M_n)$ , we obtain (e).  $\square$

## 7. Sum Formulas

The following Corollary gives sum formulas of Jacobsthal-Padovan numbers.

**COROLLARY 23.** *For  $n \geq 0$  we have the following formulas:*

$$\begin{aligned}
\text{(a): } \sum_{k=0}^n Q_k &= \frac{1}{2}(Q_{n+3} + Q_{n+2} - 2). \\
\text{(b): } \sum_{k=0}^n Q_{2k} &= \frac{1}{2}(Q_{2n+1} + 2Q_{2n} - 1). \\
\text{(c): } \sum_{k=0}^n Q_{2k+1} &= \frac{1}{2}(Q_{2n+2} + 2Q_{2n+1} - 1).
\end{aligned}$$

*Proof.* It is given in Soykan [38].  $\square$

The following Corollary presents sum formulas of Blaise and Blaise-Lucas numbers.

**COROLLARY 24.** *For  $n \geq 0$ , Blaise and Blaise-Lucas numbers have the following properties (in terms of Jacobsthal-Padovan numbers):*

$$\text{(a):}$$

(i):  $\sum_{k=0}^n B_k = \frac{1}{4}(3Q_{n+2} + 3Q_{n+1} + 2Q_n - 2(n+4)).$

(ii):  $\sum_{k=0}^n B_{2k} = \frac{1}{4}(2Q_{2n+2} + Q_{2n+1} + 2Q_{2n} - 2n - 5).$

(iii):  $\sum_{k=0}^n B_{2k+1} = \frac{1}{4}(Q_{2n+2} + 4Q_{2n+1} + 4Q_{2n} - 2n - 5).$

(b):

(i):  $\sum_{k=0}^n C_k = Q_{n+1} + 3Q_n + n.$

(ii):  $\sum_{k=0}^n C_{2k} = -Q_{2n+2} + 2Q_{2n+1} + 2Q_{2n} + n + 1.$

(iii):  $\sum_{k=0}^n C_{2k+1} = 2Q_{2n+2} + Q_{2n+1} - 2Q_{2n} + n.$

Proof. The proof follows from Corollary 23 and the identities

$$2B_n = Q_{n+2} - 1,$$

$$2C_n = -3Q_{n+2} + 2Q_{n+1} + 7Q_n + 2. \quad \square$$

### 8. Matrices and Identities Related With Generalized Blaise Numbers

If we define the square matrix  $A$  of order 4 as

$$A = \begin{pmatrix} 1 & 1 & 1 & -2 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$$

and also define

$$B_n = \begin{pmatrix} B_{n+1} & B_n + B_{n-1} - 2B_{n-2} & B_n - 2B_{n-1} & -2B_n \\ B_n & B_{n-1} + B_{n-2} - 2B_{n-3} & B_{n-1} - 2B_{n-2} & -2B_{n-1} \\ B_{n-1} & B_{n-2} + B_{n-3} - 2B_{n-4} & B_{n-2} - 2B_{n-3} & -2B_{n-2} \\ B_{n-2} & B_{n-3} + B_{n-4} - 2B_{n-5} & B_{n-3} - 2B_{n-4} & -2B_{n-3} \end{pmatrix}$$

and

$$U_n = \begin{pmatrix} W_{n+1} & W_n + W_{n-1} - 2W_{n-2} & W_n - 2W_{n-1} & -2W_n \\ W_n & W_{n-1} + W_{n-2} - 2W_{n-3} & W_{n-1} - 2W_{n-2} & -2W_{n-1} \\ W_{n-1} & W_{n-2} + W_{n-3} - 2W_{n-4} & W_{n-2} - 2W_{n-3} & -2W_{n-2} \\ W_{n-2} & W_{n-3} + W_{n-4} - 2W_{n-5} & W_{n-3} - 2W_{n-4} & -2W_{n-3} \end{pmatrix}.$$

then we get the following Theorem.

**THEOREM 25.** *For all integers  $m, n$ , we have*

(a):  $B_n = A^n$ , i.e.,

$$\begin{pmatrix} 1 & 1 & 1 & -2 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}^n = \begin{pmatrix} B_{n+1} & B_n + B_{n-1} - 2B_{n-2} & B_n - 2B_{n-1} & -2B_n \\ B_n & B_{n-1} + B_{n-2} - 2B_{n-3} & B_{n-1} - 2B_{n-2} & -2B_{n-1} \\ B_{n-1} & B_{n-2} + B_{n-3} - 2B_{n-4} & B_{n-2} - 2B_{n-3} & -2B_{n-2} \\ B_{n-2} & B_{n-3} + B_{n-4} - 2B_{n-5} & B_{n-3} - 2B_{n-4} & -2B_{n-3} \end{pmatrix}.$$

$$(b): U_1 A^n = A^n U_1.$$

$$(c): U_{n+m} = U_n B_m = B_m U_n.$$

**Proof.** Take  $r = 1, s = 1, t = 1, u = -2$  in Theorem 7.  $\square$

Using the above last Theorem and the identity

$$2B_n = Q_{n+2} - 1,$$

we obtain the following identity for Jacobsthal-Padovan numbers.

**COROLLARY 26.** *For all integers  $n$ , we have the following formula for Jacobsthal-Padovan numbers:*

$$A^n = \frac{1}{2} \begin{pmatrix} Q_{n+3} - 1 & Q_{n+4} - Q_{n+3} & -Q_{n+4} + 2Q_{n+2} + 1 & -2Q_{n+2} + 2 \\ Q_{n+2} - 1 & Q_{n+3} - Q_{n+2} & -Q_{n+3} + 2Q_{n+1} + 1 & -2Q_{n+1} + 2 \\ Q_{n+1} - 1 & Q_{n+2} - Q_{n+1} & -Q_{n+2} + 2Q_n + 1 & -2Q_n + 2 \\ Q_n - 1 & Q_{n+1} - Q_n & -Q_{n+1} + 2Q_{n-1} + 1 & -2Q_{n-1} + 2 \end{pmatrix}.$$

Next, we present an identity for  $W_{n+m}$ .

**THEOREM 27.** *For all integers  $m, n$ , we have*

$$W_{n+m} = W_n B_{m+1} + W_{n-1} (B_m + B_{m-1} - 2B_{m-2}) + W_{n-2} (B_m - 2B_{m-1}) - 2W_{n-3} B_m.$$

**Proof.** Take  $r = 1, s = 1, t = 1, u = -2$  in Theorem 8.  $\square$

As particular cases of the above theorem, we give identities for  $B_{n+m}$  and  $C_{n+m}$ .

**COROLLARY 28.** *For all integers  $m, n$ , we have*

$$B_{n+m} = B_n B_{m+1} + B_{n-1} (B_m + B_{m-1} - 2B_{m-2}) + B_{n-2} (B_m - 2B_{m-1}) - 2B_{n-3} B_m,$$

$$C_{n+m} = C_n B_{m+1} + C_{n-1} (B_m + B_{m-1} - 2B_{m-2}) + C_{n-2} (B_m - 2B_{m-1}) - 2C_{n-3} B_m.$$

## 9. Conclusions

Sequences have been fascinating topic for mathematicians for centuries. The Fibonacci and Lucas sequences are sources of many nice and interesting identities. For example, in [1], authors study on the solutions of the connection problems between Fermat and generalized Fibonacci polynomials. For rich applications of these second order sequences in science and nature, one can see the citations in [19].

As a fourth order sequence, we introduce the generalized Blaise sequence (and it's two special cases, namely, Blaise and Blaise-Lucas sequences) and we present Binet's formulas, generating functions, Simson formulas, the sum formulas, some identities, recurrence properties and matrices for these sequences.

We have shown that there are close relations between Blaise, Blaise-Lucas numbers (which are fourth order linear recurrences) and special third order linear recurrences (numbers), namely Jacobsthal-Padovan, Jacobsthal-Perrin, adjusted Jacobsthal-Padovan, modified Jacobsthal-Padovan numbers.

Linear recurrence relations (sequences) have many applications. We now present one of them. The ratio of two consecutive Padovan numbers converges to the plastic ratio,  $\alpha_P$  (which is given in (9.1) below), which have many applications to such as architecture, see [21]. Padovan numbers is defined by the third-order recurrence relations

$$P_{n+3} = P_{n+1} + P_n, \quad P_0 = 1, P_1 = 1, P_2 = 1.$$

The characteristic equation associated with Padovan sequence is  $x^3 - x - 1 = 0$  with roots  $\alpha, \beta$  and  $\gamma$  in which

$$\alpha = \left(\frac{1}{2} + \sqrt{\frac{23}{108}}\right)^{1/3} + \left(\frac{1}{2} - \sqrt{\frac{23}{108}}\right)^{1/3} \simeq 1.32471795724 \quad (9.1)$$

is called plastic number (or plastic ratio or plastic constant or silver number) and

$$\lim_{n \rightarrow \infty} \frac{P_{n+1}}{P_n} = \alpha.$$

The plastic number is used in art and architecture. Richard Padovan studied on plastic number in Architecture and Mathematics in [25, 26].

Next, we list applications of sequences which are linear recurrence relations.

First, we present some applications of second order sequences.

- For the applications of Gaussian Fibonacci and Gaussian Lucas numbers to Pauli Fibonacci and Pauli Lucas quaternions, see [2].
- For the application of Pell Numbers to the solutions of three-dimensional difference equation systems, see [5].
- For the application of Jacobsthal numbers to special matrices, see [58].
- For the application of generalized k-order Fibonacci numbers to hybrid quaternions, see [13].
- For the applications of Fibonacci and Lucas numbers to Split Complex Bi-Periodic numbers, see [55].
- For the applications of generalized bivariate Fibonacci and Lucas polynomials to matrix polynomials, see [56].
- For the applications of generalized Fibonacci numbers to binomial sums, see [54].
- For the application of generalized Jacobsthal numbers to hyperbolic numbers, see [39].
- For the application of generalized Fibonacci numbers to dual hyperbolic numbers, see [40].
- For the application of Laplace transform and various matrix operations to the characteristic polynomial of the Fibonacci numbers, see [9].
- For the application of Generalized Fibonacci Matrices to Cryptography, see [27].
- For the application of higher order Jacobsthal numbers to quaternions, see [24].
- For the application of Fibonacci and Lucas Identities to Toeplitz-Hessenberg matrices, see [12].
- For the applications of Fibonacci numbers to lacunary statistical convergence, see [4].

- For the applications of Fibonacci numbers to lacunary statistical convergence in intuitionistic fuzzy normed linear spaces, see [17].
- For the applications of Fibonacci numbers to ideal convergence on intuitionistic fuzzy normed linear spaces, see [18].

We now present some other applications of third order sequences.

- For the applications of third order Jacobsthal numbers and Tribonacci numbers to quaternions, see [7] and [6], respectively.
- For the application of Tribonacci numbers to special matrices, see [57].
- For the applications of Padovan numbers and Tribonacci numbers to coding theory, see [28] and [3], respectively.
- For the application of Pell-Padovan numbers to groups, see [10].
- For the application of adjusted Jacobsthal-Padovan numbers to the exact solutions of some difference equations, see [14].
- For the application of Gaussian Tribonacci numbers to various graphs, see [53].
- For the application of third-order Jacobsthal numbers to hyperbolic numbers, see [11].
- For the application of Narayan numbers to finite groups see [20].
- For the application of generalized third-order Jacobsthal sequence to binomial transform, see [41].
- For the application of generalized Generalized Padovan numbers to Binomial Transform, see [42].
- For the application of generalized Tribonacci numbers to Gaussian numbers, see [43].
- For the application of generalized Tribonacci numbers to Sedenions, see [44].
- For the application of Tribonacci and Tribonacci-Lucas numbers to matrices, see [45].
- For the application of generalized Tribonacci numbers to circulant matrix, see [46].

Next, we list some applications of fourth order sequences.

- For the application of Tetranacci and Tetranacci-Lucas numbers to quaternions, see [47].
- For the application of generalized Tetranacci numbers to Gaussian numbers, see [48].
- For the application of Tetranacci and Tetranacci-Lucas numbers to matrices, see [49].
- For the application of generalized Tetranacci numbers to binomial transform, see [50].

We now present some applications of fifth order sequences.

- For the application of Pentanacci numbers to matrices, see [30].
- For the application of generalized Pentanacci numbers to quaternions, see [51].
- For the application of generalized Pentanacci numbers to binomial transform, see [52].

## References

- [1] Abd-Elhameed, W.M., Youssri, Y.H., Solutions of the Connection Problems Between Fermat and Generalized Fibonacci Polynomials, JP Journal of Algebra, Number Theory and Applications, 38(4), 349-362, 2016. <http://dx.doi.org/10.17654/NT038040349>

- [2] Azak, A.Z., Pauli Gaussian Fibonacci and Pauli Gaussian Lucas Quaternions. *Mathematics*, 2022, 10, 4655. <https://doi.org/10.3390/math10244655>
- [3] Basu, M., Das, M., Tribonacci Matrices and a New Coding Theory, *Discrete Mathematics, Algorithms and Applications*, 6(1), 1450008, (17 pages), 2014.
- [4] Bilgin, N.G., Fibonacci Lacunary Statistical Convergence of Order  $\gamma$  in IFNLS, *International Journal of Advances in Applied Mathematics and Mechanics*, 8(4), 28-36, 2021.
- [5] Büyütk, H., Taşkara, N., On The Solutions of Three-Dimensional Difference Equation Systems Via Pell Numbers, *European Journal of Science and Technology, Special Issue 34*, 433-440, 2022.
- [6] Cerda-Morales, G., On a Generalization of Tribonacci Quaternions, *Mediterranean Journal of Mathematics* 14:239, 1-12, 2017.
- [7] Cerda-Morales, G., Identities for Third Order Jacobsthal Quaternions, *Advances in Applied Clifford Algebras* 27(2), 1043–1053, 2017.
- [8] Deveci, Ö., The Jacobsthal-Padovan p-Sequences and Their Applications, *Proceedings of Romanian Academy, Series A*, 20(3), 215–224, 2019.
- [9] Deveci, Ö., Shannon, A.G., On Recurrence Results From Matrix Transforms, *Notes on Number Theory and Discrete Mathematics*, 28(4), 589–592, 2022. DOI: 10.7546/nntdm.2022.28.4.589-592
- [10] Deveci, Ö., Shannon, A.G., Pell–Padovan-Circulant Sequences and Their Applications, *Notes on Number Theory and Discrete Mathematics*, 23(3), 100–114, 2017.
- [11] Dikmen, C.M., Altınsoy, M., On Third Order Hyperbolic Jacobsthal Numbers, *Konuralp Journal of Mathematics*, 10(1), 118-126, 2022.
- [12] Goy, T., Shattuck, M., Fibonacci and Lucas Identities from Toeplitz-Hessenberg Matrices, *Appl. Appl. Math.*, 14(2), 699–715, 2019.
- [13] Gül, K., Generalized k-Order Fibonacci Hybrid Quaternions, *Erzincan University Journal of Science and Technology*, 15(2), 670-683, 2022. DOI: 10.18185/erzifbed.1132164
- [14] Göcen, M., The Exact Solutions of Some Difference Equations Associated with Adjusted Jacobsthal-Padovan Numbers, *Kırklareli University Journal of Engineering and Science* 8(1), 1-14, 2022. DOI: 10.34186/klujes.1078836
- [15] Hathiwala, G.S., Shah, D.V., Binet–Type Formula For The Sequence of Tetranacci Numbers by Alternate Methods, *Mathematical Journal of Interdisciplinary Sciences* 6(1) (2017), 37–48.
- [16] Howard F.T., Saidak, F., Zhou’s Theory of Constructing Identities, *Congress Numer.* 200 (2010), 225-237.
- [17] Kişi, Ö., Tuzcuoglu, I., Fibonacci Lacunary Statistical Convergence in Intuitionistic Fuzzy Normed Linear Spaces, *Journal of Progressive Research in Mathematics* 16(3), 3001-3007, 2020.
- [18] Kişi, Ö., Debnath, P., Fibonacci Ideal Convergence on Intuitionistic Fuzzy Normed Linear Spaces, *Fuzzy Information and Engineering*, 1-13, 2022. <https://doi.org/10.1080/16168658.2022.2160226>
- [19] Koshy, T., *Fibonacci and Lucas Numbers with Applications*. Wiley-Interscience. New York, 2001.
- [20] Kuloglu, B., Özkan, E., Shannon, A.G., The Narayana Sequence in Finite Groups, *Fibonacci Quarterly*. 60(5), 212–221, 2022.
- [21] Marohnić, L., Strmečki T., Plastic Number: Construction and Applications, *Advanced Research in Scientific Areas*, 3(7), 1523-1528, 2012.
- [22] Melham, R.S., Some Analogs of the Identity  $F_n^2 + F_{n+1}^2 = F_{2n+1}^2$ , *Fibonacci Quarterly*, (1999), 305-311.
- [23] Natividad, L.R., On Solving Fibonacci-Like Sequences of Fourth, Fifth and Sixth Order, *International Journal of Mathematics and Computing*, 3(2) (2013), 38-40.
- [24] Özkan, E., Uysal, M., On Quaternions with Higher Order Jacobsthal Numbers Components, *Gazi University Journal of Science*, 36(1), 336-347, 2023. DOI: 10.35378/gujs.1002454

- [25] Padovan, R., Dom Hans van Der Laan and the Plastic Number. In: Williams K., Ostwald M. (eds) *Architecture and Mathematics from Antiquity to the Future*. Birkhäuser, Cham 407-419, 2015, [https://doi.org/10.1007/978-3-319-00143-2\\_27](https://doi.org/10.1007/978-3-319-00143-2_27). (First published as: Richard Padovan , “Dom Hans van Der Laan and the Plastic Number”, pp. 181–193 in *Nexus IV: Architecture and Mathematics*, Kim Williams and Jose Francisco Rodrigues, eds. Fucecchio (Florence): Kim Williams Books, 2002.
- [26] Padovan, R., *Dom Hans van der Laan: Modern Primitive*, Architectura and Natura Press, 1994. <http://www.nexusjournal.com/conferences/N2002-Padovan.html>
- [27] Prasad, K., Mahato, H., Cryptography Using Generalized Fibonacci Matrices with Affine-Hill Cipher, *Journal of Discrete Mathematical Sciences & Cryptography*, 25(8-A), 2341–2352, 2022. DOI : 10.1080/09720529.2020.1838744
- [28] Shtayat, J., Al-Kateeb, A., An Encoding-Decoding algorithm based on Padovan numbers, arXiv:1907.02007, 2019.
- [29] Singh, B., Bhadouria, P., Sikhwal, O., Sisodiya, K., A Formula for Tetranacci-Like Sequence, *Gen. Math. Notes*, 20(2) (2014), 136-141.
- [30] Sivakumar, B., James, V., A Notes on Matrix Sequence of Pentanacci Numbers and Pentanacci Cubes, *Communications in Mathematics and Applications*, 13(2), 603–611, 2022. DOI: 10.26713/cma.v13i2.1725
- [31] Sloane, N.J.A., The on-line encyclopedia of integer sequences, <http://oeis.org/>
- [32] Soykan, Y., Simson Identity of Generalized m-step Fibonacci Numbers, *International Journal of Advances in Applied Mathematics and Mechanics*, 7(2), 45-56, 2019.
- [33] Soykan, Y., Properties of Generalized (r,s,t,u)-Numbers, *Earthline Journal of Mathematical Sciences*, 5(2), 297-327, 2021. <https://doi.org/10.34198/ejms.5221.297327>
- [34] Soykan, Y., A Study On the Recurrence Properties of Generalized Tetranacci Sequence, *International Journal of Mathematics Trends and Technology*, 67(8), 185-192, 2021. doi:10.14445/22315373/IJMTT-V67I8P522
- [35] Soykan, Y., A Study on Generalized Fibonacci Polynomials: Sum Formulas, *International Journal of Advances in Applied Mathematics and Mechanics*, 10(1), 39-118, 2022. (ISSN: 2347-2529)
- [36] Soykan, Y., On Generalized Fibonacci Polynomials: Horadam Polynomials, *Earthline Journal of Mathematical Sciences*, 11(1), 23-114, 2023. E-ISSN: 2581-8147. <https://doi.org/10.34198/ejms.11123.23114>
- [37] Soykan, Y. On the Recurrence Properties of Generalized Tribonacci Sequence, *Earthline Journal of Mathematical Sciences*, 6(2), 253-269, 2021. <https://doi.org/10.34198/ejms.6221.253269>
- [38] Soykan, Y., A Study on Generalized Jacobsthal-Padovan Numbers, *Earthline Journal of Mathematical Sciences*, 4(2), 227-251, 2020. <https://doi.org/10.34198/ejms.4220.227251>
- [39] Soykan, Y., Taşdemir, E., A Study On Hyperbolic Numbers With Generalized Jacobsthal Numbers Components, *International Journal of Nonlinear Analysis and Applications*, 13(2), 1965–1981, 2022. <http://dx.doi.org/10.22075/ijnaa.2021.22113.2328>
- [40] Soykan, Y., On Dual Hyperbolic Generalized Fibonacci Numbers, *Indian J Pure Appl Math*, 2021. <https://doi.org/10.1007/s13226-021-00128-2>
- [41] Soykan, Y., Taşdemir, E., Göcen, M., Binomial Transform of the Generalized Third-Order Jacobsthal Sequence, *Asian-European Journal of Mathematics*, 15(12), 2022. <https://doi.org/10.1142/S1793557122502242>.
- [42] Soykan, Y., Taşdemir, E., Okumuş, İ., A Study on Binomial Transform of the Generalized Padovan Sequence, *Journal of Science and Arts*, 22(1), 63-90, 2022. <https://doi.org/10.46939/J.Sci.Arts-22.1-a06>
- [43] Soykan, Y., Taşdemir, E., Okumuş, İ., Göcen, M., Gaussian Generalized Tribonacci Numbers, *Journal of Progressive Research in Mathematics(JPRM)*, 14 (2), 2373-2387, 2018.
- [44] Soykan, Y., Okumuş, İ., Taşdemir, E., On Generalized Tribonacci Sedenions, *Sarajevo Journal of Mathematics*, 16(1), 103-122, 2020. ISSN 2233-1964, DOI: 10.5644/SJM.16.01.08

- [45] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, *Communications in Mathematics and Applications*, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102
- [46] Soykan, Y. Explicit Euclidean Norm, Eigenvalues, Spectral Norm and Determinant of Circulant Matrix with the Generalized Tribonacci Numbers, *Earthline Journal of Mathematical Sciences*, 6(1), 131-151, 2021. <https://doi.org/10.34198/ejms.6121.131151>
- [47] Soykan Y, Tetranacci and Tetranacci-Lucas Quaternions, *Asian Research Journal of Mathematics*, 15(1): 1-24, 2019; Article no.ARJOM.50749.
- [48] Soykan, Y., Gaussian Generalized Tetranacci Numbers, *Journal of Advances in Mathematics and Computer Science*, 31(3): 1-21, Article no.JAMCS.48063, 2019.
- [49] Soykan, Y., Matrix Sequences of Tetranacci and Tetranacci-Lucas Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 57-69, 2019, (ISSN: 2347-2529).
- [50] Soykan, Y., On Binomial Transform of the Generalized Tetranacci Sequence, *International Journal of Advances in Applied Mathematics and Mechanics*, 9(2), 8-27, 2021.
- [51] Soykan, Y., Özmen, N., Göcen, M., On Generalized Pentanacci Quaternions, *Tbilisi Mathematical Journal*, 13(4), 169-181, 2020.
- [52] Soykan, Y., Binomial Transform of the Generalized Pentanacci Sequence, *Asian Research Journal of Current Science*, 3(1), 209-231, 2021.
- [53] Sunitha, K., Sheriba. M., Gaussian Tribonacci R-Graceful Labeling of Some Tree Related Graphs, *Ratio Mathematica*, 44, 188-196, 2022.
- [54] Ulutaş, Y.T., Toy, D., Some Equalities and Binomial Sums about the Generalized Fibonacci Number  $u_n$ , *Notes on Number Theory and Discrete Mathematics*, 28(2), 252–260, 2022. DOI: 10.7546/nntdm.2022.28.2.252-260
- [55] Yilmaz, N., Split Complex Bi-Periodic Fibonacci and Lucas Numbers, *Commun.Fac.Sci.Univ.Ank.Ser. A1 Math. Stat.* 71(1), 153–164, 2022. DOI:10.31801/cfsuasmas.704435
- [56] Yilmaz, N., The Generalized Bivariate Fibonacci and Lucas Matrix Polynomials, *Mathematica Montisnigri*, Vol LIII, 33-44, 2022. DOI: 10.20948/mathmontis-2022-53-5
- [57] Yilmaz, N., Taskara, N., Tribonacci and Tribonacci-Lucas Numbers via the Determinants of Special Matrices, *Applied Mathematical Sciences*, 8(39), 1947-1955, 2014.
- [58] Vasanthi, S., Sivakumar, B., Jacobsthal Matrices and their Properties. *Indian Journal of Science and Technology* 15(5): 207-215, 2022, <https://doi.org/10.17485/IJST/v15i5.1948>
- [59] Waddill, M.E., Another Generalized Fibonacci Sequence, *M. E., Fibonacci Quarterly*, 5(3) (1967), 209-227.
- [60] Waddill, M.E., The Tetranacci Sequence and Generalizations, *Fibonacci Quarterly*, (1992), 9-20.