

Original Research Article

Effects of Different Solvents and their Purity on the Extraction of Total Phenolic Content, Total Flavonoid Content and Antioxidant Activity from the Peels of Lotkon (*Baccaurea Motleyana Müll. Arg.*) and Longan (*Dimocarpus Longan Lour.*)

ABSTRACT

The objectives of this study was to investigate how three types of solvent (ethanol, methanol, and acetone) affected the yield of total phenolic content (TPC), total flavonoids content (TFC) and DPPH radical scavenging activity of extracts from lotkon and longan peels. The results showed that the 50% methanol extracts for longan peel displayed the highest levels of TPC, TFC, and DPPH free radical scavenging activities. The TPC varied for both peel extracts (longan and lotkon) from 343.39 ± 0.22^j to 425.55 ± 0.19^a , 121.63 ± 0.29^j to 321.54 ± 0.12^a mg GAE/100g dry weight; TFC from 48.24 ± 0.09^j to 86.24 ± 0.11^a , 6.47 ± 0.37^g to 15.37 ± 0.11^a mg QE/100g dry weight and DPPH radical scavenging activity from 30.79 ± 0.03^j to 60.01 ± 0.02^a %, 25.84 ± 0.01^j to 54.76 ± 0.03^a % respectively. These findings showed that total phenolic, total flavonoid and antioxidant activity of lotkon and longan peel extracts are significantly influenced by the extraction solvents and their purity. The findings of this study serve as a valuable guide for the food sector in determining the ideal conditions for extracting antioxidants and desirable phenolic compounds from lotkon and longan peels, as well as a solid foundation for more research in the future.

Keywords: antioxidant; extraction solvent; phenolic substances; lotkon and longan peels.

1. INTRODUCTION

Lotkon (*Baccaurea Motleyana Müll. Arg.*) is one of the most widely consumed fruits in Bangladesh and is a family member of the Phyllanthaceae. Lotkon is a significant evergreen, sluggish, dioecious fruit tree native to tropical and subtropical regions. Longan (*Dimocarpus Longan Lour.*), sometimes referred to as Kath Lichu regionally is a popular fruit that originates from China and Southeast Asia that that is a subtropical evergreen tree belongs to the Sapindaceae family. Due to the juicy and rich acidic flesh of longan, it is widely prized for its distinctive taste and flavor and is denoted as a snack eaten frequently [27]. According to Srumsiri and Silman [24], longan contains approximately 12.4-19.6% peel. While lotkon contains approximately 36.11% peel of the whole weight [21]. According to Rakariyatham et al. [22], the peels of longans were found to contain more than 50 phenolic chemicals. Phenolics, flavonoids, tannins, related hydrolysis products, and variations are among the various types of phenolic chemicals. As stated by Nurmayani et al. [19], the fruit and its peel both contain antioxidants and phenolics, two naturally occurring protectors, in addition to the lotkon's nutritional and dietary worth. The major bioactive compounds present in lotkon peel are polyphenols, flavonoids, and tannins [12].

These bio-wastes can be used to make these antioxidants for practical use because longan and lotkon peels are both abundant sources of bioactive substances, and phenolic compounds are well known for their health benefits related to antioxidant properties as well as their potential application in food processing industries as bio preservatives that enable the production of food without synthetic chemicals for consumers [15]. Numerous investigations showed that the kind of solvent used has a significant impact on whether bioactive chemicals may be successfully extracted from plant material. Various phenolic compounds, including simple phenolics, phenolic acids, coumarins, flavonoids, stilbenes, hydrolyzable and condensed tannins, lignans, and lignins, are present in plants [17]. Extraction of bioactive compounds from plant material is typically performed using solvent extraction. Since the solvent

system is simple to use and effective for the extraction of various chemicals, different solvents are employed to prepare extracts from plant material. The extraction yield of the compounds is affected by the physical and chemical properties of the sample, the type of solvent used, the length of time and temperature of the extraction process, and the ratio of the sample to the solvent [7]. Depending on the solvent's polarity and the solubility of phenolics in the solvent, polyphenols can be recovered from plant material [16]. It is common practice to extract antioxidants from plants using water and aqueous solutions of ethanol, methanol, and acetone [25]. Solvent diffuses into the solid plant material during extraction, solubilizing substances of a comparable polarity [18]. The majority of polyphenols are extracted from plant material using polar solvents. Solvents like acetone, ethanol, methanol, and ethyl acetate are optimal for extracting polyphenolic chemicals [7]. The type of solvent used is crucial to the efficient extraction of bioactive chemicals from organic material. Solvent diffuses into the solid plant material during extraction, solubilizing substances of a comparable polarity [1].

Methanol was used to extract tea polyphenols [30] and acetone was used to extract wheat total phenolics [31], both of which were more successful than water. The polarity of the solvents was important to the extraction process, according to Alothman et al. [4] since it boosted the total solubility of antioxidant compounds from three tropical fruits pulps (honey pineapple, banana and thai seedless guava). According to Muhamad et al. [16], the extraction parameters and solvent polarity had an impact on the antioxidant, total flavonoid, and total phenolic content yields from *Averrhoa bilimbi* extract. There is a lack of research comparing how various solvents affect the extraction and antioxidant activity of longan and lotkon peel extract. This study aims to assess the effects of solvent type and their purity on the yields of total phenolic, total flavonoids and antioxidant activity in extracts of longan and lotkon peels.

2. MATERIAL AND METHODS

2.1 Chemical reagent

Aluminum chloride, sodium hydroxide, ethanol, acetone, Follin's reagent were bought from Merck (Darmstadt, Germany). All additional chemicals and reagents utilized in the experiment were of analytical grade.

2.2 Sample preparation

From the Jashore area of Khulna, Bangladesh, fruit samples of the lotkon (*Baccaurea Motleyana Müll. Arg.*) and longan (*Dimocarpus Longan Lour.*) were collected. Fruits were selected to ensure ripe as well as uniformity in size and color. The peels were hand detached after the fruits had been carefully cleansed with distilled water. Lotkon and longan peels were dried in an oven (Hanyangs Scientific Equipment Co. Ltd., Korea) at 60 °C for 48 hrs and 60 °C for 24 hrs respectively. Then peels were ground (maximum particle size 0.4 mm) into powder in a lab mill (FW100, Taisite Instrument Co., Ltd, Tianjin, China). The supplies were stored in a desiccator at room temperature before to use.

2.3 Extraction Procedures

With a small adjustment, extracts from samples of lotkon and longan fruits were made using the technique outlined by Addai et al. [2].

Using the maceration process, antioxidant and bioactive chemicals were extracted from fruit samples. For this procedure, 10 ml of solvent (1:10 w/v) was applied to universal bottles containing 1 gm of peel powder. Pure acetone, ethanol, and methanol are among the solvents, along with their corresponding aqueous solutions at 50% and 70% concentrations. After that, an ultra-turrax homogenizer was used to thoroughly blend the samples (peel powder with solvents) for 5 min. A tabletop centrifuge (DSC-200A-2, Digisystem Laboratory Instrument Inc., Taiwan) was used to centrifuge all of the extracted samples for 10 min at 3000 rpm. The supernatants were collected and stored at -20°C for further analysis.

2.4 Analysis

2.4.1 Total Phenolic Content (TPC)

The total phenolic content was calculated using a modified Folin-Ciocalteu method [28]. Briefly, 5 ml of Folin-Ciocalteu reagent (1:10 v/v in distilled water) and 4 ml of 7.5% (w/v) sodium carbonate solution were

mixed with 1 ml of each extract (1 g/ml). The mixture was vortexed for 15 sec to enhance the color before being allowed to stand at 40°C for 30 min. In order to measure the absorbance at 765nm, a double beam UV-Vis spectrophotometer from Thermo Scientific was used (T60, UVV-is Spectrophotometer, USA). In place of the sample, water was used to prepare the blank. A collection of gallic acid reference solutions was used to compare a blank too. A calibration curve using gallic acid was used to determine the result as mg GAE/100g dry sample.

2.4.2 Total Flavonoid Content (TFC)

The total amount of flavonoid was measured by spectrophotometric method as described by Csepregi et al. [6]. After being combined with 0.3 mL of 5% NaNO₂, 1 mL of the sample was diluted with 3.5 mL of demineralized water. 0.4 mL of AlCl₃ (10% w/v) was added after 6 min. Following the addition of 2.1 mL of 1 M NaOH, 6 min later, 2.7 mL of demineralized water was added right away, bringing the total to 10 mL. After thorough mixing, the solution's absorbance was determined using a standardized ultraviolet-visible spectrophotometer at 510nm. By comparing the result to the quercetin calibration curves, which were created under the same circumstances, the result was represented in mg quercetin equivalent/100gdried sample.

2.4.3 Antioxidant Activity

The modified approach reported by [23] was used to measure the stable DPPH radical-scavenging activity. In order to conduct this test, 2 ml of extract solutions in a range of concentrations were added to 2 ml of 0.1 mM DPPH solutions. The mixture was then vigorously stirred for 15 sec. The solutions were then allowed to stand for 30 min at room temperature in a dark environment in order for a reaction to happen. Utilizing a twin-beam Scientific UV-Vis Spectrophotometer at 517nm (T60, UV-Vis Spectrophotometer, USA), absorbance was measured after 30 min against a blank. Equation (1) was used to direct the calculation,

$$\% \text{ DPPH free radical scavenging activity} = \frac{Ab(\text{blank}) - Ab(\text{sample})}{Ab(\text{blank})} \times 100 \dots\dots\dots(1)$$

2.4 Statistical analysis

Statistical Tool for Agricultural Research (STAR) software system was used to do an ANOVA on the triplicate data at a 5% level of significance.


3. RESULTS AND DISCUSSION

3.1 Total phenolic content (TPC)

Natural phenolics' major health benefits are mostly brought about by their antioxidant activity [9]. There is a correlation between the extraction yields of phenolic compounds and the kind of solvent used for the extraction, which might range from polar to non-polar [13]. Total phenolic content extraction yields from lotkon and longan peels were significantly ($p < 0.05$) affected by solvent type and their purity. The findings demonstrated that longan peel extract had significantly more total phenolic content than lotkon peel extract (Fig.1). Comparing the different solvent concentrations, 50% methanol had the highest amounts of TPC (425.55 ± 0.19^a and 321.54 ± 0.12^a mg GAE/100g dry weight) in longon and lotkon peel extracts respectively. While compared to other solvents, water resulted in lower yields of TPC (343.39 ± 0.22^j and 121.63 ± 0.29^j mg GAE/100g dry weight) in longon and lotkon peel extracts respectively. The TPC value was influenced by the extracting solvents in the following order, from high to low: 50% methanol > 50% ethanol > 50% acetone. These results agree with Park et al. (20), who discovered that methanol provided the best TPC recovery from orange peel. Similar findings were reported by Ali et al. [3], who found that methanol was the most effective solvent for extracting phenolic content from ginger fruit. According to Turkmen et al. [26], solvents with various polarities considerably affected phenolic compounds. Muhamad et al. [16] also found that *Averrhoa bilimbi* had the maximum total phenolic levels at a solvent concentration of 50% methanol. Our findings also agree with Addai et al. [2], who discovered that methanol was the most effective solvent for extracting phenolic content from papaya fruit. The highest extraction yield was observed with methanol, which may be a result of the reduced polyphenol oxidase

(PPO) these which is that is

activity in extracts an enzyme



responsible for the oxidation of phenolic substances [1].

Fig.1: Effect of different solvents and their purity on TPC of lotkon and longan peels extract

3.2 Total flavonoid content (TFC)

Flavonoids, which include flavones, flavanols, and condensed tannins, are common secondary metabolites of plants. Fig.2 shows the total flavonoid content extraction yields from lotkon and longan peels were significantly ($p < 0.05$) affected by solvent type and their purity. Methanol at 50% had the highest amounts of TFC (86.24 ± 0.11^a and 15.37 ± 0.11^a mg QE/100g dry weight) in longon and lotkon peel extracts respectively. While water resulted in lower yields of TFC (48.24 ± 0.09^j and 6.47 ± 0.37^g mg QE/100g dry weight) in longon and lotkon peel extracts respectively. The TFC yields by extraction solvents were as follows, from highest to lowest: 50% methanol > 50% ethanol > 50% acetone.

These findings agree with Ma et al. [14], who found that methanol was the most efficient solvent for extraction for the flavonoid compound from Penggan (*Citrus reticulata*) peel. Similar findings were reported by Gonzalez et al. [10], who found that extraction efficiency of TFC changes with polarity of solvents in lemon peels. Muhamad et al. [16] reported that *Averrhoa bilimbi* had the maximum total flavonoid content at a solvent concentration of 50% methanol. Our findings also agree with Addai et al. [2], who discovered that methanol proved to be the most efficient solvent for extracting phenolic content from papaya fruit. Abad-García et al. [1] reported that methanol produced highest for extraction for the flavonoid compound due to reduced the activity of polyphenol oxidase (PPO) enzymes which responsible for the oxidation of flavonoid compounds.




Fig. 2: Effect of different solvents and their purity on TFC of lotkon and longan peels extract

3.3 DPPH free radical scavenging activity

Antioxidants are recognized to have a crucial function in preventing oxidative cell damage [5]. Dietary antioxidants, which can scavenge free radicals, can reduce the risk of the disease. As a result, determining the radical scavenging efficacy of antioxidants is essential [8]. Fig. 3 shows that extraction solvents and their purity significantly affected the antioxidant activity of the lotkon and longan peel extracts. The results showed that longan peel extract had significantly ($p < 0.05$) higher scavenging activity than lotkon peel extract. The DPPH values of both extracted peels (lotkon and longan) decrease as the concentration of the organic solvent increases until it reaches 100%. Methanol at 50% had the highest value of DPPH (60.01 ± 0.02^a and 54.76 ± 0.03^a %) in longan and lotkon peel extracts respectively. Water extracts produced the lowest DPPH values for longan (30.79 ± 0.03^l %) and lotkon (25.84 ± 0.01^l %). The DPPH value was influenced by the extracting solvents in the following order, from high to low: 50% methanol > 50% ethanol > 50% acetone.


Our results agree with Muhamad et al. [16], who discovered that extracting antioxidant activity from *Averrhoa bilimbi* fruit needed a solvent concentration of 50% methanol. Similar findings were reported by Chavan et al. [7], who found that methanol had the highest level of antioxidant capacity for Saptarangi (*Salacia chinensis* L.) fruit pulp, whereas ethanol, acetone, and water had the lowest levels. According to Addai et al. [2] and Grant et al. [11], the lowest levels of DPPH radical scavenging activity were seen in pure solvents in the extraction process due to polarity of solvents plays an indirect role since it can increase the solubility of antioxidant chemicals. Alothman et al. [4] demonstrated that the chemical composition of the solvent has a substantial effect on the yield of DPPH value recovered from tropical fruits in Malaysia. According to Addai et al. [2], antioxidant activity results varied depending on the type of fruit, growth conditions, harvest maturity stage, storage settings, extraction time, and sample preparation method. Turkmen et al. [26] also reported that different extraction solvents probably had an impact in producing inconsistent results of antioxidant activity. The DPPH test is known as an electron transfer assay; the high value produced by methanol can be attributed to the quick transfer of electrons from the phenoxide anion to the radical as a result of partial ionization. [29].

Fig. 3: Effect of different solvents and their purity on DPPH% of lotkon and longan peels extract

4. CONCLUSION

In summary, our findings clearly demonstrated that different solvent extracted significantly varied phenolic and flavonoid levels as well as antioxidant activity of both peel (lotkon and longan) extracts. The yield of total phenolic, total flavonoid and antioxidant activity was influenced by the characteristics of the longan and lotkon

as well as the extracting solvents. All of available solvents were used to evaluate extracts of longan and lotkon peels, methanol was shown to be most effective extracting total phenolic, total flavonoid and antioxidant activity (DPPH).



COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Abad-García B, Berrueta LA, Lopez MDM, Crespo FI, Gallo B and Vicente F. Optimization and validation of a methodology based on solvent extraction and liquid chromatography for the simultaneous determination of several polyphenolic families in fruit juices. *Journal of Chromatography A*. 2007; 1154(1-2): 87-96. doi: <https://doi.org/10.1016/j.chroma.2007.03.023>
2. Addai ZR, Abdullah A and Abd S. Effect of extraction solvents on the phenolic content and antioxidant properties of two papaya cultivars. *Journal of Medicinal Plants Research*. 2013; 7: 3354-3359. <https://doi.org/10.5897/JMPR2013.5116>
3. Ali G, Hawa ZE, and Asmah R. Effects of solvent type on phenolics and flavonoids content and antioxidant activities in two varieties of young ginger (*Zingiber officinale Roscoe*) extracts. *J. Med. Plants Res*. 2011; 5(7):1147-1154. <https://doi.org/10.5897/JMPR.9000540>
4. Alothman M, Bhat R and Karim A.A. Antioxidant capacity and phenolic content of selected tropical fruits from Malaysia, extracted with different solvents. *Food chemistry*. 2009;115(3): 785-788. <https://doi.org/10.1016/j.foodchem.2008.12.005>
5. Alothman M, Bhat R and Karim AA. UV radiation-induced changes of antioxidant capacity of fresh-cut tropical fruits. *Innov. Food Sci. Emerg. Technol*. 2009;10:512-516 <https://dx.doi.org/10.1016/j.ifset.2009.03.004>
6. Csepregi K, Kocsis M and Hideg É. On the spectrophotometric determination of total phenolic and flavonoid contents. *Acta Biologica Hungarica*. 2013;64(4):500-509. <https://doi.org/10.1556/abiol.64.2013.4.10>
7. Chavan JJ, Jagtap UB, Gaikwad NB, Dixit GB and Bapat VA. Total phenolics, flavonoids and antioxidant activity of Saptarangi (*Salacia chinensis L.*) fruit pulp. *Journal of plant biochemistry and biotechnology*. 2013; 22(4): 409-413. <https://doi.org/10.1007/s13562-012-0169-3>
8. Dai j and Mumper rj. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. *Molecules*. 2010; 21;15(10):7313-7352. <https://doi.org/10.3390/molecules151073>
9. Fang YZ, Yang S, and Wu G. Free radicals, antioxidants, and nutrition. *Nutrition*. 2002 ;18(10):872-9. [https://doi.org/10.1016/s0899-9007\(02\)00916-4](https://doi.org/10.1016/s0899-9007(02)00916-4)
10. González ME, Domínguez PR, Moreno DA and García VC. Natural bioactive compounds of Citrus limon for food and health. *Journal of pharmaceutical and biomedical analysis*.2010; 51(2): 327-345. doi: <https://doi.org/10.1016/j.jpba.2009.07.027>
11. Grant GH and Robert GB. A preliminary evaluation of antioxidant compounds, reducing potential, and radical scavenging of pawpaw (*Asimina tribloba*) fruit pulp from different stages of ripeness. *LWT. Food Sci. Technol*. 2009;42:275-279. <https://doi.org/10.1016/J.LWT.2008.05.006>
12. Hossain MA and Hossain MS. Optimization of Antioxidant Extraction from Freeze-dried Pulp, Peel, and Seed of Burmese grape (*Baccaurea ramiflora Lour.*) by Response Surface Methodology. *Research Square*; 2021. <https://doi.org/10.21203/rs.3.rs-347432/v1>

13. Khoddami A, Wilkes MA and Roberts TH. Techniques for analysis of plant phenolic compounds. *Molecules*. 2013; 18(2): 2328-2375.
14. Ma Y, Ye X, Hao Y, Xu G, Xu G and Liu D. Ultrasound-assisted extraction of hesperidin from Penggan (*Citrus reticulata*) peel. *Ultrasonics Sonochemistry*. 2008;15(3), 227-232. doi: <https://doi.org/10.1016/j.ultsonch.2007.03.006>
15. Martillanes S, Rocha PJ, Cabrera BM, Martín VD and Delgado AJ. Application of Phenolic Compounds for Food Preservation: Food Additive and Active Packaging. In M. Soto-Hernandez, M. Palma-Tenango, & M. d. R. Garcia-Mateos (Eds.), *Phenolic Compounds - Biological Activity*. IntechOpen. 2017; <https://doi.org/10.5772/66885>
16. Muhamad N, Muhmed SA, Yusoff MM and Gim bun J. Influence of solvent polarity and conditions on extraction of antioxidant, flavonoids, and phenolic content from Averrhoa bilimbi. *Journal of Food Science and Engineering*. 2014;4: 255-260. <https://doi.org/10.17265/2159-5828/2014.05.006>
17. Naczki M and Shahidi F. Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. *Journal of pharmaceutical and biomedical analysis*. 2006; 28;41(5):1523-1542. <https://doi.org/10.1016/j.jpba.2006.04.002>
18. Ncube NS, Afolayan AJ and Okoh AI. Assessment techniques of antimicrobial properties of natural compounds of plant origin: current methods and future trends. *African journal of biotechnology*. 2008; 7(12). <https://doi.org/10.5897/AJB07.613>
19. Nurmayani S, Widodo WD and Matra DD. Characterization of rambai (*Baccaurea motleyana*) genes putatively involved in sugar metabolism . IOP Conference Series: Earth and Environmental Science.2021;<https://doi.org/10.1088/1755-1315/694/1/012067>
20. Park JH, Lee M and Park E. Antioxidant activity of orange flesh and peel extracted with various solvents. *Preventive Nutrition and Food Science*. 2014; 19(4): 291. doi: <https://doi.org/10.3746%2Fpnf.2014.19.4.291>
21. Rafi KN, Sen P, Uddin Z and Aziz MG. Dietary Fiber Profiling of Different Parts of Burmese Grape (*Baccaurea ramiflora*) and Their Application. *Food Science and Engineering*. 2020; 95-106. <https://doi.org/10.37256/fse.122020582>
22. Rakariyatham K, Zhou D, Rakariyatham N and Shahidi F. Sapindaceae (*Dimocarpus longan* and *Nephelium lappaceum*) seed and peel by-products: Potential sources for phenolic compounds and use as functional ingredients in food and health applications. *Journal of Functional Foods*. 2020; 67:103846. <https://doi.org/10.1016/j.jff.2020.103846>
23. Shimada K, Fujikawa K, Yahara K and Nakamura T. Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *Journal of agricultural and food chemistry*. 1992 ;40(6):945-948.
24. Sruamsiri S and Silman P. Chemical composition and in vitro digestibility of by-products from longan production. *Journal of Agricultural Research and Extension*. 2015; 32(2): 50-58.

25. Sun T and Ho CT. Antioxidant activities of buckwheat extracts. Food chemistry. 2005;90(4):743-749. <https://doi.org/10.1016/j.foodchem.2004.04.035>
26. Turkmen N, Sari F and Velioglu YS. Effect of extraction solvents on concentration and antioxidant activity of black and black mate polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods. Food Chem. 2006; 99:838-841 <https://doi.org/10.1016/j.foodchem.2005.08.034>
27. Uma DB, Ho CW and Wan AWM. Optimization of extraction parameters of total phenolic compounds for Henna (*Lawsonia inermis*) leaves. Sains Malaysiana.2010; 39:119-128
28. Wootton BPC, Moran A and Ryan L. Stability of the total antioxidant capacity and total polyphenol content of 23 commercially available vegetable juices before and after in vitro digestion measured by FRAP, DPPH, ABTS and Folin-Ciocalteu methods. Food research international. 2011;44(1):217-224.
29. Yang B, Jiang Y, Shi J, Chen F and Ashraf M. Extraction and pharmacological properties of bioactive compounds from longan (*Dimocarpus longan Lour.*) fruit A review. Food Research International. 2011; 44(7): 1837-1842.
30. Yao LH, Jiang YM, Caffin N, Darcy B, Datta N and Liu X, Singanusong R and Xu Y. Phenolic compounds in tea from Australian supermarkets. Food Chemistry. 2006;96(4):614-620. <https://doi.org/10.1016/j.foodchem.2005.03.009>.
31. Zhou K and Yu L. Effects of extraction solvent on wheat bran antioxidant activity estimation. LWT-Food science and Technology. 2004;37(7): 717-721.