

## **Original Research Article**

### **Effect of seasonal changes on photosynthetic pigments and proline contents of some plants growing naturally in Tayma region, Saudi Arabia**

#### **Abstract**

The present study was undertaken to understand adaptive behavior and the possibility of osmotic adjustment adopted by three plant species to tolerate harsh environmental conditions during the winter and summer seasons based on some soil and plant variables as well as the relationships among them. The plant species studied are *Hyoscyamus muticus* L. (*H. muticus*), *Anabasis setifera* Moq. (*A. setifera*) and *Teucrium polium* L. (*T. polium*) and were collected from Tayma Governorate, Tabuk region, Saudi Arabia. ANOVA results showed that the plant species, depths, and their interaction significantly affected most physicochemical properties of the soil supporting three plant species. Also, the plant species, seasons, and their interaction highly significantly affected photosynthetic pigments and proline contents. The highest values for pH and Ec in *A. setifera* soil, for Na<sup>+</sup>, Mg<sup>++</sup> and Cl<sup>-</sup> in *H. muticus* soil, and for other studied chemical properties in *T. polium* soil were recorded. Significantly increased water content % was observed in *A. setifera* soil at 20-40 depth across the winter season. Chlorophyll a (Chl.a), chlorophyll b (Chl.b) and Chl.a+b contents in *H. muticus* plants and Chl.a/b, total carotenoids, total pigment and proline contents in *T. polium* plants were significantly increased in comparison to the other plant species. Compared to the winter season, the amount of proline and photosynthetic pigments increased significantly in the summer season. The principal component (PC) and biplot analysis displayed that the three plant species were quite separated based on the variables studied, and showed positive correlations among most soil and plant variables, but these associations varied in their degree and consistency. The positive correlations were observed for PC1 with K<sup>+</sup>, Ca<sup>++</sup>, HCO<sub>3</sub><sup>-</sup> and SO<sub>4</sub><sup>-</sup> in soil as well as Chl.a/b, total carotenoids, total pigment and proline in *T. polium* plants. While PC2 showed highly positively correlated to Na and Mg in soil as well as Chl.a, Chl.b and Chl.a+b in *H. muticus* plants. As a result, the above soil chemical properties are associated closely with the above plant variables in the two previous plant species. The statistical study shows that due to its adaptive behavior and potential for osmotic adjustment, *T. polium* type generally adapts better to the dry desert environment and seasonal changes.

**Key words:** *H. muticus*, *A. setifera*, *T. polium*, soil properties, photosynthetic pigments, proline, PCA.

#### **1. Introduction**

The productivity of ecosystems is decreased, along with the diversity and relative abundance of species, by factors such as soil degradation, drought, global warming, extinction of perennially palatable species, overgrazing, and human-caused activities (Louhaichi et al. 2012; Ouled Belgacem et al. 2013). Numerous researchers have noted seasonal variations in the chemical or biological components of a variety of plant

species, for example Peters et al. (2018), Ramírez-Briones et al. (2019), Huang et al. (2020), El-Absy (2021). Seasonal variations affect the different biochemical characteristics of the wild plant species, these variations significantly were noticed in the active components of these species, and these variations are caused by variations in environmental factors including rainfall, temperature, and other factors (Reddy and Reddy, 1977; Kumar et al. 2007). Kamel and El-Absy (2020) and El-Absy (2021) conducted a study on the seasonal variation in the chemical composition of soil and plants and found that significant differences were seen to represent real environmental variations in chemical composition throughout the seasons, species, and their interactions.

All soils contain a variety of soluble salts, the most prevalent cations being calcium ( $\text{Ca}^{++}$ ), magnesium ( $\text{Mg}^{++}$ ), and sodium ( $\text{Na}^+$ ), which are associated with soil salinity (Alam, 1999). Misra and Tyler (1999) reported that although nutrient content may increase as the volume of soil water declines and results in enhanced uptake by plants for a brief period, low water availability in soils may have diverse impacts on it. Plants require rather high amounts of the so-called macroelements  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ , and  $\text{K}^+$  with values of  $>0.1\%$  of dry mass, in addition to the metabolic mineral elements nitrogen, phosphorus, and sulfur (Maathuis, 2009).

Desertification is a severe issue in semi-arid climates, where there is a rapid increase in soil erosion and a gradual decline in vegetation cover (El-Absy and Kamel, 2022). Two important environmental conditions that affect plant productivity and spread are drought and salinity factors (Bray et al., 2000). Plant species have developed a variety of techniques to cope with harsh environmental changes in all types of climates and terrains and to lessen negative impacts of ions from those areas of the plants where they may be damaging, in order to adapt and flourish to salt and drought conditions (Aslam et al. 2011; Wang et al., 2014). Plant species can regulate their osmotic pressure by accumulating osmolytes that are compatible with their cells' metabolic processes (Hasegawa et al. 2000). Aldamegh et al., (2013), Pradhan et al., (2020) and Salama et al. (2021) mentioned that dry weather conditions may activate the production of secondary phytochemical compounds or accumulation of numerous reactive oxygen species and osmolytes with high concentrations such as proline, which will play a critical role during stress acclimation in plants. Proline has been suggested to have a variety of roles in osmotically challenged plant tissues, including osmotic adjustment, protection of plasma membrane integrity, a source of carbon and nitrogen, as well as a sink of energy or decreasing power (Ahmad & Hellebust, 1988; Mansour et al. 1998; Verbruggen et al. 1996). The higher plant tissues that absorb light include important pigments called chlorophylls and carotenoids. Through the process of photosynthesis in the cell, they are able to convert the radiant energy of sunlight into the chemical energy of organic carbon compounds (Sims & Gamon, 2002). These compounds may have improved the plants' ability to withstand the effects of salinity and drought conditions (Al-Ghamdi, 2015). Due to the plant species capacity to maintain turgidity and water absorption, desert plants can withstand drought conditions (Sayed et al. 2013). Additionally, to achieve this the plant species undergo morphological, physiological, and molecular modifications (Levitt, 1980; Bartels & Sunkars, 2005).

The physical, chemical, and biological characteristics of the soil as well as plant growth are all impacted by the pH of the soil (Al-Mujahidy et al., 2013). Where

understanding the relationship between geographically heterogeneous soil characteristics and fertility is necessary due to public concern over boosting soil productivity and crop input efficiency (Abd-Elmabod et al. 2019). Because soil and plant species are interdependent, the amount of organic matter in the soil as well as the type and nature of vegetation affect its physical, chemical, and biological characteristics (Kim et al., 1995; Singwane & Malinga, 2012). Numerous research, including those by Al-Ghamdi (2015), Al-Mutairi (2017), Salama et al. (2017), and El-Ghani & Amer (2003) investigated the effects of soil factors on different plant species. El-Ghani & Amer (2003) stated that the correlations between the recorded soil characteristics and the vegetation are examined using multivariate statistical methods. The principal component analysis (PCA) has been used to study the relationship between soil parameters and plant variables by many researchers, for example El-Ghani & Amer 2003; Ferraz et al., 2019; Metwally et al., 2019; Abdel-Fattah et al., (2021) and El-Absy (2021 and 2022).

The major aim of this research was to calculate soil physicochemical properties and study seasonal variations of photosynthetic pigments and proline contents in three plant species growing under natural conditions in Tayma Governorate, Tabuk region, Saudi Arabia, as well as, elucidate the relationship between soil and plant variables by PCA and Biplot analysis. These plant species are *Hyoscyamus muticus* L. (*H. muticus*), *Anabasis setifera* Moq. (*A. setifera*) and *Teucrium polium* L. (*T. polium*).

## **2. Materials and methods**

### **2.1. Study area**

The current study was conducted between January 2022 (winter) and August 2022 in Tayma Governorate, Tabuk Region, Saudi Arabia (summer). Tayma is a city located on the western edge of the great sand dune desert, the Nafoud Al-Kebir. The oasis of Tayma is located in the Province of Tabuk, 255 km southeast of the city of Tabuk and extends between latitude: 27°37'47"N and longitude: 38°32'38"E (Fig. 1). This area has an arid climate with year-round temperatures that range from 0 °C in the winter to approximately 50 °C in the summer. Less than 150 mm of precipitation per year on average is relatively little (Al-Qahtani, 2019).



**Fig. 1. Location map of Tayma Governorate, Tabuk Region, Saudi Arabia.**  
([https://geanderson.files.wordpress.com/2014/11/maps\\_tayma.jpg](https://geanderson.files.wordpress.com/2014/11/maps_tayma.jpg)).

The weather information for 2022 regarding temperature, relative humidity, and rainfall is shown in Fig. 2. Average high and low temperatures, relative humidity, and rainfall for the historical period of 2022 were reported with values of 28.41 °C, 16.05 °C, 29.44%, and 3.16 mm, respectively. The months of June, July, and August have the highest high and low temperatures, while January had the lowest temperature readings. The months with the highest relative humidity % were January and December, while the ones with the lowest percentages were May and June. The highest rainfall was observed between January and October months, but the lowest rainfall occurs in the months of June and July.




Fig. 2. Monthly average high and low temperatures (°C), relative humidity (%) and rainfall (mm) in the region studied (<https://tcktcktck.org/saudi-arabia/tabuk/tayma>).

### 2.3. Soil physical and chemical properties

In the study area, soil samples were carefully taken from the soil associated with *H. muticus*, *A. setifera*, and *T. polium* at two random depths of 0-20 cm and 20-40 cm. Each sample was divided into three replicates, which were then transported to the lab in sealed tins to be utilized for soil physical and chemical parameters. As recommended by Jackson (1967) and Rowell (1994) for soil texture, soil samples were air-dried, sieved, and utilized for the physical analysis of soil particles. The results of the physical analysis are reported as a percentage of the original weight. The Rowell (1994) described approach was used to determine the soil moisture content. According to Jackson (1962), the soil-water paste was used to measure electrical conductivity (EC;  $\text{ds m}^{-1}$ ) and pH value for each sample. Using a saturation paste devised by Tuzuner (1990), the soil chemical properties including sodium, potassium, calcium, magnesium (Cation;  $\text{meq L}^{-1}$ ), chlorides, bicarbonate, and sulphate (Anion;  $\text{meq L}^{-1}$ ), were determined.

### 2.4. Plant analysis

The samples of *H. muticus* (Solanaceae), *A. setifera* (Amaranthaceae), and *T. polium* (Lamiaceae) were manually randomly collected in triplicates at the region under investigation in both the winter and summer seasons of 2022. Then they were put in plastic bags before being transported right away to the lab for preparing and processing. The contents of the photosynthetic pigments were measured spectrophotometrically, and using the wavelengths of 663, 645 and 470 nm, the chlorophyll a (Chl.a), chlorophyll b (Chl.b) and total carotenoids were determined by equations of Lichtenthaler (1987), respectively, as follows:

$$\text{Chl. a (mg g}^{-1} \text{ FW)} = [(12.7 \times A_{663}) - (2.69 \times A_{645})] \times V / (1000 \times W)$$

$$\text{Chl. b (mg g}^{-1} \text{ FW)} = [(22.9 \times A_{645}) - (4.68 \times A_{663})] \times V / (1000 \times W)$$

$$\text{Carotenoids (mg g}^{-1} \text{ FW)} = [(1000 A_{470}) - (2.27 \text{ Chl. a}) - (81.4 \text{ Chl. b}) / 226] \times V / (1000 \times W)$$

Where, V: is the volume of the extracting liquid, W: is the weight in grams of the fresh leaf sample, and  $A_{663}$ ,  $A_{645}$  and  $A_{670}$ : are the corresponding wavelengths of the light density value, respectively.

The proline content of samples in the three plant species was determined at 16 DAA according to the standard method of Bates et al., (1972). Following sample collection, preparation, and processing, a spectrophotometer (SPECTRO UV-VIS RS Spectrophotometer, Labo Med, Inc.) was used to determine the proline content on a fresh weight basis by utilizing the equation below:

$$\text{Proline } (\mu\text{moles/g of fresh plant materials}) = \{(\mu\text{g proline/mL} \times \text{mL toluene})/115.5 \mu\text{g}/\mu\text{moles}\}/(\text{g sample}/5)$$

## 2.5. Statistical analysis:

The Komolgorov-Smirnov test was used to confirm the normality of the data distribution in this study. Then, using the Steel & Torrie (1980) method, the measured data were subjected to two- and three-way ANOVA tests as well as the coefficient of variation (CV%) to identify any significant differences ( $p \leq 0.05$  and  $p \leq 0.01$ ) in the impact of the experimental factors and their interactions. The resulting data were presented as average  $\pm$  standard error (SE), and multiple comparisons were decided using the least significant difference test (L.S.D.) at the 0.05 level of probability, according to Steel & Torrie (1980). For a better understanding of the correlation between the soil and plant variables in both seasons, principal component analysis (PCA) was used. All statistical analysis was performed using the computer software program OriginPro 2018 b9.5.0.193.

## 3. Results and Discussion

### 3.1. Soil analysis

The physical properties % of the soil supporting *H. muticus*, *A. setifera*, and *T. polium* in the two depths at the studied area are shown in Tables (1). The two factors of plant species ( $P < 0.01$ ) and depths ( $P < 0.05$ ) significantly influenced the percentages of sand, silt, and clay in their soil. While the species x depths interaction showed insignificant ( $P > 0.05$ ) effects for these soil properties. These results were consistent with earlier research by Salama et al. (2014), Moustafa et al. (2021), and Al-Taisan (2022). The plant type of *T. polium* recorded a significantly higher sand % compared with the other two plant species investigated. However, the highest percentage of silt and clay was found in the soil associated with *A. setifera* type in the region studied. During the 0-20 and 20-40 depths, the sand % in the 20-40 depth as well as the silt and clay % in the 0-20 depth had recorded the highest values at plants soil. As for the interaction between the two factors, the highest values of sand% in *T. polium* soil at the 20-40 depth, and silt and clay % in *A. setifera* at the 0-20 depth were observed in this area studied. The soil of three plant species generally had higher sand percentages in the two soil depths than other soil particles, indicating a sandy texture in the soil at the study location. According to USDA (2002), due to rapid water infiltration and percolation, as well as strong soil aeration, sandy soil, sometimes referred to as light-textured soil, has a low ability to hold water. Moustafa et al. (2021) and El-Absy (2022) mentioned that the proportions of sand, silt, and clay in soils collected from various environments differed significantly.

Table 1. Physical proprieties % of the soil supporting three plant species in two depths at Tayma Governorate, Tabuk Region, Saudi Arabia.

Factors	Sand	Silt	Clay	Soil Texture
Species				
<i>H. muticus</i> (H)	82.85±0.92b	11.17±0.61a	5.98±0.32b	Sandy
<i>A. setifera</i> (A)	81.37±0.45b	11.67±0.21a	6.97±0.31a	Sandy
<i>T. polium</i> (T)	91.17±0.55a	5.90±0.45b	2.93±0.15c	Sandy
<i>P</i> -Values	0.00**	0.00**	0.00**	
Depths				
0 – 20	84.38±1.55b	10.06±0.88a	5.56±0.69a	Sandy
20-40	87.88±1.63a	8.10±1.06b	4.03±0.58b	Sandy
<i>P</i> -Values	0.07*	0.08*	0.08*	
Species x Depths Interaction				
H x (0 – 20)	82.26±1.39a	11.71±0.88a	6.03±0.50a	Sandy
H x (20 – 40)	83.44±1.40a	10.63±0.89a	5.93±0.51a	Sandy
A x (0 – 20)	80.63±0.53a	11.76±0.34a	7.62±0.19a	Sandy
A x (20 – 40)	82.10±0.45a	11.58±0.30a	6.32±0.19a	Sandy
T x (0 – 20)	90.25±0.51a	6.72±0.27a	3.03±0.27a	Sandy
T x (20 – 40)	92.09±0.65a	5.09±0.50a	2.83±0.17a	Sandy
<i>P</i> -Values	0.95 <sup>ns</sup>	0.49 <sup>ns</sup>	0.20 <sup>ns</sup>	
CV%	1.87	10.74	11.11	

Statistically significant differences at \* $p \leq 0.05$  and \*\* $p \leq 0.01$ ; ns: indicate the non-significant difference. Different lowercase letters in the same column indicate statistically significant differences at  $p \leq 0.05$  according to the LSD test

The chemical proprieties of the soil supporting three plant species in two depths at the region studied are shown in Table 2. The two-way ANOVA demonstrated a significant effect ( $P < 0.05$  or  $0.01$ ) of plant species, depths and their interaction on all soil chemical properties under study, with the exception of Ec,  $\text{HCO}_3^-$  and  $\text{SO}_4^{--}$  by depth factor, as well as pH, Ec and  $\text{Ca}^{++}$  by species x depths interaction. Our results were in line with El-Absy (2021), El-Absy and Kamel (2022), Han et al. (2022) and Jamin et al. (2022).

The highest values of  $\text{Na}^+$ ,  $\text{Mg}^{++}$  and  $\text{Cl}^-$  in *H. muticus* soil, pH and Ec in *A. setifera* soil and other soil chemical proprieties in *T. polium* were recorded at the region under study. While the values of pH, Ec and  $\text{Ca}^{++}$  were higher in the second depth compared with the first dept. The opposite is true for other soil chemical proprieties evaluated from the two depths. During the two depths, the values of  $\text{Na}^+$  and  $\text{Mg}^{2+}$  in *H. muticus* soil as well as  $\text{K}^+$ ,  $\text{Ca}^{++}$ ,  $\text{HCO}_3^-$  and  $\text{SO}_4^{--}$  in *T. polium* soil were higher than in the other soil chemical proprieties. On the other hand, the highest values of pH and Ec in the second depth and  $\text{Cl}^-$  in the first depth were observed in *A. setifera* soil. The study site tended to be alkaline because the pH values were higher than 7.7 in the two depths across the plant species soil. Marschner (1995) mentioned that cation elements including magnesium, calcium, and potassium as well as other elements become less soluble at alkaline pH values higher than 7.5 due to chemical interactions with particular

substances, such as  $\text{HPO}_4$  and  $\text{CaCO}_3$ . Because soluble salts are more prone to accumulate in arid soils due to low precipitation and temperature rise, decreases in EC values may have a significant influence on plant health and nutrient availability (Smith and Doran 1996; Corwin and Lesch 2005). The rise in total soluble salts in plant species soil could be the cause of alkalinity, as reported by Salama et al. (2021).

Local variation in the soil properties surrounding single plants was what caused the diversity in plant species (Shaltout et al. 2020), and significant variations in environmental variables reflect the variability in the soil texture and properties (Al-Mutairi, 2017). The location with respect to the distance from the sea and the types of surrounding mountains may be the cause of the variance in values of soil chemical properties between various environments, thus both of which have a substantial impact on soil characteristics (Moustafa et al. 2021). In general, soil properties like texture, pH, the presence of microaggregates, major cations, and organic matter can have an impact on the rhizospheric community either directly or indirectly (Garbeva et al. 2004). The chemical and physical properties of the soil have a significant impact on plant growth patterns (Al-Zahrani and Al-Amer, 2006) due to their impact on water supplies (Batanouny & Baeshin, 1983), thus the plants are able to adjust and flourish in environments with various soil characteristics (Comole et al. 2021).

Table 2. Chemical proprieties of the soil supporting three plant species in two depths at Tayma Governorate, Tabuk Region, Saudi Arabia.

Factors	pH	Ec (ds m <sup>-1</sup> )	Cation (milliequivalent/Liter)				Anion (milliequivalent/Liter)		
			Na <sup>+</sup>	K <sup>+</sup>	Ca <sup>++</sup>	Mg <sup>++</sup>	Cl <sup>-</sup>	HCO <sub>3</sub> <sup>-</sup>	SO <sub>4</sub> <sup>-</sup>
Species									
<i>H. muticus</i> (H)	8.00±0.06a	4.90±0.47c	11.76±0.33a	0.60±0.03b	5.06±0.29b	114.33±0.43a	9.45±0.38a	2.45±0.11c	2.55±0.12c
<i>A. setifera</i> (A)	8.01±0.01a	6.68±0.40a	5.92±0.56b	0.29±0.04c	3.70±0.46c	112.30±0.72b	9.42±0.75a	3.17±0.34b	3.24±0.03b
<i>T. polium</i> (T)	7.71±0.04b	5.08±0.29b	6.30±0.38b	1.30±0.06a	7.65±0.39a	109.35±0.32c	8.01±0.25b	7.07±0.27a	4.41±0.14a
<i>P-Values</i>	0.00 <sup>**</sup>	0.00 <sup>**</sup>	0.00 <sup>**</sup>	0.00 <sup>**</sup>	0.00 <sup>**</sup>	0.00 <sup>**</sup>	0.01	0.00	0.00
Depths									
0 – 20	7.90±0.07a	4.76±0.30b	7.47±0.95a	0.81±0.16a	4.94±0.63b	112.39±0.73a	9.38±0.55a	4.24±0.65a	3.35±0.28a
20 – 40	7.91±0.06a	6.34±0.33a	8.52±0.99b	0.65±0.14b	6.00±0.65a	111.60±0.90b	8.53±0.29b	4.22±0.83a	3.45±0.29a
<i>P-Values</i>	0.94 <sup>ns</sup>	0.00 <sup>**</sup>	0.09	0.00 <sup>**</sup>	0.05	0.00 <sup>**</sup>	0.04	0.92 <sup>ns</sup>	0.41 <sup>ns</sup>
Species x Depths Interaction									
H x (0-20)	8.02±0.10a	3.90±0.20a	11.24±0.38a	0.64±0.02c	4.55±0.27a	113.83±0.55b	9.70±0.58a	2.37±0.19c	2.49±0.18c
H x (20-40)	7.98±0.10a	5.90±0.29a	12.29±0.36a	0.55±0.05c	5.57±0.29a	114.83±0.61a	9.20±0.59a	2.53±0.12c	2.61±0.17c
A x (0-20)	8.00±0.01a	5.82±0.17a	5.40±0.80b	0.37±0.03d	3.15±0.63a	113.70±0.58b	10.90±0.52a	3.72±0.41c	3.19±0.03b
A x (20-40)	8.02±0.02a	7.53±0.18a	6.43±0.81b	0.21±0.01e	4.25±0.61a	110.90±0.59c	7.94±0.53c	2.63±0.32c	3.29±0.02b
T x (0-20)	7.69±0.07a	4.57±0.30a	5.78±0.46b	1.41±0.06a	7.12±0.49a	109.63±0.03d	7.55±0.33c	6.62±0.40b	4.36±0.22a
T x (20-40)	7.73±0.06a	5.60±0.29a	6.83±0.47b	1.18±0.04b	8.18±0.52a	109.07±0.67d	8.47±0.09b	7.51±0.08a	4.46±0.21a
<i>P-Values</i>	0.80 <sup>ns</sup>	0.17 <sup>ns</sup>	0.04	0.08	1.00 <sup>ns</sup>	0.00 <sup>**</sup>	0.00	0.00	0.03
<i>C.V.%</i>	1.23	7.45	13.69	7.79	15.91	0.40		7.42	7.62

Statistically significant differences at  $*p \leq 0.05$  and  $**p \leq 0.01$ ; ns: indicate the non-significant difference. Different lowercase letters in the same column indicate statistically significant differences at  $p \leq 0.05$  according to the LSD test.

The results in Fig. 3 revealed a highly significant effect ( $P < 0.01$ ) of the plant species, seasons and depths on the water content % in soil associated with the three plant species. The same results were obtained by El-Lamey (2020), El-Absy (2021), and El-Absy and Kamel (2022), who stated that the depths and seasons factors were significant effects on the soil water content (%) of plant species in various habitats studied. While the effects of the first and second interaction on the soil supporting three plant species in two depths across winter and summer seasons were insignificant. Significantly, the water content % increased with the soil of *A. setifera* type, followed by *T. polium* and *H. muticus* species. Additionally, the water content % was higher in the winter season and 20-40 depth compared with the other season and depth. Misra and Tyler (1999) reported that seasonal effects may lead to beneficial changes in the water content of the soil, for example, an increase in the availability of phosphorus required for plant nutrition. In comparison to other interactions among the experimentally tested factors, the soil of *A. setifera* in the second depth throughout the winter season displayed increased water content (%). However, it was discovered that *H. muticus* soil at the first depth throughout the summer season had the lowest water content (%). According to Moustafa and Zayed (1996), the moisture gradient is complex and related to a variety of environmental factors, including climatic drought, soil texture, the character of the soil surface, height and slope.




Fig. 3. Water content % of the soil supporting three plant species in two depths across winter and summer seasons at Tayma Governorate, Tabuk Region, Saudi Arabia. Statistically significant differences at  $**p \leq 0.01$ ; ns: indicate the non-significant difference. Different lowercase letters in the same column indicate statistically significant differences at  $p \leq 0.05$  according to the LSD test.

### 3.2. Plant analysis

Seasonal variations are the primary factor influencing the chemical composition of plants because they reflect seasonal variations in physiological needs and activities rather than availability in plant content (Estevez et al. 2010; Sgarbossa et al. 2019).

Therefore, rich and diverse metabolic networks that enable the plant to produce a variety of chemicals support various strategies for adaptation (Wang et al. 2014). The statistical analysis in Table 3 showed that plant species, seasons, and their interactions with the amounts of all photosynthetic pigments in the three plant species had highly significant effects ( $P < 0.01$ ). Our findings were consistent with those of Uvalle Saucedo et al. (2008), Devi et al. (2015), El-Absy (2021), Khater et al. (2022), Malik et al. (2022), Neamah and Jdayea (2022), in which the findings likewise revealed seasonal variations had significant effects on all photosynthetic pigments in different plant species. The contents of chlorophyll a (Chl.a), chlorophyll b (Chl.b) and Chl.a+b showed a significant increase in *H. muticus* type, followed by *T. polium* and *A. setifera* species. While the contents of Chl.a/b, total carotenoids, and total pigment were significantly greater in *T. polium* than that in the other two plant species investigated. Moreover, all photosynthetic pigment contents displayed a significant increase in the summer season compared to the winter season, except Chl.a/b ratio ( $>1$ ). In both of the studied seasons, the highest values were noticed for Chl.a, Chl.b and Chl.a+b contents in *H. muticus* plant and for total carotenoids and total pigment contents in *T. polium* plant. On the other hand, the Chl.a/b ratio had highest in *T. polium* and *A. setifera* across the winter season. The Chl.a/b ratios for the three plant species under investigation were higher than 1 because the Chl.a content was higher than the Chl.b content in those species across both seasons. Salama et al., (2017), Huang et al. (2021), El-Absy (2022), El-Absy and Kamel (2022) all reported similar findings to our study.

Morsy et al. (2008) stated that desert plants were able to adapt to changes in light conditions and stressors thanks to the higher amounts of chlorophyll and carotenoids they attained in dry environments. While a decrease in Chl.a might be seen as a protective adaptation strategy that stops increasing photon absorption, based on a study by Ait Said et al. (2013). Also, chlorophyll can demonstrate a plants endurance for high pH environments (Cimen et al. 2014). The amount of chlorophyll in each leaf area shows how different plant species have adapted to the regional environmental circumstances (Ivanov et al. 2020).

Table 3. Photosynthetic pigments contents (g/100g fr. wt.) of the three plant species across winter and summer seasons at Tayma Governorate, Tabuk Region, Saudi Arabia.

Factors	Chlorophyll a (Chl.a)	Chlorophyll b (Chl.b)	Chl.a+b	Chl.a/b	Total Carotenoids	Total Pigment
Plant Species						
<i>H. muticus</i> (H)	2.58±0.05a	2.26±0.06a	4.84±0.11a	1.14±0.01c	408.15±3.37b	412.99±3.40b
<i>A. setifera</i> (A)	1.51±0.04c	1.23±0.01c	2.74±0.05c	1.23±0.02b	239.22±4.47c	241.96±4.50c
<i>T. polium</i> (T)	1.80±0.15b	1.33±0.26b	3.13±0.41b	1.53±0.19a	565.35±28.25a	568.47±28.66a
<i>P</i> -Values	0.00	0.00	0.00	0.00	0.00	0.00
Seasons						
Winter	1.84±0.16b	1.38±0.20b	3.22±0.36b	1.46±0.12a	384.69±37.03b	387.91±36.99b
Summer	2.08±0.18a	1.83±0.17a	3.91±0.35a	1.14±0.01b	423.79±57.55a	427.70±57.74a
<i>P</i> -Values	0.00	0.00	0.00	0.00	0.00	0.00
Plant Species x Season Interaction						

H x Winter	2.48±0.06b	2.14±0.02b	4.62±0.08b	1.16±0.01cd	403.62±1.73d	408.24±1.65d
H x Summer	2.68±0.02a	2.37±0.04a	5.05±0.06a	1.13±0.01ce	412.68±5.77c	417.73±5.71c
A x Winter	1.59±0.03d	1.24±0.02d	2.83±0.04d	1.28±0.01b	248.09±4.62e	250.92±4.58e
A x Summer	1.43±0.01e	1.21±0.01d	2.64±0.01d	1.18±0.01ce	230.35±0.20f	232.99±0.19f
T x Winter	1.46±0.04e	0.75±0.03e	2.21±0.06e	1.95±0.04a	502.35±1.15b	504.56±1.10b
T x Summer	2.13±0.07c	1.91±0.01c	4.04±0.07c	1.12±0.03de	628.34±4.62a	632.38±4.56a
<i>P</i> -Values	0.00	0.00	0.00	0.00	0.00	0.00
C.V.%	3.60	2.86	3.03	2.70	1.70	1.66

Statistically significant differences at  $**p \leq 0.01$ . Different lowercase letters in the same column indicate statistically significant differences at  $p \leq 0.05$  according to the LSD test.

Proline is an essential osmolyte produced at a high cost of energy, and it is increased in response to stress conditions (Parida et al. 2016). As shown in Fig. 4, the results of the two-way ANOVA analysis revealed very significant effects ( $P < 0.01$ ) of plant species, season, and their interaction on the proline content of the three plant species investigated as part of the region study. These findings agreed with those of El-Absy and Kamel (2019), Khater et al. (2022). The type of *T. polium* had a proline concentration that was significantly higher than that of *H. muticus* and *A. setifera* species. Significantly increased proline content of plants was observed in the summer season compared to the winter season. According to the interaction between the plant species and seasons, *T. polium* in the summer season demonstrated significantly higher proline content compared with other plant species in both seasons. In keeping with these findings, increased proline levels have also been observed under stressful conditions may increase *T. polium* plants tolerance to oxidative stress and aid in scavenging free radicals by El-Absy and Kamel (2019) and Khater et al. (2022). Also, Halperin and Flores (1997) reported *H. muticus* accumulated proline at a high rate when osmotically stressed. During *A. setifera* type, the proline accumulation and degeneration responded sensitively to climatic changes, as shown by Treichel et al. (1984). In most plant species, the proline can be utilized as a signal of altered physiological conditions including salt and drought stresses (Dhaka and Meena 2018; Al-Qahtani et al. 2020; El-Absy, 2021; El-Absy and Kamel 2022). When it comes to osmotic adjustment and photoprotection, proline accumulation and chlorophyll content reduction in plant species appear to play a part, helping the plants resist salinity, hypoxia, and their combinations in their natural habitats (Tounekti et al. 2018).

Generally, it can be believed that weather conditions were the root of the notable variations in photosynthetic pigments and proline concentrations in the plant species under study because the variances across seasons were weather-related. Seasonal variation in defenses is frequent in plant species, and seasonal changes effects on plant metabolites have been well-documented (Huang et al. 2020).




Fig. 4. Proline contents (mg/g) of the three plant species across winter and summer seasons at Tayma Governorate, Tabuk Region, Saudi Arabia. Statistically significant differences at \*\* $p \leq 0.01$ . Different lowercase letters in the same column indicate statistically significant differences at  $p \leq 0.05$  according to the LSD test.

### 3.3. Principal Component and Biplot Analyses:

For a clear understanding of correlations between soil and plant variables for plant species in both seasons, the principal component (PC) and Biplot analyses were performed (Fig. 5 and 6). The PC1 and PC2 displayed eigenvalues higher than one ( $>1$ ) and they recorded 62.73% and 37.27% variability of the total variation of all evaluated variables, respectively. Therefore, the two first components express higher variability and support the choice of the variable with a positive loading factor because they better explain variability than a single variable. The PCA results obtained here are in line with Kooch et al., (2008), Ferraz et al., (2019), Metwally et al., (2019) and El-Absy (2021 and 2022).

Positive loadings were noticed for K, Ca, carotenoids, total pigment and proline in both PCs, for water content,  $\text{HCO}_3^-$ ,  $\text{SO}_4^-$  and Chl.a/b in PC1, as well as for  $\text{Na}^+$ ,  $\text{Mg}^{++}$ , Chl.a, Chl.b and Chl.a+b in PC2. Generally, the PC1 with  $\text{HCO}_3^-$ ,  $\text{SO}_4^-$ , and Chl.a/b contents as well as PC2 with Chl.a, Chl.b, and Chl.a+b contents showed the largest positive loadings (Fig. 5). These results harmony with findings by Gil et al., (2014) and El-Absy (2021 and 2022), also who indicated these soil and plant variables are related to water stress and to salt stress. It is clear that these components represent a response to soil and plant variables evaluated, which has both positive and negative effects on both seasons at the study location. In order to increase the soil and plant variable examined under drought stress circumstances, the PC1 is therefore thought to be crucial. Similar to how PC1 and PC2 described various soil variables in this study, the PCA revealed distinct differences in soil characteristics across the study region (Kooch et al., 2008). In other studies, the soil variables EC,  $\text{Ca}^{++}$ ,  $\text{K}^+$ ,  $\text{Na}^+$ ,  $\text{Mg}^{++}$  and Cl<sup>-</sup> (Khafagi et al., 2013) and the soil variables EC,  $\text{Na}^+$  and Cl<sup>-</sup> (Al-Mutairi 2017) were the most important factors controlling the community structure of plants in the region under study.

The PC-Biplot for the soil and plant variables was made using PC1 and PC2, as shown in Fig. 6. Angles between variable vectors that are less than or equal to  $90^\circ$  (sharp angles) and greater than  $90^\circ$  (obtuse angles) represent a positive and negative correlation, respectively. The

Biplot analysis showed strong positive correlations among soil variables (Ferraz et al., 2019; Neina 2019; Abdel-Fattah et al., 2021; El-Absy 2021 and 2022) as well as among plant variables (Nejat and Sadeghi 2016; Yinping et al., 2021). Hendry and Price (1993) and Kaspariy et al., (2020) also reported a positive correlation between total chlorophyll and carotenoid contents which are crucial for assessing a plant's capacity to absorb light in the shade and defending it against stresses through photo-oxidation. Positive correlations were observed between water content (%) in soil and Chl.a/b, among pH,  $Mg^{++}$ ,  $Cl^-$ , Chl.a, Chl.b, and Chl.a+b contents, as well as among  $K^+$ ,  $Ca^{++}$ ,  $HCO_3^-$ ,  $SO_4^-$ , Chl.a/b, carotenoids, total pigment, proline contents. Na of soil was highly positively correlated with Chl.a, Chl.b, and Chl.a+b contents, but it had positively correlated with carotenoids and total pigment contents. Our results are as well in agreement with Gil et al., (2014) and El-Absy (2021 and 2022). In this regard, Gil et al. (2011) found that major soil variables ( $Ca^{++}$  and  $Mg^{++}$ ) and proline content in plants had positive associations using PCA, indicating that proline serves a functional purpose in the stress tolerance systems of the plant species under investigation.

The PC1 had higher soil and plant variables than the PC2. Furthermore, the first two PCs primarily distributed and distinguished these variables with the plant species into four groups. The first quarter (the highest for both PCs) comprised the  $K^+$  and  $Ca^{++}$  soil variables as well as carotenoids, total pigment and proline contents, which showed related to *T. polium* type. As for the *H. muticus* type, which is found in the second quarter (the lowest PC1 and the highest PC2), has high amounts of the soil variables  $Na^+$  and  $Mg^{++}$  as well as the plant variables Chl.a, Chl.b, and Chl.a+b. On the other hand, *A. setifera* type was linked with the soil variables Ec, pH, and  $Cl^-$  in the third quarter (the lowest PC1 and PC2). The fourth quarter (the highest PC1 and lowest PC2) included the water content %,  $HCO_3^-$ , and  $SO_4^-$  soil variables, which related to the Chl.a/b ratio. El-Absy (2022) after using PCA and Biplot analysis, reported a strong positive association between the soil and plant variables suggesting that the chemical properties of the soil have an impact on the plant variables and that elevating the soil variables will elevate the plant variables. According to Salama et al., (2013), there were clear negative associations between the soil variables including EC,  $Na^+$ ,  $K^+$ ,  $Ca^{++}$ ,  $Mg^{++}$ , and  $Cl^-$  and the richness of plant species. Generally, PC1 was positively correlated to soil variables ( $K^+$ ,  $Ca^{++}$ , water content %,  $HCO_3^-$ , and  $SO_4^-$ ) and plant variables (Chl.a/b ratio, carotenoids, total pigment and proline contents), which related to the *T. polium* plants. These findings suggest that *T. polium* type may survive and sustain a variety of environmental conditions throughout the growing year thanks to its adaptive behavior and potential for osmotic adjustment.




Fig. 5. The loading of the first two components with soil and plant variables of the three plant species. WC: water content %.




Fig. 6. Biplot diagram between PC1 and PC2 shows the correlation between soil (brown color) and plant (green color) variables with the three plant species (point red color) in both seasons. WC: Water content %.

## 5. Conclusions

In this study, significant effects ( $P < 0.05$  or  $0.01$ ) by the plant species, depths and their interaction on most soil physicochemical properties, and by the plant species, seasons and their interaction on photosynthetic pigments and proline contents were observed. Most soil and plant variables examined in both seasons had a high positive association, according to PCA and Biplot analysis. These findings show how the three plant species varied in their responses to seasonal changes in the area under investigation. In comparison to *H. muticus* and *A. setifera* species, *T.*

*polium* type generally adapts better to the arid desert environment and seasonal fluctuations because of its adaptive behavior and potential for osmotic adjustment.

## References

- Abdel-Fattah, M.K., Mohamed, E.S.;Wagdi, E.M., Shahin, S.A., Aldosari, A.A., Lasaponara, R. and Alnaimy, M.A. (2021). Quantitative Evaluation of Soil Quality Using Principal Component Analysis: The Case Study of El-Fayoum Depression Egypt. *Sustainability*, 13:1824. <https://doi.org/10.3390/su13041824>
- Abd-Elmabod SK, Bakr N, Muñoz-Rojas M, Pereira P, Zhang Z, Cerdà A, Jordán A, Mansour H, De La Rosa D, Jones L. Assessment of soil suitability for improvement of soil factors and agricultural management. *Sustainability*. 2019;11(6):1588. <https://doi.org/10.3390/su11061588>
- Ahmad I, Hellebust JA. 1988. The relationship between inorganic nitrogen metabolism and proline accumulation in osmoregulatory responses of two euryhaline microalgae. *Plant Physiology* 88, 348-354.
- Ait Said S, Torre F, Derridi A, Gauquelin T, Mevy JP. Gender, Mediterranean drought, and seasonality: photosystem II photochemistry in *Pistacia lentiscus*. *Photosynthetica*, 2013;51:552-564. DOI: 10.1007/s11099-013-0055-9
- Alam SM. Nutrient uptake by plants under stress conditions. In Pessaraki M (ed) *Handbook of plant and crop stress*. 1999;285-313. Marcel Dekker, New York.
- Aldamegh, M.A., Abdallah, E.M. and Hsouna, A.B. (2013). Evaluation of antimicrobial and antioxidant properties of leaves of *Emex spinosa* and fruits of *Citrullus colocynthis* from Saudi Arabia. *African Journal of Biotechnology*, 2(34):5308-5313. DOI: 10.5897/AJB2013.12987
- Al-Ghamdi, A.A.M. (2015). Ecological studies on the colocynth, *Citrullus colocynthis* (L.) (Cucurbitaceae) from Shada, Saudi Arabia and its insect repellent properties. *Life Science Journal*, 12(1):125-133.
- Al-Mujahidy, S.M.J., Hassan, M.M., Rahman, M.M. and Mamun-or-Rashid, A. (2013): Study on measurement and statistical analysis of adherent soil chemical compositions of leguminous plants and their impact on nitrogen fixation. *International Journal of Biosciences*, 3:112-119. <http://dx.doi.org/10.12692/ijb/3.6.112-119>
- Al-Mutairi, K.A. (2017). Influence of soil physical and chemical variables on species composition and richness of plants in the arid region of Tabuk, Saudi Arabia. *Ekológia (Bratislava)*, 36(2):112-120. <https://doi.org/10.1515/eko-2017-0010>
- Al-Qahtani S.M. (2019). Effect of soil properties on the diversity and distribution of weeds in citrus farms in arid region. *Applied Ecology AND Environmental Research* 17(1):723-732. [http://dx.doi.org/10.15666/aeer/1701\\_723732](http://dx.doi.org/10.15666/aeer/1701_723732)
- Al-Qahtani H, Alfarhan AH, Al-Othman ZM. Changes in chemical composition of *Zilla spinosa* Forssk. medicinal plants grown in Saudi Arabia in response to spatial and seasonal variations. *Saudi Journal of Biological Sciences*. 2020;27(10):2756-2769, <https://doi.org/10.1016/j.sjbs.2020.06.035>
- Aslam R, Bostan N, Nabgha-e-Amen M., Safdar, M, Safdar W. A critical review on halophytes: salt tolerant plants. *J Med Plants Res*. 2011;5:7108-7118. DOI: 10.5897/JMPRx11.009

- Al-Taisan W. A. (2022). Floristic diversity and vegetation of the az Zakhnuniyah Island, Arabian Gulf, Saudi Arabia. *Heliyon*, 8(7), e09996. <https://doi.org/10.1016/j.heliyon.2022.e09996>
- Al-Zahrani, H.S. and Al-Amer, K.H., (2006). A comparative study on *Citrullus Colocynthis* plant grown in different altitudinal locations in Saudi Arabia. *American-Eurasian Journal of Scientific research*, 1(1):1-7.
- Bartels, D. and Sunkars, R. (2005). Drought and salt tolerance in plants. *Cr Rev Plant Science* 24:23-58. <https://doi.org/10.1080/07352680590910410>
- Batanouny, K.H. and Baeshin, N.A. (1983). Plant communities along the Medina-Badr road across the Hejaz mountains, Saudi Arabia. *Vegetatio*, 53:3343. <https://doi.org/10.1007/BF00039769>
- Bates, L.S., Waldren, R.P. and Teare I.D. (1972). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39:205-207.
- Bray, E.A., Bailey-Serres, J. and Weretilnyk, E. (2000) Responses to abiotic stress. *Biochemistry & molecular biology of plants*. In: Grisse, W. and Jones, R., Eds., American Society of Plant Physiologists, Rockville, 1158-1203.
- Cimen, B., Yesiloglu, T., Incesu M, Yilmaz B. Growth and photosynthetic response of young 'Navelina' trees budded on to eight citrus rootstocks in response to iron deficiency. *N. Z. J. Crop Hortic. Sci.* 2014;42:170-182.
- Comole, A.A., Malan, P.W. and Tiawoun, M.A.P. (2021): Effects of *Prosopis velutina* invasion on soil characteristics along the riverine system of the molopo river in north-west province, south africa". *International Journal of Ecology*, vol. 2021, Article ID 6681577, 11. <https://doi.org/10.1155/2021/6681577>
- Corwin DL, Lesch SM. Apparent soil electrical conductivity measurements in agriculture. *Computers and Electronics in Agriculture*. 2005;46(1-3),11-43. <https://doi.org/10.1016/j.compag.2004.10.005>
- Devi K, Kapila S, Rao A. Seasonal variations in photosynthetic pigments of three species of Marchantiaceae. *International Journal of Advances in Pharmacy, Biology and Chemistry*. 2015;4(3): 713-718.
- Dhaka V, Meena KL. Seasonal variation in free proline content in some species of family euphorbiaceae of the rajasthan, India. *Journal of Experimental Biology and Agricultural Sciences*. 2018;6(1):249-252. DOI:10.18006/2018.6(1).249.252
- El-Absy KM and A.M Kamel (2019). Physiological and anatomical responses of *Teucrium polium* L. growing under different habitat conditions at North West Coast and South Sinai. *Journal of Biodiversity and Environmental Sciences (JBES)*, 15(6):40-52.
- El-Absy, K. M. (2021). Seasonal Changes of Some Metabolites in *Hyoscyamus boveanus* (Dunal) Asch. & Schweinf – Saint Katherine, South Sinai, Egypt. *Asian Journal of Biology*, 13(2), 49-66. <https://doi.org/10.9734/ajob/2021/v13i230184>
- El-Absy, K. M. (2022). Effect of Different Habitats Conditions on *Citrullus colocynthis* (L.) Schrad. Growing Naturally in Egypt and Kingdom of Saudi Arabia. *Journal of Advances in Biology & Biotechnology*, 25(2), 8-29. <https://doi.org/10.9734/jabb/2022/v25i230265>
- El-Absy, K. M., & Kamel, A. M. (2022). Ecophysiological Studies on *Salvadora persica*. *Asian Journal of Research in Botany*, 8(2), 35-52. Retrieved from <https://journalajrib.com/index.php/AJRIB/article/view/157>

- El-Lamey TM. Changes in some chemical compounds of *Retama raetam* (Forssk.) Webb & Berthel. in response to different environmental conditions. *Journal of Biodiversity and Environmental Sciences*. 2020;16(2):78-91.
- Estevez, J.A., Landete-Castillejos, T., GarcíaB, A.J., Ceacero, F., Martínez, A., Gaspar-López, E., Calatayud, A. and Gallego, L. (2010). Seasonal variations in plant mineral content and free-choice minerals consumed by deer. *Animal Production Science*, 50(3):177-185. <https://doi.org/10.1071/AN09012>
- Ferraz, G.A.S., PFerra, P F.P., Martins, F.B., Silva, F.M., Damasceno, F.A., Barbari, M. (2019). Principal components in the study of soil and plant properties in precision coffee farming. *Agronomy Research* 17(2):418-429. <https://doi.org/10.15159/AR.19.114>
- Garbeva, P., van Veen, J. A., & van Elsas, J. D. Microbial diversity in soil: selection microbial populations by plant and soil type and implications for disease suppressiveness. *Annual review of phytopathology*, 2004;42:243–270. <https://doi.org/10.1146/annurev.phyto.42.012604.135455>
- Gil, R., Bautista, I., Boscaiu, M., Lidon, A., Wankhade, S., Sa´nchez, H., Llinares, J., Vicente, O. (2014). Responses of five Mediterranean halophytes to seasonal changes in environmental conditions. *AoB Plants* 6: plu049; <https://doi.org/10.1093/aobpla/plu049>
- Halperin, S.J., Flores, H.E. (1997). Hyoscyamine and proline accumulation in water-stressed *Hyoscyamus muticus* 'hairy root' cultures. *In Vitro Cellular and Developmental Biology. Plant* 33(3): 240-244. DOI: 10.1007/s11627-997-0030-x
- Han W, Chen L, Su X, Liu D, Jin T, Shi S, Li T and Liu G (2022) Effects of Soil Physico-Chemical Properties on Plant Species Diversity Along an Elevation Gradient Over Alpine Grassland on the Qinghai-Tibetan Plateau, China. *Frontiers Plant Science*, 13:822268. <https://doi.org/10.3389/fpls.2022.822268>
- Hasegawa, P. M., Bressan, R. A., Zhu, J. K., & Bohnert, H. J. (2000). PLANT CELLULAR AND MOLECULAR RESPONSES TO HIGH SALINITY. *Annual review of plant physiology and plant molecular biology*, 51, 463–499. <https://doi.org/10.1146/annurev.arplant.51.1.463>
- Hendry, G.A.F. and Price, A.H. (1993). Stress Indicators: Chlorophylls and Carotenoids. In: Hendry, G.A.F. and Grime, J.P., Eds., *Methods in Comparative Plant Ecology*, Chapman Hall, London 148-152. <http://dx.doi.org/10.1007/978-94-011-1494-3>
- Huang, W., Bont, Z., Hervé, M.R. C.A. M. Robert & M. Erb. (2020). Impact of Seasonal and Temperature-Dependent Variation in Root Defense Metabolites on Herbivore Preference in *Taraxacum officinale*. *J Chem Ecol* 46, 63–75 (2020). <https://doi.org/10.1007/s10886-019-01126-9>
- Huang Z, Liu Q, An B, Wu X, Sun L, Wu P, Liu B, Ma X. Effects of planting density on morphological and photosynthetic characteristics of leaves in different positions on *Cunninghamia lanceolata* saplings. *Forests*. 2021;12: 853. <https://doi.org/10.3390/f12070853>
- Ivanov LA, Ronzhina DA, Yudina PK, Zolotareva NV, I. V. Kalashnikova IV, Ivanova LA. Seasonal dynamics of the chlorophyll and carotenoid content in the leaves of steppe and forest plants on species and community level. *Russ Journal Plant Physiology*. 2020;67:453-462. <https://doi.org/10.1134/S1021443720030115>
- Jackson, M.L. (1962). *Soil chemical analysis constable and co. Ltd. London.*

- Jackson, M.L. (1967). Soil chemical analysis. Pritice Hall of India Private., New Delhi., India.
- Jamin, J.; Diehl, D.; Meyer, M.; David, J.; Schaumann, G.E.; Buchmann, C. Physico-Chemical Soil Properties Affected by Invasive Plants in Southwest Germany (Rhineland-Palatinate)-A Case Study. *Soil System*, 2022, 6, 93. <https://doi.org/10.3390/soilsystems6040093>
- Kamel AM, El-Absy KM. Seasonal variations in protein patterns and mineral contents of *Lycium showii* under different habitat conditions. *Asian Plant Research Journal*. 2020;6(4): 91-103. <https://doi.org/10.9734/apri/2020/v6i430141>
- Kaspary, T.E., Cutti, L., Bellé, C., Casarotto, G. and Ramos R.F. (2020). Nondestructive analysis of photosynthetic pigments in forage radish and vetch. *Crop Production Rev. Ceres*, 67(6):424-431. <https://doi.org/10.1590/0034-737X202067060001>
- Khafagi, O.A., Sharaf, A.A., Hatab E.E. and Moursy, M.M. (2013). Vegetation Composition and Ecological Gradients in Saint Katherine Mountain, South Sinai, Egypt. *American-Eurasian J. Agric. & Environment Science*, 13(3):402-414. DOI: 10.5829/idosi.aejaes.2013.13.03.11313.
- Khater AK, T.M. El-Lamey, H.A. El-khamissi and M.M. Eldanasoury (2022). Seasonal variation in Photosynthetic Pigments, Phytohormones, and phenols of *Teucrium Polium* L. Growing in Wadi Halazien, Egypt. *Al-Azhar Journal of Agricultural Research*, 47(1):159-171. DOI: 10.21608/ajar.2022.266495
- Kim, C., Sharik, T. and Jurgensen, M. (1995). Canopy cover effects on soil nitrogen mineralization in northern red oak (*Quercus rubra*) stands in northern Lower Michigan. *Forest Ecology and Management*, 76(1-3):21-28. [http://doi.org/10.1016/0378-1127\(95\)03563-P](http://doi.org/10.1016/0378-1127(95)03563-P)
- Kooch, Y., Jalilvand, H., Bahmanyar, M.A. and Pormajidian, M.R., (2008). The Use of Principal Component Analysis in Studying Physical, Chemical and Biological Soil Properties in Southern Caspian Forests (North of Iran). *Pakistan Journal of Biological Sciences*, 11:366-372. DOI: 10.3923/pjbs.2008.366.372
- Kumar A, Kaul MK, Bhan MK, Punit K. Khanna PK, Suri KA. Morphological and chemical variation in 25 collections of the Indian medicinal plant, *Withania somnifera* (L.) Dunal (Solanaceae). *Genetic Resources and Crop Evolution*. 2007;54:655-660. <https://doi.org/10.1007/s10722-006-9129-x>
- Levitt J (1980) Responses of plants to environmental stress: chilling, freezing and high temperature stresses, 2<sup>nd</sup> Ed New York: Academic Press.
- Lichtenthaler HK. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. In: Packer, L.; Douce, R., Editors. *Methods in enzymology*. London: Academic Press. 1987;148: 350-82. [https://doi.org/10.1016/0076-6879\(87\)48036-1](https://doi.org/10.1016/0076-6879(87)48036-1)
- Louhaichi M., F. Ghassali, A.K. Salkini, S.L. Petersen (2012). Effect of sheep grazing on rangeland plant communities: Case study of landscape depressions within Syrian arid steppes, *Journal of Arid Environments*, 79: 101-106, <https://doi.org/10.1016/j.jaridenv.2011.11.024>
- Maathuis FJM. Physiological functions of mineral macronutrients. *Current Opinion in Plant Biology*, 2009;12:250-258. <https://doi.org/10.1016/j.pbi.2009.04.003>
- Malik, J.A.; AlQarawi, A.A.; AlZain, M.N.; Dar, B.A.; Habib, M.M.; Ibrahim, S.N.S. Effect of Salinity and Temperature on the Seed Germination and Seedling Growth of

- Desert Forage Grass *Lasiurus scindicus* Henr. Sustainability 2022, 14, 8387. <https://doi.org/10.3390/su14148387>
- Mansour MMF. 1998. Protection of plasma membrane of onion epidermal cells by glycinebetaine and proline against NaCl stress. Plant Physio. Biochem 36, 767-772.
- Marschner H. Mineral nutrition of higher plants. second edition. London: Academic Press. 1995;889. <https://doi.org/10.1016/C2009-0-63043-9>
- Metwally, M. S., Shaddad, S. M., Liu, M., Yao, R. J., Abdo, A. I., Li, P., Jiao, J. and Chen, X. (2019). Soil properties spatial variability and delineation of site-specific management zones based on soil fertility using fuzzy clustering in a hilly field in Jianyang, Sichuan, China. Sustainability, 11(24):7084. <https://doi.org/10.3390/su11247084>
- Misra A, Tyler G. Influence of soil moisture on soil solution chemistry and concentrations of minerals in the calcicoles *Phleum phleoides* and *Veronica spicata* grown on a limestone soil. Annals of Botany, 1999;84(3):401-410. <https://doi.org/10.1006/anbo.1999.0941>
- Morsy AA, Youssef AM, Mosallam HAM, Hashem AM. Assessment of Selected Species along Al-Alamein-Alexandria International Desert Road, Egypt. Journal of Applied Sciences Research. 2008;4(10):1276-1284.
- Moustafa AM, Zayed A. Effect of environmental factors on the flora of alluvial fans in southern Sinai, Journal of Arid Environments. 1996;32(4):431-443. <https://doi.org/10.1006/jare.1996.0036>
- Moustafa M, Alamri S, Al-Emam A, Alghamdi H, Shati A, Alrumman S, Sulayli A, Al-Khatani M, Abbas A. Biological, Physical and Chemical Properties of Nanosilver Particles Collected from Soil in Asir, Saudi Arabia. Arabian Journal for Science and Engineering. 2021;46:129-140. <https://doi.org/10.1007/s13369-020-04833-8>
- Neamah, S.I. and N.A. Jdayea, "Positive Response of *Hyoscyamus pusillus* Callus Cultures to Exogenous Melatonin on Biochemical Traits and Secondary Metabolites under Drought Conditions", International Journal of Agronomy, vol. 2022, Article ID 7447024, 10 pages, 2022. <https://doi.org/10.1155/2022/7447024>
- Neina, D. (2019). The Role of Soil pH in Plant Nutrition and Soil Remediation", Applied and Environmental Soil Science, vol. 2019, Article ID 5794869, 9 pages,9. <https://doi.org/10.1155/2019/5794869>
- Nejat, N. and Sadeghi, H. (2016). Finding out relationships among some morpho-biochemical parameters of christ's thorn (*Ziziphus spina-christi*) under drought and salinity stresses. Planta Daninha, Viçosa-MG, 34(4):667-674. doi: 10.1590/S0100-83582016340400006
- Ouled Belgacem A, Tarhouni M, and Louhaichi M. Effect of protection on plant community dynamics in the Mediterranean arid zone of southern Tunisia: a case study from Bouhedma national park. Land Degrad Dev. 2013;24(1):57-62. <https://doi.org/10.1002/ldr.1103>
- Parida AK, Veerabathini SK, Kumari A, Agarwal PK. Physiological, Anatomical and Metabolic Implications of Salt Tolerance in the Halophyte *Salvadora persica* under Hydroponic Culture Condition. Front. Plant Sci. 2016;7:351. doi:10.3389/fpls.2016.00351

- Peters, K., Gorzolka, K., Bruelheide, H., & Neumann, S. (2018). Seasonal variation of secondary metabolites in nine different bryophytes. *Ecology and evolution*, 8(17), 9105–9117. <https://doi.org/10.1002/ece3.4361>
- Pradhan, A. K., Rehman, M., Saikia, D., Jyoti, S.Y., Poudel, J. and Tanti, B. (2020). Biochemical and molecular mechanism of abiotic stress tolerance in plants. *Plant ecophysiology and adaptation under climate change: Mechanisms and Perspectives I*. Springer; 825-853. <https://doi.org/10.1007/978-981-15-2156-0>
- Ramírez-Briones E, Rodríguez-Macías R, Salcedo-Pérez E, Ramírez-Chávez E, Molina-Torres J, Tiessen A, Ordaz-Ortiz J, Martínez-Gallardo N, Délano-Frier JP, Zañudo-Hernández J. Seasonal Changes in the Metabolic Profiles and Biological Activity in Leaves of *Diospyros digyna* and *D. rekoii* “Zapote” Trees. *Plants*. 2019; 8(11):449. <https://doi.org/10.3390/plants8110449>
- Reddy RK. Reddy SJ. Elemental concentrations in medicinally important leafy materials, *Chemosphere*. 1997;34(9-10): 2193-2212. [https://doi.org/10.1016/S0045-6535\(97\)00078-7](https://doi.org/10.1016/S0045-6535(97)00078-7)
- Rowell, D.L. (1994). *Soil science methods and applications*. Longman Publishers, Singapor. 229.
- Salama, F.M., Abd El-Ghani, M.M., El-Naggar, S.M. and Aljarroushi, M.M. (2013). Vegetation analysis and species diversity in the desert ecosystem of coastal wadis of South Sinai, Egypt. *Journal of Biology and Earth Sciences*, (3)2:B214-B227.
- Salama F, El-Ghani MA, Gadallah M, El-Naggar S, Amro A. Variations in Vegetation Structure, Species Dominance and Plant Communities in South of the Eastern Desert-Egypt. *Notulae Scientia Biologicae*, 2014;6(1):41-58. <https://doi.org/10.15835/nsb619191>
- Salama, F.M., El-Ghani, M.M., El-Tayeh, N.A., Amro, A.A., and El-Naggar, S. (2017). Some aspects of drought resistance in *Citrullus colocynthis* L. in the Egyptian deserts. *Taekholmia*, 37(1):52-66. DOI:10.21608/TAEC.2017.11935
- Salama, F.M, Abd El-Ghani , M. M. Gaafar , A. E., Hasanin D. M. Abd El- Wahab D. A. (2021). Adaptive eco-physiological mechanisms of *Alhagi graecorum* in response to severe aridity in the Western desert of Egypt, *Plant Biosystems*. An International Journal Dealing with all Aspects of Plant Biology, <https://doi.org/10.1080/11263504.2021.1887957>
- Sayed, S.A., Gadallah, M.A.A. and Salama F.M. (2013). Ecophysiological studies on three desert plants growing in Wadi Natash, Eastern Desert, Egypt. *Journal of Biology and Earth Sciences*, 3(1):B135-B143.
- Sgarbossa, J., Schmidt, D., Schwerz, F., Schwerz, L., Prochnow, D. and Caron, B.O. (2019). Effect of season and irrigation on the chemical composition of *Aloysia triphylla* essential oil. *Revista Ceres*, 66(2):85-93. <https://dx.doi.org/10.1590/0034-737x201966020002>
- Shaltout, K.H., Al-Sodany, Y.M., Eid, E.M., [Heneidy](#), S.Z. and [Taher](#), M.A. (2020). Vegetation diversity along the altitudinal and environmental gradients in the main wadi beds in the mountainous region of South Sinai, Egypt. *Journal of Mountain Science*, 17(10). <https://doi.org/10.1007/s11629-020-6153-9>

- Sims DA, Gamon JA. 2002. Relationships between leaf pigment content and spectral reflectance across a wide range of species, leaf structures and developmental stages. *Remote Sens. Environment* 81, 337-354.
- Singwane, S.S., and Malinga, P.S. (2012). Impacts of Pine And Eucalyptus Forest Plantations on Soil Organic Matter Content in Swaziland - Case of Shiselweni Forests. *Journal of Sustainable Development in Africa*, 14(1):137-151. Corpus ID: 53406634.
- Smith JL, Doran JW. Measurement and use of pH and electrical conductivity for soil quality analysis. In *Methods for assessing soil quality*. Soil Science Society of America Special Publication. 1996;49:169-185. <https://doi.org/10.2136/sssaspecpub49.c10>
- Steel, R. G. D., & Torrie J. H. (1980). *Principles and procedures of statistics*. 2nd edition. McGraw Hill Book Company Inc., New York.
- Tounekti T, Mahdhi M, Al-Turki TA, Khemira H. Physiological responses of the halophyte *Salvadora persica* to the combined effect of salinity and flooding. *International Journal of Agriculture & Biology*. 2018;20(10): 2211–2220. DOI: 10.17957/IJAB/15.0764
- Treichel, S., Brinckmann, E., Scheitler, B., & von Willert, D. J. (1984). Occurrence and changes of proline content in plants in the southern Namib Desert in relations to increasing and decreasing drought. *Planta*, 162(3), 236–242. <https://doi.org/10.1007/BF00397445>
- Tuzuner, A. (1990). *Soil and water laboratory analysis guide*. Ankara: General Directorate of Rural Services Publications.
- Uvalle Saucedo JI, Gonzalez Rodriguez H, Ramirez Lozano RG, Silva IC, Gomez Meza MV. Seasonal trends of chlorophylls a and b and carotenoids in native trees and shrubs of Northeastern Mexico. *Journal of Biological Sciences*. 2008;8: 258-267. DOI: 10.3923/jbs.2008.258.267
- USDA. Natural Resources Conservation Service. *Soil Conservationists. Salinity Management Guide - Salt Management*. Available at <http://www.launionsweb.org/salinity.htm> 2002.
- Verbruggen N, Hua XJ, May M, VanMontagu M. 1996. Environmental and developmental signals modulate proline homeostasis: evidence for a negative transcriptional regulator. *Proc. Natl. Acad. Sci. USA* 93, 8787-8791.
- Wang, Z., Hu, H., Goertzen, L.R., McElroy, J.S. and Dane F. (2014). Analysis of the *Citrullus colocynthis* Transcriptome during Water Deficit Stress. *PLoS ONE*, 9(8):e104657. <https://doi.org/10.1371/journal.pone.0104657>
- Yinping, L., Tian, H., He, Q., Geng, Z., Yan, S., Tuniyazi, G. and Bai, X. (2021). Investigation of the geographical environment impact on the chemical components of *Peganum harmala* L. Through a Combined Analytical Method. *ACS Omega*, 6 (39):25497-25505. <https://doi.org/10.1021/acsomega.1c03420>

